

Constraining Neutrino Magnetic Moments at the FPF

Roshan Mammen Abraham

Department of Physics, Oklahoma State University

NuTau2021

September 29, 2021

Based on

- arXiv:2109.05032, A. Ismail, S. Jana, and R. Mammen Abraham
- In preparation, S. Foroughi-Abari, Y.-D. Tsai, and R. Mammen Abraham




OKLAHOMA STATE
UNIVERSITY

Neutrino Magnetic Moments

- Neutrino mixing parameters have been measured with incredible precision in recent years.
- **Electromagnetic properties of neutrinos can be used to distinguish Dirac and Majorana neutrinos**, and probe new physics.
- SM predicts a very small value for the active neutrino transition magnetic moments for Majorana neutrinos, $\sim 10^{-23} \mu_B$.
- The minimally-extended SM with right-handed neutrinos can acquire a diagonal magnetic moment proportional to their mass, $\mu_\nu^D \sim 10^{-19} \left(\frac{m_\nu}{1\text{eV}}\right) \mu_B$.
- These predictions are **several orders of magnitude smaller than the present experimental and astrophysical upper bounds** on the neutrino effective magnetic moments, motivating our study of NMM using LHC neutrinos.

Neutrino Magnetic Moments - Experimental Signature



A striking experimental signature of the magnetic moment operator is an electron recoiling. Incoming active neutrinos interact with the electrons in the target atom causing the electron to recoil. We consider two cases:

- Final state neutrino, ν_f is the same as the incoming active neutrino species, ν_i .
- ν_f is a Heavy Neutral Lepton (HNL) a.k.a sterile neutrino, N_R .

Neutrino Magnetic Moments - Cross-Section Expression

A characteristic feature of neutrino magnetic moment interaction is an enhancement in signal cross-section at low recoil energies,

$$d\sigma/dE_{rec} \sim 1/E_{rec}$$

For the scattering $\nu_\alpha e^- \rightarrow \nu_\alpha e^-$ we have,

$$\frac{d\sigma}{dE_{rec}} = \frac{\pi\alpha^2}{m_e^2} \left(\frac{1}{E_{rec}} - \frac{1}{E_\nu} \right) \left(\frac{\mu_{\nu_\alpha}}{\mu_B} \right)^2$$

and for $\nu_\alpha e^- \rightarrow N_R e^-$

$$\frac{d\sigma}{dE_{rec}} = \alpha (\mu_\nu^\alpha)^2 \left[\frac{1}{E_{rec}} - \frac{1}{E_\nu} + M_N^2 \frac{E_{rec} - 2E_\nu - M_e}{4E_\nu^2 E_{rec} M_e} + M_N^4 \frac{E_{rec} - M_e}{8E_\nu^2 E_{rec}^2 M_e^2} \right].$$

So if we have a **source of neutrinos and a detector with sufficiently low energy thresholds** then we can study neutrino magnetic moments. FPF!!!

Before we study the prospects at FPF we have to reduce the backgrounds.

- Muon-induced backgrounds: Muons can emit photons through bremsstrahlung which subsequently undergo pair conversion. If one of the resulting e^\pm is missed, the event would mimic our neutrino-electron scattering process. With timing, however, these events could be associated with the accompanying muon and vetoed.
- ν -induced backgrounds: The dominant background is ν interactions where only an e recoils. This is both NC interactions for all flavors, and CC interaction for ν_e only. We use differences in kinematic distributions to reduce this background.



At the FPF we focus on the following detectors:

- FASER ν 2 : 0.5 m x 0.5 m x 2 m tungsten detector with a mass of 10 tonnes, and $E_{th} = 300\text{MeV}$.
- FLArE : Liquid argon detector with $E_{th} = 30\text{MeV}$ and dimensions
 - 1 m x 1 m x 7 m with a mass of 10 tonnes.
 - 1.6 m x 1.6 m x 30 m with a mass of 100 tonnes.

With a lower energy threshold we expect FLArE to be more sensitive.

Active Neutrino Magnetic Moment

$$\nu_\alpha e^- \longrightarrow \nu_\alpha e^-$$



Left: The SM background has a flat distribution but the NMM contribution is enhanced at low recoil energies. **Right:** Expected number of SM and $\mu_{\nu_\tau} + \text{SM}$ events at FLArE10.

Active Neutrino Magnetic Moment

We employ a simple cut and count analysis with *cuts* corresponding to $E_{thresh} < E_{rec} < 1$ GeV.

Detector	SM backgrounds		$\mu_{\nu_e} = 5 \cdot 10^{-8} \mu_B$		$\mu_{\nu_\mu} = 5 \cdot 10^{-9} \mu_B$		$\mu_{\nu_\tau} = 5 \cdot 10^{-8} \mu_B$	
	no cuts	cuts	no cuts	cuts	no cuts	cuts	no cuts	cuts
FASTER ν 2	86	0.1	142	11.5	9.5	0.8	3.8	0.3
FLArE-10	51	0.1	97	24.1	7.3	1.9	4.0	1.0
FLArE-100	332	1.0	704	177.4	56.4	15.0	39.1	9.8



Bounds on μ_{ν_α} :

Detector	μ_{ν_e} ($10^{-8} \mu_B$)	μ_{ν_μ} ($10^{-9} \mu_B$)	μ_{ν_τ} ($10^{-7} \mu_B$)
FASTER ν 2	2.2	8.4	1.3
FLArE-10	1.5	5.5	0.75
FLArE-100	0.62	2.1	0.26

DONUT bounds are at $\mu_{\nu_\tau} < 3.9 \times 10^{-7} \mu_B$. FLArE-100 can do an order of magnitude better.

Active to Sterile Neutrino Transition Magnetic Moment

$$\nu_\alpha e^- \longrightarrow N_{R\alpha} e^- \quad ; \quad \mathcal{L}_{dipole} \supset \frac{1}{2} \mu_\nu^\alpha \bar{\nu}_L^\alpha \sigma^{\mu\nu} N_R F_{\mu\nu}$$



Qualitatively it is the same as before so we can employ a similar cut and count analysis. But **sterile neutrinos can undergo decays.**

Active to Sterile Neutrino Transition Magnetic Moment

The decay length of N_R in the lab frame is given by

$$l_{decay} = \frac{16\pi}{\mu_\nu^2 M_N^4} \sqrt{E_N^2 - M_N^2}, \text{ where } E_N = \text{energy of the outgoing } N_R.$$

l_{prompt} = minimum decay length for the decay vertex to appear displaced, and hence distinguishable from the production vertex.

- $l_{decay} > l_{detector}$: N_R decays outside the detector and the decay vertex is not observable.
- $l_{prompt} < l_{decay} < l_{detector}$: The decay vertex is sufficiently displaced from the production vertex and results in “double-bang” events.
- $l_{decay} < l_{prompt}$: The decay occurs promptly, leading to an electron and photon produced at the same point.

Active to Sterile Neutrino Transition Magnetic Moment

Of the possible signatures above, we focus on those with a single electron track emerging from the production vertex, **with no other nearby activity in the detector**.


We ignore events where the N_R decays promptly, which could have different backgrounds than the ones we consider.

We take l_{prompt} to be the mean free path λ for pair production by the photon in the detector material: $\lambda = 4.5$ mm (18 cm) for FASER ν 2 (FLArE).

Loose (strong) cuts correspond to $E_{thresh} < E_{rec} < 10$ (1) GeV.

Detector	SM backgrounds			$\mu_{\nu_e} = 10^{-7} \mu_B$			$\mu_{\nu_\mu} = 10^{-8} \mu_B$			$\mu_{\nu_\tau} = 10^{-7} \mu_B$		
	no cuts	loose	strong	no cuts	loose	strong	no cuts	loose	strong	no cuts	loose	strong
FASER ν 2	86	2.5	0.1	480	134.1	39	30	8.6	2.5	12.7	3.6	1.0
FLArE-10	51	2	0.1	320.5	144	79.6	22.3	10.4	5.9	13.1	5.9	3.3
FLArE-100	332	15	1.0	2285	1037	575.7	165.1	78.2	44.6	126.1	57.2	31.8

Active to Sterile Neutrino Transition Magnetic Moment




The colored dotted lines show $l_{decay} = l_{detector}$ for various detectors assuming $E_N = 100$ GeV, and the black dotted lines show $l_{decay} = \lambda$ in various detector materials. The red dashed line is the sensitivity at 90% C.L. coming from considering only double bang events at FLArE-10.

- The existence of nonzero neutrino magnetic moments is implied by neutrino masses.
- The need for BSM physics in the neutrino sector suggests the importance of searches for magnetic moments that could be larger than the typical expectation given the neutrino mass scale.
- The intense beam of ν 's, and detectors with low energy thresholds make FPF suited for such searches.
- FLArE-10 can constrain active tau neutrino magnetic moments to $\sim 10^{-7}\mu_B$, and FLArE-100 to $\sim 10^{-8}\mu_B$, and order of magnitude better than DONUT.
- Highly competitive bounds on active to sterile neutrino transition moments are also obtained at FPF.


Backup slides

Global picture: μ_{ν_e}



Backup slides

Global picture: $\mu_{\nu\mu}$



Backup slides

Global picture: $\mu_{\nu\tau}$

