



**IR2@EIC 2021-03-18**



# **Parton Propagation and Induced Dijet Production at EIC**

**Yuan-Yuan Zhang (CCNU)**

**Collaborator : Xin-Nian Wang (LBL)**


# Contents

- Introduction: Jet Transport Coefficient  $\hat{q}$  in cold nuclei
- Single Scattering Dijet & Double Scattering Dijet Cross Section
- Nuclear Modification of Dijet in e+A (  $\Delta\phi$ ,  $|y_{l_q} - y_l|$ ,  $R_A$  dependence )
- Summary and Outlook

# $\hat{q}$ in QGP and Cold Nuclei

-average squared pT broadening per unit length

Electron Ion Collision




$$\hat{q} \approx 0.015 \text{ GeV}^2/\text{fm}$$

P. Ru et al, arXiv:2004.00027


NB Chang et al, PRC **89.3** (2014): 034911

E Wang, XN Wang, PRL **89.16** (2002): 162301



JET Collaboration, PRC **90**, 014909

Heavy Ion Collision



$$\hat{q} \approx 1.2 \pm 0.3 \text{ GeV}^2/\text{fm} \quad \text{RHIC}$$

$$\hat{q} \approx 1.9 \pm 0.7 \text{ GeV}^2/\text{fm} \quad \text{LHC}$$

JET Collaboration, PRC **90**, 014909

# Existing efforts on $\hat{q}$ in cold nuclei

## High Twist Approach

- Use the pT broadening of final-state hadron (DIS) to extract  $\hat{q}$

$$\hat{q} \approx 0.015 \text{ GeV}^2/\text{fm}$$

P. Ru et al, arXiv:2004.00027

- Use the the suppression of leading hadron (DIS) to extract  $\hat{q}$

$$\hat{q} \approx 0.02 \text{ GeV}^2/\text{fm}$$

NB Chang et al, PRC **89.3** (2014): 034911


## Generalized High Twist Approach

Zhang, Y. Y., Qin, G. Y., & Wang, X. N. (2019). *PRD*, 100(7), 074031.

- $l_{\perp}, l_{q\perp} \sim k_{\perp}$ , no collinear expansion
- medium induced radiation spectra contain medium gluon TMD pdf  $\phi(x_G, k_{\perp})$  or TMD  $\hat{q}(k_{\perp})$


$$\hat{q} = \int d^2k_{\perp} \hat{q}(k_{\perp}) = C \int d^2k_{\perp} \rho \phi(x_g, k_{\perp})$$

# Dijet to probe gluon TMD pdf or TMD $\hat{q}(k_\perp)$



- Generalized High Twist Approach deals with hard splitting
- Further include initial quark transverse momentum  $v_\perp$

# Dijet in e+A: Single scattering



nuclear modification


$$\frac{d\hat{\sigma}_{eA}^S}{dx_B dQ^2 dz d^2l_\perp d^2l_{q\perp}} = \frac{2\pi\alpha_{em}^2}{Q^4} \sum_q e_q^2 \left[ 1 + \left( 1 - \frac{Q^2}{x_B S} \right)^2 \right] \frac{\alpha_s}{2\pi} \frac{1+z^2}{1-z} \frac{C_F}{\pi} \frac{q_A(x_B, \vec{v}_\perp)}{[\vec{l}_\perp - (1-z)\vec{v}_\perp]^2}$$

- multiple soft interaction (eikonalized as gauge link) → quark pT broadening
- pT broadening embedded in effective  $q_N(x, \vec{v}_\perp, \vec{b}_\perp)$ , gaussian broadening with width

$$\Delta_F(\vec{b}_\perp) = \int dy_0^- \hat{q}_F(y_0^-, \vec{b}_\perp) \quad \text{depend on } \hat{q} \quad \text{Liang, Z. T., Wang, X. N., \& Zhou, J. (2008). } PRD, 77(12), 125010.$$

# Dijet in e+A : Double scattering

Under two-parton correlation factorization



$$\frac{d\hat{\sigma}_{eA}^D}{dx_B dQ^2 dz d^2l_\perp d^2l_{q\perp}} = \frac{2\pi\alpha_{em}^2}{Q^4} \sum_q e_q^2 \left[1 + \left(1 - \frac{Q^2}{x_B S}\right)^2\right] \frac{\alpha_s}{2\pi} \frac{1+z^2}{1-z} \frac{2\pi\alpha_s}{N_c} \int \frac{d^2k_\perp}{(2\pi)^2} \int d^2b_\perp dy_0^- dy_1^-$$

$$\rho_A(y_0^-, \vec{b}_\perp) \rho_A(y_1^-, \vec{b}_\perp) q_N(x_B, \vec{v}_\perp, \vec{b}_\perp) \frac{\phi_N(x_G, \vec{k}_\perp)}{k_\perp^2} \left[ \mathcal{N}_g^{\text{nonLPM}} + \mathcal{N}_g^{\text{qLPM}} + \mathcal{N}_g^{\text{gLPM}} \right]$$


- initial quark  $q_N(x, \vec{v}_\perp, \vec{b}_\perp)$  contain pT broadening, depend on  $\hat{q}$  indirectly
- depend on  $\phi(x_G, k_\perp) \sim \hat{q}(k_\perp)$  directly

# Different contribution in double scattering

Contribution divided by how gluon radiated, understand from central cut diagrams


$$\mathcal{N}_g^{\text{nonLPM}} = C_F(\dots)$$

no LPM interference




$$\mathcal{N}_g^{\text{qLPM}} = \frac{1}{N_c}(\dots) \left[ 1 - \cos\left(\frac{y_1^- - y_0^-}{\tau_{gf}}\right) \right]$$

$$\tau_{qf} = \frac{2q^-z(1-z)}{[\vec{l}_\perp - (1-z)\vec{v}_\perp]^2}$$



$$\mathcal{N}_g^{\text{gLPM}} = C_A(\dots) \left[ 1 - \cos\left(\frac{y_1^- - y_0^-}{\tau_{gf}}\right) \right]$$

$$\tau_{gf} = \frac{2q^-z(1-z)}{[\vec{l}_\perp - (1-z)\vec{v}_\perp - \vec{k}_\perp]^2}$$





# A simple model for gluon saturation in $\phi(x_G, k_\perp) \sim \hat{q}(k_\perp)$


Simple model to include saturation

$$\phi_N(x_G, k_\perp, \mu^2) = \begin{cases} \phi_N^0 \text{ at } Q_s, & k_\perp < Q_s; \\ \phi_N^0(\mu^2 = k_\perp^2), & k_\perp > Q_s, \end{cases}$$

Calculate saturation scale self-consistently

$$Q_s^2(x_B, Q^2, b_\perp) = C \int dy_0^- \rho(y_0^-, b_\perp) \int \frac{d^2 k_\perp}{(2\pi)^2} \alpha_s(\mu) \phi_N(x_G, k_\perp, \mu^2)$$

Scale in  $\alpha_s(\mu)$  and  $\phi_N(x_G, k_\perp, \mu^2)$  is  $\mu^2 = k_\perp^2$



# Nuclear Modification Ratio

Ratio of dijet cross section in e+A and e+p

$$R_{eA}^{S(D)}(l_{\perp}, l_{q\perp}, \Delta\phi, z) = \frac{d\hat{\sigma}_{eA}^{S(D)}}{d\mathcal{P}} / A \frac{d\hat{\sigma}_{ep}}{d\mathcal{P}} \quad d\mathcal{P} \equiv dx_B dQ^2 dz d^2l_{\perp} d^2l_{q\perp}$$

Don't distinguish quark jet from gluon jet

$$\hat{\sigma} \equiv \hat{\sigma}(l_{\perp}, l_{q\perp}, \Delta\phi, z) + \hat{\sigma}(l_{q\perp}, l_{\perp}, \Delta\phi, 1 - z)$$

Kinematics for calculation:



$$E_e = 10 \text{ GeV}, E_N = 100 \text{ GeV}, x_B = 0.2, Q^2 = 200 \text{ GeV}^2, A = 208$$

Kinematic constraints from approximations, experiments


# Azimuthal angle $\Delta\phi$ dependence : $R_{eA}^D$

$$|\vec{l}_\perp + \vec{l}_{q\perp}| = 2l_\perp^2(1 + \cos \Delta\phi)$$


for  $l_\perp = l_{q\perp}$



nonLPM: finite(relative contribution increase with z increase)




qLPM: negligible(suppress by color and LPM factor)




gLPM: dominant

$R_{eA}^D$  magnitude decrease with increase of  $l_\perp(l_{q\perp})$ , double scattering contribution power suppressed by  $l_\perp(l_{q\perp})$

# Azimuthal angle $\Delta\phi$ dependence : $R_{eA}^S$ & $R_{eA}^{S+D}$



$$Q_s = 1.2 \text{ GeV}$$





$$Q_s = 2 \text{ GeV}$$

- Single scattering dominates in dijet Xsection,  $R_{eA}^S$  dominate in  $R_{eA}^{S+D}$
- $Q_s$  artificially increase, peak in  $R_{eA}^S$  moves,  $R_{eA}^D$  contribution increase

# Rapidity gap $|y_{l_q} - y_l|$ dependence

$$y_{l_q} - y_l = \ln\left(\frac{z}{1-z}\right) \text{ for } l_{\perp} = l_{q\perp}$$





○  $R_{eA}^S$  only due to quark pT broadening, independent of rapidity gap  $|y_{l_q} - y_l|$

○  $R_{eA}^D$  dominant - gLPM term with LPM suppression factor  $1 - \cos \frac{y_1^- - y_0^-}{\tau_{gf}}$



$|y_{l_q} - y_l| \uparrow, z \downarrow, \tau_{gf} \downarrow$ , when  $2R_A/\tau_{gf} \geq \pi$  LPM suppression disappear, increased incoherent contribution

# Nuclear size $R_A$ dependence



- $R_{eA}^S$  linear in  $R_A$
  - $R_{eA}^D$  nonLPM  $\propto \frac{1}{A} \int d^2b_{\perp} dy_0^{-} \rho_A(y_0^{-}, b_{\perp}) \int dy_1^{-} \rho_A(y_1^{-}, b_{\perp}) \sim \frac{9R_A}{16\pi r_0^3}$
  - $R_{eA}^D$  : qLPM , gLPM  $\propto \int_0^{R_A} dy_1^{-} \rho_A(y_1^{-}, 0_{\perp}) [1 - \cos \frac{y_1^{-}}{\tau_f}] = \frac{3}{4\pi r_0^3} R_A (1 - \frac{\sin(R_A/\tau_f)}{R_A/\tau_f})$
- Non-linear  $R_A$  dependence

# Summary




- Use dijet correlation in e+A to probe gluon TMD pdf  $\phi(x_G, k_{\perp})$  or TMD  $\hat{q}(k_{\perp})$
- Nuclear modification in single and double scattering depend on  $\phi(x_G, k_{\perp})$
- LPM effect and gluon saturation embedded in  $\phi(x_G, k_{\perp})$  bring unique features in  $\Delta\phi, |y_{l_q} - y_l|, R_A$  dependence of nuclear modification ratio

**Thanks for your attention!**




# Backup: Saturation scale $Q_s^2$

$$Q_s^2(x_B, Q^2, b_\perp) \approx Q_{s0}^2(x_B, Q^2) A^{1/3} \sqrt{1 - \frac{b_\perp^2}{R_A^2}}$$



$x_B = 0.1 - 1, Q^2 = 2 - 100 \text{ GeV}^2$



$x_B = 0.001 - 0.01, Q^2 = 2 - 100 \text{ GeV}^2$

can reach  $1 \text{ GeV}^2$  at  $x_B = 0.001, Q^2 = 100 \text{ GeV}^2$