

e+A Diffraction "Plan"

Mark D. Baker*

8 Sep 2021

Key tools, questions & critical path items


* - [mdbaker@ mdbpads.com](mailto:mdbaker@mdbpads.com), [bnl.gov](mailto:mdbaker@bnl.gov), [jlab.org](mailto:mdbaker@jlab.org)

Key plots to aim for

- "Foreground" plots – especially impact of beam effects.
- Reproduce background rejection for Pb @ IP6.
 - **Need all forward detectors simulated at some level.**
- Background rejection for Zr @ IP8
 - **Need all forward detectors simulated at some level.**
 - With Au or p 10σ for all RPs?
 - With the 10σ we need to get good A-1 tagging?

Sartre for Pb (Peter)

- Reproduce/understand classic J/ψ & ϕ plots from Au for Pb w/ simple 5% t smearing.
 - Extra smooth or realistic plots?
- Impact of perfect detector, but beam effects?
 - **Is the show already stopped?**
 - Do we have e & Pb p -spread included?
 - Alternatives to $d\sigma/dt$?




Toll, Ullrich,
PRC 87, 024913 (2013)

Sartre for Zr (Axel)

- Key question: coherent/incoherent vs. A.
Is Zr harder or easier in terms of background rejection needs?
- This seems on the critical path.

Background Rejection IR6 w/ Pb (Dhevan)

- Prove that we have all of the tools working.
Critical path. Uses full Pb J/ψ BeAGLE sample.



Chang et al.,
<https://arxiv.org/abs/2108.01694>

FIG. 7. Momentum transfer $|t|$ distribution in incoherent J/ψ production in ePb collisions at the EIC energy. Different lines indicate with or without beam pipe.

Back. Rejection IR8 w/ Pb & Zr (Dhevan)

- Uses full Pb & Zr J/ψ BeAGLE samples.
- Needs all forward detectors simulated at some level.
- Needs good understanding of 10σ for all RPs.
- Needs analysis code.
- **Very tight for end of September. Are we on track?**

Conclusions

- We have a lot to do and time is tight.
- Lots of big picture questions open:
 - Can we handle the beam effects?
 - Does IP8 help with Pb?
 - Is IP8 dramatically better for Zr due to A-1 tagging?
- What have I missed?