

Quark structure of pion from lattice QCD

CFNS Workshop: High Luminosity-EIC

June 23, 2022


Nikhil Karthik

Jefferson Lab - William & Mary



Understanding the internal structure of Nambu-Goldstone boson

- * String tension in pure-gauge theory, massive proton even in massless limit,...
- * Spontaneous Chiral symmetry breaking in QCD coupled to massless quarks

$$SU_L(N_f) \times SU_R(N_f) \times U_V(1)$$



$$SU_V(N_f) \times U_V(1)$$




- * $\Sigma = \langle \bar{\psi}_i \psi_i \rangle \neq 0$ Chiral-condensate


- * $N_f^2 - 1$ Massless Nambu-Goldstone bosons (“Pions”)

- * $\langle 0 | J_{5\mu}(0) | \pi \rangle = F_\pi p_\mu$ Decay constant

Valence PDF of $\pi^+(u\bar{d})$



JAM18, P. C. Barry et al, 2018




**No pion targets to perform DIS on.
PDF determination via indirect processes and assumptions**

Direct determination on the lattice

Factorization of form-factors: Distribution amplitude

Lepage, Brodsky '80


Efremov, Radyushkin '80



$$\langle P' | \mathcal{O}(Q) | P \rangle \sim \int_0^1 \int_0^1 dx dy \phi(x) E_{\mathcal{O}}(x, y) \phi(y)$$

EM current, stress tensor,...

Beyond what values of Q^2 do FFs approach their partonic description?



Projection for EIC
(20fb⁻¹, 5 GeV e⁻, 100 GeV p)

Aguilar et al, EPJA 55 (2019) 10, 190


What do we talk about when we talk about “x-dependent PDF from lattice”



I will restrict myself to one particular approach in this talk.

PDF as light-like separated $q\text{-}\bar{q}$ correlation

$$\mathcal{M}_{\text{LF}}(z^- P^+, \mu) = \frac{1}{2P^+} \langle H(P) | \bar{\psi}(z^-) W_+(z^-, 0) \gamma^+ \psi(0) | H(P) \rangle$$




$$f(x, \mu) = \int \frac{d\nu}{2\pi} e^{-ix\nu} \mathcal{M}_{\text{LF}}(\nu, \mu)$$


Light-Front physics from equal-time matrix elements

(conformal OPE for DA)
V. Braun, D. Muller '08

PDFs, GPDs, TMDs,..
X. Ji '13
J. Qiu, Y. Ma '14
A. Radyushkin '17



**Ultra-relativistic
Pion or any hadron H**



$$\mathcal{M}(zP_z, z^2) \stackrel{\text{Def}}{=} \langle H(\vec{P}) | \text{---}^z \text{---} | H(\vec{P}) \rangle$$

Twist expansion as the unifying concept

Distill leading-twist terms from Euclidean matrix elements at large-hadron momentum and/or short-distances

(Renormalized quantity from LQCD
e.g: ratios of qPDF matrix elements)

(IR-safe
Wilson coeff.)

($\overline{\text{MS}}$ PDF
moments)

$$\mathcal{M}(zP_z, z^2) = \sum_n c_n(z^2\mu^2) \frac{(-izP_z)^n}{n!} \langle x^n \rangle(\mu)$$

+ twist-2 trace $\text{tr} [M^k P_z^{n-k} z^n$

+ Higher twist $\sim \Lambda_{\text{QCD}}^{2k} P_z^{2n-2k} z^{2n}$

Suppress

***** $\langle x^n \rangle = \int_0^1 x^n f(x) dx$

****** “Twist” = [Dimension] – [Spin]

$$h(z, P_z) = \left\langle P_z \left| \begin{array}{c} \bar{\psi}(0) \\ \bullet \text{---} \text{---} \text{---} \bullet \\ W(0, z) \end{array} \right| P_z \right\rangle$$

Renormalize

$$\mathcal{M}(zP_z, z^2) = \frac{h(z, P_z)}{h(z, 0)}$$

Fourier Transform
at fixed P_z

$$q(x, P_z)$$

Twist-2 OPE $\sum_n \langle x^n \rangle c_n(\mu^2 z^2) \frac{(-iP_z z)^n}{n!}$

$$\int_{-1}^1 \frac{dy}{|y|} C\left(\frac{x}{y}, \frac{\mu}{|y|P_z}\right) f(y, \mu)$$

Compare in a range
of zP_z

$$f(x, \mu^2)$$

An experimental determination

$$\langle x^n \rangle$$

Access moments
directly

Fit Ansätze
 $x^\alpha (1-x)^\beta \mathcal{G}(x)$


$$f(x, \mu^2)$$

x-space
convolution

Valence PDF of pion: multiple perspectives on analysis

The basic object that lattice QCD measures:


Ratio of qPDF M.E.



(Variable that is Fourier conjugate to Bjorken x)


Comparing with experimental results without data processing

Pion matrix element
at momentum P_z





Extracting PDF data with minimal modeling: Mellin moments

Fit PDF moments as free parameters without modeling the x-dependence



(maximum of z-range used)






$$\mathcal{M}^{\text{tw}2}(zP_z, z^2) = \sum_n \frac{(-izP_z)^n}{n!} c_n(\mu^2 z^2) \langle x^n \rangle$$

Fit z & zP_z dependencies to model PDFs

$$f_v^\pi(x, \mu) = \mathcal{N} x^\alpha (1-x)^\beta (1 + s\sqrt{x} + tx)$$


Corresponding PDF



**Extracting x-dependent PDF
Using Ansätze**


Towards continuum limit

X. Gao, NK et al, PRD 102, 094513 (2020)
T. Izubuchi, NK et al, PRD 100, 034516 (2019)




- Two fine lattice spacing with a^2 corrections
- Systematic errors: range of quark-antiquark distance, ren. scheme, PDF ansatz..

Roadmap for precision lattice studies




Hybrid LaMET x-space matching



Recent work using exactly the same lattice data I showed in the last few slides, but analysis in x-space

Lattice meets experiments: universality




Lattice meets experiments: universality

Lattice data for pion matrix element (from BNL group)



Lattice data for proton matrix element (from ETMC)

Twist-2 OPE

Twist-2 OPE



JAM and ASV pion PDF data



**Towards exclusive processes at large- Q^2 :Distribution
amplitude of pion**

Pion distribution amplitude from leading-twist OPE

Theoretical framework:

V.Braun, D.Müller '07

X.Ji '13

A.Radyushkin '17

Y.S.Liu et al, '18

$$\langle 0 | \bar{u}(-\frac{z_3}{2}) \gamma_z \gamma_5 d(\frac{z_3}{2}) | \pi^+; P_3 \rangle = f_\pi P_3 \mathcal{M}(\lambda = z_3 P_3, z_3^2)$$

(equal-time quasi-DA matrix element)

$$\phi(x, \mu)$$

pert. matching in x-space

Mellin OPE at NLO

Conformal OPE at LL

$$\sum_{n=0} \frac{(-i\lambda)^n}{n!} \sum_{m=0}^n C_{n,m}(\mu^2 z^2) \langle x^m \rangle$$

Same up to
Finite NLO terms + truncation

$$\sum_{n=0} a_n \mathcal{F}_n \left(\frac{\lambda}{2}, \mu^2 z^2 \right)$$


Ratio of pion quasi-DA matrix elements and OPE fits

Renormalize: $\mathcal{M}(z_3 P_3, z_3^2, P_3^0) = \frac{h^B(z_3, P_3)}{h^B(z_3, P_3^0)}$

Fit OPE: $z \in [z_{\min}, z_{\max}]$


Lattice artifact?

Higher twist?




Estimating first two Mellin moments of pion DA

Variations in fit choices and corrections to twist-2



0.2866(62)(56)




0.138(28)(34)

Reconstruction of x-dependent pion DA

$$\phi(x, \mu) = \mathcal{N} x^\alpha (1-x)^\alpha \sum_n s_n C_n^{\frac{1}{2}+\alpha}(1-2x) \quad s_n \lesssim \delta$$


[See works by HadStruc collaboration
on using Jacobi polynomial basis]



Factorization of form-factors: lattice can play a part

$$F_{\pi}(Q^2) = \int_0^1 \int_0^1 dx dy \phi(x, Q) E(x, y, Q^2) \phi(y, Q)$$

Input fitted DA from lattice



Outlook

- There is both theoretical and experimental Interest in pion physics, especially from the perspective of EIC — PDF of pion, form factors at large Q^2
- The leading-twist perturbative framework applied to lattice matrix elements is a promising direction to do parton physics on the lattice. Comes with its own drawbacks — fit systematics, perturbative convergence, large P_z , small z . Progress underway in all these directions.
- Presented lattice results on valence PDF and DA for pion using quasi-operator using fine lattices.
- Is it essential to extract x -dependent PDF from twist-2 saturated lattice matrix elements? How to extract relevant experimental quantities in model independent manner (e.g., large- x behavior, convolutions that enter FF) ?
- What is the best way to make use of experimental results in lattice QCD Parton computations?
- The question on emergence of mass-gap and hadron-spin is general to Yang-Mills theories. How to develop relevant theoretical models that lattice can probe?