

SIDIS reconstruction with ML at the EIC

Connor Pecar

RHIC/AGS User Group Meeting

June 7, 2022

Research supported by:



U.S. DEPARTMENT OF
ENERGY


Office of
Science



Duke


SIDIS observables and coverage at the EIC

- Broad kinematics and PID coverage available at EIC/ATHENA
 - large lever arm for SIDIS multiplicities and asymmetries
- Many SIDIS projections made for proposals:
 - A_{LL} with kaons, gluon saturation with dihadrons, Sivers



ATHENA detector

- ATHENA (A Totally Hermetic Electron Nucleon Apparatus)
- Precise tracking and PID coverage in wide momentum range – great potential for SIDIS
- Studies shown in this presentation use ATHENA full simulation as Detector 1 full simulation is still in development



SIDIS kinematic reconstruction at EIC


- SIDIS variables: reliant on reconstruction of virtual photon four-momentum, typically determined using

$$q = l - l'$$

- Reliable for larger y , but begins to fail for $y < \sim 0.05$
 Low- y : region of interest for TMDs, and important for evolution studies

- **To utilize full EIC kinematic reach for SIDIS studies, need improved methods to determine SIDIS variables**


- CC – would require first method without electron



Phys.Rev. D70 (2004) 117504


ATHENA full simulation:

pT mean relative error, ele. method



Reconstruction with hadronic final state

- Through conservation of momentum and energy, **hadronic final state (HFS) should also contain enough information to constrain q**
- Methods utilizing hadronic final state should be more robust with respect to radiative corrections
 - Impact of radiative effects expected to be large for SIDIS observables at EIC, studies ongoing



Reconstruction with hadronic final state

- Method used in EIC YR and ATHENA proposal to reconstruct virtual photon using hadronic final state (HFS)
 - x and y components - summed HFS momentum
 - z and t components - solved for algebraically using

$$y = \frac{p \cdot q}{p \cdot l} \quad Q^2 = -q^2$$




and DIS variables from any DIS reconstruction method

i) <i>Leptonic variables</i>	$q \equiv q_l = k_2 - k_1, \quad y_l = p_1 \cdot (k_1 - k_2) / p_1 \cdot k_1$	
ii) <i>Hadronic variables</i> [81]	$q \equiv q_h = p_2 - p_1, \quad y_l = p_1 \cdot (p_2 - p_1) / p_1 \cdot k_1$	
iii) <i>Jacquet-Blondel variables</i> [82]	$Q_{JB}^2 = (\vec{p}_{2,\perp})^2 / (1 - y_{JB}), \quad y_{JB} = \Sigma / (2E(k_1))$ $\Sigma = \sum_h (E_h - p_{h,z})$	Prog. Part. Nucl. Phys. 2013, Blümlein
iv) <i>Mixed variables</i> [81]	$q = q_l, y_m = y_{JB}$	
v) <i>Double angle method</i> [83]	$Q_{DA}^2 = \frac{4E(k_2)^2 \cos^2(\theta(k_2)/2)}{\sin^2(\theta(k_2)/2) + \sin(\theta(k_2)/2) \cos(\theta(k_2)/2) \tan(\theta(p_2)/2)},$ $y_{DA} = 1 - \frac{\sin(\theta(k_2)/2)}{\sin(\theta(k_2)/2) + \cos(\theta(k_2)/2) \tan(\theta(p_2)/2)},$	

- + solution in quadratic equation found to always be closer to MC truth
- Resolution improved if this is carried out in head-on frame, then transformed to lab frame

ATHENA full simulation SIDIS resolution, p_T


Transverse momentum (w.r.t. q),
 10×275 , π^+ , $z > 0.2$:




DA, hybrid method ->
 p_T resolution more
 acceptable at low y

<- JB, HFS only
 Potential for CC



ATHENA full simulation SIDIS resolution, ϕ_h



ATHENA full simulation, 10x275,
pi+, z > 0.2



Angular resolution still
poor at low-y with all
methods




Machine learning reconstruction

- Based on hybrid HFS-electron SIDIS reconstruction, using ML to combine information from both to reconstruct q
 - Potential to correct overall HFS and electron momentum and to more accurately reconstruct z and t components than exact formula
 - ML models used for DIS reconstruction have been shown to be able to naturally account for radiative effects
 - (arXiv:2108.11638 Diefenthaler, Farhat, Verbytskyi, Xu, as well as NIM-A 1025 (2022) 166164, Arratia, Britzger, Long, Nachman)
- Currently utilizing graph-like neural network architectures designed for jet reconstruction
 - Treating DIS HFS at EIC similar to a jet

Particle flow networks

- Particle flow networks (PFN) developed by Komiske et al., (JHEP 01 (2019) 121, Komiske, Metodiev, Thaler)
 - Accepts unordered set of particles
 - Particles \rightarrow input to layers Φ
 - Summed over to create latent space of ℓ variables
 - Global features of event concatenated with latent space variables
 - Latent space variables and global features fed to layers F , produce final output
- Designed to be general purpose, universal approximator on sets of particles



JHEP 01 (2019) 121

ML SIDIS model and training



- Model combining electron and HFS:
 - Particle features for PFN: momentum, energy, η , φ in lab frame
 - Event-wide features: electron four momentum, DIS variables from JB, DA, electron methods
 - DIS variables will eventually be replaced with final reconstructed Q^2 and x likely using another ML method, but in this study statistics for training were limited
 - Target: MC virtual photon four-momentum in lab frame
- Training sample: ATHENA full simulation
 - HFS at the level of reconstructed particles
 - Version of dd4hep ATHENA full sim. used for detector proposal
 - Still some features missing, e.g. proper scattered electron ID
 - 10 GeV electron beam, 275 GeV proton beam, crossing angle -25 mrad
 - Trained on 3 million events with $Q^2 > 1 \text{ GeV}^2$, 2 million with $Q^2 > 10 \text{ GeV}^2$
 - 1 million $Q^2 > 1 \text{ GeV}^2$ events for validation

ATHENA full simulation,
10x275, pi+, z > 0.2



$$\frac{p_T - p_{T,true}}{p_{T,true}}$$

$$\phi_H - \phi_{H,true}$$

Electron
method



Neural
network



PFN able to
correct electron
method in almost
all of x-Q2







ATHENA full simulation,
10x275, pi+, $z > 0.2$

Mean:

- Comparison with other HFS/hybrid methods vs y_{true}
- NN clearly best performance for low y , and at least equaling electron method for large y

RMS:


Duke




ATHENA full simulation,
10x275, pi+, z > 0.2

- Comparison with other HFS methods vs true pT
- NN again clearly outperforming other methods for all pT, with no y cut

Mean:



RMS:



Summary

- Projections for the ATHENA and ECCE detectors demonstrate the exciting capabilities and kinematic coverage of the EIC for SIDIS measurements
- The electron method fails for $y < 0.05$, but can be improved using the hadronic final state and DIS variables to reconstruct virtual photon axis
- We demonstrate a machine learning approach combining the hadronic final state and scattered electron which surpasses existing methods for all of x - Q^2 and p_T
- Next steps in reconstruction:
 - Currently working on replacing the particle flow network with an architecture which can learn correlations between HFS particles (such as a GNN), as well as exploring other deep learning approaches
 - Method will need to be tested with better implementation of radiative effects

