The Climate Emergency: can Particle Physics ever be sustainable?

Véronique Boisvert (b. 329.7 CO2 ppm)

@VBoisvertRHULPP



7th March 2023 (421.4 CO2 ppm)

Outline

- The climate emergency
- CO2e emissions & solutions from:
 - Accelerators (construction/operation)
 - Detectors
 - Computing
 - Rest (travel, conferences, buildings, etc.)
- Possible recommendations
- Disclaimer:
 - I'm not a climate/energy scientist!
 - My research is on ATLAS, so energy frontier bias!




Climate Change: an emergency

IPCC AR6


Recent changes in the climate are widespread, rapid, and intensifying, and unprecedented in thousands of years.

IDCC WE UNDER CHANGE WHO UNDER



Unless there are immediate, rapid, and large-scale reductions in greenhouse gas emissions, limiting warming to 1.5°C will be beyond reach.






IPCC AR5

Climate Change: an emergency

Climate change is already affecting every inhabited region across the globe with human influence contributing to many observed changes in weather and climate extremes




IPCC AR6


Climate Change: an emergency


- UK parliament first to approve a motion to declare an "environment and climate emergency" on 1st May 2019
- Of the top 10 GHG emitters, only Japan, Canada and the EU have legally binding target of "net zero emissions by 2050 (2045)"
 - The pandemic was a blip (<u>lessons</u>)
- IPCC 2015 Paris agreement: aim to stay "below 2°C" so focus on 1.5 °C
 - NDC: Countries make pledges for how to achieve this (and then increase those pledges over time)
 - Climate Action Tracker: "With all target pledges, including those made in Glasgow, global greenhouse gas emissions in 2030 will still be around twice as high as necessary for the 1.5 °C limit.

IPCC AR6




Ice ages: ~ -5°C +4°C: civilization breakdown...







Emissions from accelerators: operations

- CERN now releases Environment reports (1st: 2017-18, 2nd: 2019-20)
- CERN peak power: ~180 MW (~ 1/3 of Geneva)
- Per year: ~ 1.2 TWh (~ 2% of Switzerland, o.o3% of Europe)
- LHC: ~55% of CERN's E consumption
- Electricity mainly comes from France: 88% carbon free (2017)




Electrical power distribution 2018






Emissions from accelerators: construction

- Potential future of energy frontier: <u>FCC</u> (ee then hh)
- Civil engineering:
 - Machine tunnel: one of the longest tunnels in the world: 97.75 km in circumference
 - 8 km of bypass tunnels
 - 18 shafts
 - 12 large caverns
 - 12 new surface sites
 - Excavation: 9 million cubic metres of spoil (mixture of marls and sandstone)
 - New roads






Emissions from accelerators: construction

- Concrete needed for the tunnel, which means (Portland) cement!
- Half of emissions from Portland clinker (<u>ref</u>)
- Ken Bloom and my rough calculation:
 - ~26ok tonnes of CO2 emissions
- Paper on emissions from road tunnels:
 - Lowest estimate: ~500k tonnes CO2 emissions
- Comparison: Using <u>report</u> for CO2e for construction of buildings: = building 8 London Shards!
- 1.4% of CH CO2e emissions (2016)
- Plant 6 million trees!





 $CaCO_3$ + heat \longrightarrow CaO + CO_2





Fig. 1. Schematic overview of the system boundary.

Emissions from accelerators: solutions

District heating:


- From 2022 hot water from LHC cooling at Point 8 will heat 8000 homes in Ferney-Voltaire, CERN also looking at Point 2 and 5, and Point 1 could heat CERN building on Meyrin site
- Since 2011 series of workshops: Energy for Sustainable Science at Research Infrastructures, 6th one: September 2022 at ESRF
 - Seems very Europe-centric
- Long-standing R&D in lowering accelerator power requirements
 - Eg Energy-Recovery in a Laser-Driven Plasma Wakefield Acceleration





Emissions from detectors

- Dominant CO2e emissions from CERN: gases used in experiments!
- Scope 1: direct emissions from organization/vehicles etc.
- Scope 2: indirect emissions from electricity generation, heating, etc.
- Scope 3: all other indirect emissions, upstream and downstream (business travel, personnel commutes, catering, etc.)




GROUP	GASES	tCO₂e 2017	tCO₂e 2018
PFC	CF ₄ , C ₂ F ₆ , C ₃ F ₈ , C ₄ F ₁₀ , C ₆ F ₁₄	61 984	69 611
HFC	CHF $_3$ (HFC-23), C $_2$ H $_2$ F $_4$ (HFC-134a), HFC-404a, HFC-407c, HFC-410a, HFC R-422D, HFC-507	106 812	96 624
	SF ₆	10 192	13 087
	CO ₂	14 612	12 778
TOTAL SCOPE 1		193 600	192 100


CERN environmental report 2017-18

Emissions from detectors

Particle detection


- SF6, HFCs and PFCs: particle detection
- HFCs and PFCs: detector cooling
- HFCs: air conditioning systems
- SF6: also used for electrical insulation in power supply systems





Gas recirculation is 90%

Emissions from detectors




Beatrice Mandelli, CERN Gas Systems

EU HFC phase-down policy

European Union "F-gas regulation":

- Limiting the total amount of the most important F-gases that can be sold in the EU from 2015 onwards and phasing them down in steps to one-fifth of 2014 sales in 2030.
- Banning the use of F-gases in many new types of equipment where less harmful alternatives are widely available.
- Preventing emissions of F-gases from existing equipment by requiring checks, proper servicing and recovery of the gases at the end of the equipment's life.




Prices could increase in EU and availability in the future is not known.

Reduction of the use of F-gases is fundamental for future particle detector applications


Beatrice Mandelli 4 9 Nov 2021


2020: CERN launched a working group on managing F-gases, with representatives
from the departments concerned and the large LHC experiments. The group looked at
issues such as the implementation of a centralised F-gas procurement policy, leak
detection, replacement alternatives, training courses for personnel handling F-gases,
and improving traceability and reporting.



- Alternative gas example: replace CF4
- CF4 prevents ageing, improves timing resolution and is a scintillator
- CMS CSC: currently 10% CF4
 - Reduce concentration to 5%
 - Replace with CF3l or HFO1234ze
- LHCb RICH studies:
 - CF4 or C4F10 used for good refractive index
 - Could replace C₄F₁₀ with C₄H₁₀ but flammable
 - Replace CF₄ with CO₂: under study
 - Use of SiPM to reduce the chromatic error and increase the yield







Why it is so difficult to find good GHG alternatives

When looking for alternatives eco-friendly gases, several factors have to be taken into account

Safety

Safety first for detector operations

- Gas mixture not flammable
- Gas components cannot have high toxicity levels



Safety

Performance



Tradeoff between flammability and GWP

- Replacing F with Cl or H: it shortens atmospheric lifetime BUT increase flammability limit
- Adding C=C bound: it increases reaction with O₂

Environment

GWP represents the main environment concern

Performance

GWP is related to IR absorption over time. Low GWP gases have short atmospheric lifetimes

- Water solubility -> rain out
- OH reactivity -> oxidation
- UV absorbance -> photolysis

RPC short and long term performance are affected

- Good quenching gases required
- Radiation-hard gas required
- Gases cannot heavily react with H₂O or UV radiation

FCC-ee CDR

- FCC-ee detectors:
- CLD (similar to CLIC): RPCs for the muon detector
- IDEA: large drift chamber (similar to KLOE and MEG2) containing He:iC4H10 90:10, iC4H10 GWP: ~3, for muon: large area μ-Rwell chambers, also use 10% iC4H10
- Crucial to do R&D in finding replacements or ensure 100% leak-free and 100% recirculation






Fig. 7.4. The CLD concept detector: end view cut through (left), longitudinal cross section of the top right quadrant (right).

G. Bencivenni et al 2017 JINST 12 C06027



Emissions from Computing

- Global IT sector <u>could be</u> 2-6% of global CO2e emissions, growing to 20% by 2030
- 70% from data centres and communication networks
- HEP uses Grid centres all over the world, yet emissions from electricity vary wildly





Solutions:

- Choose sites with green electricity...
- Green500 list
- Optimize your code ;-)

Far future (2040):

- All OECD electricity grids will be fully green...
- But huge demand for electricity

Emissions from FNAL

- In the US, DOE requirements to report yearly on environmental impacts including emissions
- **REC:** Renewable Energy Certificates

Scope 1 & 2 Greenhouse Gas Emissions
Goal: Reduce direct GHG emissions by 50 percent by FY 2025 relative to FY 2008 baseline
Interim Target (FY 2019): -31.0%

Current Performance: -62.5%

	FY 2008	FY 2019	% Change
Facility Energy	343,366.8	161,122.7	-53.1%
Non-Fleet V&E Fuel	142.6	186.6	30.9%
Fleet Fuel	691.6	0.0	-100.0%
Fugitive Emissions	40,165.1	139.1	-99.7%
On-Site Landfills	0.0	0.0	N/A%
On-Site WWT	0.0	0.0	N/A%
Renewables	0.0	0.0	N/A%
RECs	0.0	-17,435.4	N/A
Total (MtCO2e)	384,366.1	144,013.0	-62.5%


Scope 3 Greenhouse Gas Emissions Goal: Reduce indirect GHG emissions by 25 percent by FY 2025 relative to FY 2008 baseline Interim Target (FY 2019): -13.0%

Current Performance: -51.0%

	FY 2008	FY 2019	% Change
T&D Losses*	22,287.8	7,306.8	-67.2%
T&D RECs Credit	0.0	-1,148.5	N/A
Air Travel	2,215.8	2,530.1	14.2%
Ground Travel	168.9	128.5	-23.9%
Commute	4,633.3	5,392.5	16.4%
Off-Site MSW	191.8	247.7	29.1%
Off-Site WWT	4.8	11.0	129.2%
Total (MtCO2e)	29,502.4	14,468.1	-51.0%

^{*} Includes T&D losses for purchased renewable electricity

Emissions from FNAL




Emissions from Universities

- Heating/cooling + hot water + light/appliances = 40% of energy consumption of a UK citizen
- Not only helps Climate, but is cheaper to run
- B Corporations certification
- Green Labs: "research labs consume 10 times more energy and 4 times more water than office spaces", green lab certification, <u>LEAF</u>




Remaining carbon budget:

(50% chance of staying < 1.5°C)
460 GtCO2
Per year per person: 2.2t



Emissions from Travel

- Commuting, conferences, etc.
- A nearly carbon-neutral conference model
- Although aviation is 2.4% (2018) of global emissions (more than Australia or Italy or France!), rate of growth is large and carbon neutral flights long way off (CO₂ emissions increased by 32% from 2013-2018)
- Environmental groups calling for frequent flyers levy since eg in 2015 only 12% of people in England took 3 flights or more!
- Carbon offsetting as short term mitigation? <u>controversial</u>





One day in May 2019: 162,637 take offs recorded

Emissions from food

- IPCC report in August on Land Usage
- How about migrating our PP catering (meetings, conf, workshops) in that direction?

Carbonfootprint.com				
My Carbon footprint:	tonnes of CO2e		pesc.	vegan
House	0.92	9%	0.90	0.90
Flights	2.74	27%	1.37	1.37
Car	2.64	26%	0.96	0.96
Bus+Rail	0.07	1%	0.14	0.14
Food	1.58	15%	1.32	0.98
Secondary (clothes, etc.)	2.37	23%	2.38	1.19
Total	10.32		7.07	5.54
Target:	6.192		-	

Carbon Intensity of Eating: g CO2e/kcal				
Drinks	2.2			
Snacks, sugar	0.6			
Oils, spreads	0.8			
Fruit	4.6			
Vegetables	2.8			
Cereals, breads	1.3			
Dairy	4.5			
Chicken, fish, pork	3.8			
Beef, lamb	14.1			
Note: Figures are grams of carbon dioxide equivalents per kilocalorie of food eaten				

Note: Figures are grams of carbon dioxide equivalents per kilocalorie of food eaten (g CO2e/kcal). Intensities include emissions for total food supplied to provide each kilocarie consumed. This accounts for emissions from food eaten as well as consumer waste and supply chain losses. All figures are based on typcial food production in the USA. Estimates are emissions from cradle to point of sale, they do not include personal transport, home storage or cooking, or include any land use change emissions

Sources: ERS/USDA, LCA data, IO-LCA data, Weber & Matthews

Challet That Footnales

Pre-pandemic


Average UK	6.5	
Average EU	6.4	
Average world	5	
Target:	2	60%

My ATLAS footprint? Scope1+2 ~240 ktCo2e/4 = 60 ktCO2e/3k = 20 tCO2e

POSSIBLE RECOMMENDATIONS

European Strategy Update

 "The Strategy is due to be updated by May 2020 to guide the direction of the field to the mid-2020s and beyond."



Input to the ESU

VB (RHUL) Cham Ghag (UCL) Francesco Spano (RHUL) David Waters (UCL)




CERN Courier article

ECFA group of early career researchers asked to give input to European Strategy, document released which includes statements supporting the environmental sustainability of the field

Outcome

European Strategy Update web site



_Annex 3: The Working Groups

Working Group 1: Social and career aspects for the next generation

Chair: Professor Eric Laenen (Netherlands)

Working Group 2: Issues related to Global Projects hosted by CERN or funded

through CERN outside Europe

Chair: Professor Mark Thomson (United Kingdom)

Working Group 3: Relations with other groups and organisations

Chair: Professor Tatsuya Nakada (Switzerland)

Working Group 4: Knowledge and Technology Transfer

Chair: Professor Leander Litov (Bulgaria)

Working Group 5: Public engagement, Education and Communication

Chair: Professor Sijbrand de Jong (Netherlands)

Working Group 6: Sustainability and Environmental impact

Chair: Professor Dirk Ryckbosch (Belgium)

Outcome

7 3

Environmental and societal impact

- A. The energy efficiency of present and future accelerators, and of computing facilities, is and should remain an area requiring constant attention. Travel also represents an environmental challenge, due to the international nature of the field. The environmental impact of particle physics activities should continue to be carefully studied and minimised. A detailed plan for the minimisation of environmental impact and for the saving and re-use of energy should be part of the approval process for any major project. Alternatives to travel should be explored and encouraged.
- B. Particle physics, with its fundamental questions and technological innovations, attracts bright young minds. Their education and training are crucial for the needs of the field and of society at large. For early-career researchers to thrive, the particle physics community should place strong emphasis on their supervision and training. Additional measures should be taken in large collaborations to increase the recognition of individuals developing and maintaining experiments, computing and software. The particle physics community commits to placing the principles of equality, diversity and inclusion at the heart of all its activities.
- C. Particle physics has contributed to advances in many fields that have brought great benefits to society. Awareness of knowledge and technology transfer and the associated societal impact is important at all phases of particle physics projects. Particle physics research centres should promote knowledge and technology transfer and support their researchers in enabling it. The particle physics community should engage with industry to facilitate knowledge transfer and technological development.
- D. Exploring the fundamental properties of nature inspires and excites. It is part of the duty of researchers to share the excitement of scientific achievements with all stakeholders and the public. The concepts of the Standard Model, a wellestablished theory for elementary particles, are an integral part of culture. Public engagement, education and communication in particle physics should continue to be recognised as important components of the scientific activity and receive adequate support. Particle physicists should work with the broad community of scientists to intensify engagement between scientific disciplines. The particle physics community should work with educators and relevant authorities to explore the adoption of basic knowledge of elementary particles and their interactions in the regular school curriculum.

a) The energy efficiency of present and future accelerators, and of computing facilities, is and should remain an area requiring constant attention. Travel also represents an environmental challenge, due to the international nature of the field. The environmental impact of particle physics activities should continue to be carefully studied and minimised. A detailed plan for the minimisation of environmental impact and for the saving and re-use of energy should be part of the approval process for any major project. Alternatives to travel should be explored and encouraged.

SnowMass 2021

- Community Engagement Frontier: <u>TGo7</u>
 Environmental & Societal Impacts
- Coordinators: VB, Ken Bloom (U of Nebraska),
 Mike Headley (SDSTA-SURF)
- White papers released 15th March, Community meeting in Seattle in July
 - Summary report:
 - Now P5 needs to write their recommendations to HEPAP who then makes recommendations to DOE/NSF...

SnowMass2021

In Summary Report:

Finally, **HEP must take greater responsibility for its impacts on climate change** by addressing and mitigating these impacts through DOE project policies and individual community member actions.

Climate impacts of particle physics

Kenneth Bloom^{1,*}, Veronique Boisvert^{2,**}, Daniel Britzger³, Micah Buuck⁴, Astrid Eichhorn⁵, Michael Headley⁶, Kristin Lohwasser⁷, and Petra Merkel⁸

Abstract. The pursuit of particle physics requires a stable and prosperous society. Today, our society is increasingly threatened by global climate change. Human-influenced climate change has already impacted weather patterns, and global warming will only increase unless deep reductions in emissions of CO2 and other greenhouse gases are achieved. Current and future activities in particle physics need to be considered in this context, either on the moral ground that we have a responsibility to leave a habitable planet to future generations, or on the more practical ground that, because of their scale, particle physics projects and activities will be under scrutiny for their impact on the climate. In this white paper for the U.S. Particle Physics Community Planning Exercise ("Snowmass"), we examine several contexts in which the practice of particle physics has impacts on the climate. These include the construction of facilities, the design and operation of particle detectors, the use of large-scale computing, and the research activities of scientists. We offer recommendations on establishing climate-aware practices in particle physics, with the goal of reducing our impact on the climate. We invite members of the community to show their support for a sustainable particle physics field [1].

Submitted to the Proceedings of the US Community Study on the Future of Particle Physics (Snowmass 2021)

¹University of Nebraska-Lincoln, Lincoln, NE, USA

²Royal Holloway University London, United Kingdom

³Max-Planck-Institute for Physics, Munich, Germany

⁴SLAC National Accelerator Laboratory, Menlo Park, CA, USA

⁵CP3-Origins, University of Southern Denmark, Denmark

⁶Sanford Underground Research Facility (SURF), Lead, SD, USA

⁷University of Sheffield, United Kingdom


⁸Fermi National Accelerator Laboratory (Fermilab), Batavia, IL, USA

^{*}Contact author, e-mail: kenbloom@unl.edu

^{**}Contact author, e-mail: Veronique.Boisvert@rhul.ac.uk

Show support!

https://indico.fnal.gov/event/53795/



Recommendations

- New experiments and facility construction projects should report on their planned emissions and energy usage as part of their environmental assessment
 - Eg LHCb TDR for Phase II
- Review across all international laboratories to ascertain whether emissions are reported clearly and in a standardized way
- Take steps to mitigate impact on climate change by setting concrete reduction goals and defining pathways to reaching them
 - spend a portion of research time on directly tackling challenges related to climate change

Recommendations

- Minimize the travel emissions of users
- Long-term projects should consider the evolving social and economic context
- Actively engage in learning about the climate emergency and about the climate impact of particle-physics research
- Promote and publicize their actions surrounding the climate emergency to the general public and other scientific communities
- Engage with the broader international community to collectively reduce emissions

Recommendations

Eg ATLAS Sustainability Forum!

THE CLIMATE EMERGENCY: CAN PARTICLE PHYSICS EVER BE SUSTAINABLE?

DISCUSSION/QUESTIONS

BACK UP

Gas Recuperation systems at LHC experiments

Sometimes it is not possible to recirculate 100% of the gas mixture due to detector constrains

- Air **permeability**, max recirculation **fraction**, impurities, etc.
- A fraction of gas has to be renewed
 - Some gas is sent to the atmosphere
- This fraction of gas mixture can be sent to a recuperation plant where the GHG is extracted, stored and re-used
 - Challenges: R&D, custom development, operation and recuperated gas quality
- Gas recuperation also to empty/fill the detectors during LS

LHCb RICH1 and RICH2


During LS2 need to empty the detectors for maintenance/upgrade:

RICH1: 4 m³ of C₄F₁₀

Gas recuperation system for empty/filling and cleaning from Air contamination

RICH2: 100 m³ of CF₄

New gas recuperation system developed for the empty and filling of the detector



CMS CSC

Small Permeability to Air: accumulation of N₂

Need to inject fresh gas continuously to keep N₂ stable and guarantee detector performance

~80 l/h of CF₄ would be lost in exhaust without gas recuperation system


The R134a recuperation system for RPCs

ATLAS and CMS RPC Gas Systems


- Detector volume ~15 m³
- Gas mixture: ~95% C₂H₂F₄, ~5% iC₄H₁₀, 0.3% SF₆
- Gas recirculation: ~90%
 - Maximum recirculation validated for RPC detectors
- Fundamental to repair detector leaks
 - To have the gas at the exhaust of the gas system

R134a and iC₄H₁₀ form an azeotrope

A mixture of liquids whose proportions cannot be altered or changed by simple distillation



C₂H₂F₄ recuperation prototype system under study in CMS Experiment



Gas Chromatograph/Mass Spectrometer

Recuperation efficiency ~80%

First C₂H₂F₄ recuperation system under construction: installation foreseen beginning of 2023 in CMS experiment

Gas disposal

Abatement plants are employed when GHGs are polluted and therefore are not reusable

In case all studies on recuperation will not bring to efficient recuperation plants, industrial system able to destroy GHGs avoiding their emission into the atmosphere have been considered

Quite heavy infrastructure required:


- CH₄/city gas + O₂ supply + N₂ supply
- Waste water treatment
- PFC/HFC are converted in CO₂ + HF acid dissolved in water
- disposal of remaining waste/mud
 - To have the gas at the exhaust (600-1000 l/h)



Found also companies available to take PFC/HFC based mixture for disposal: but extremely expensive

Carbon footprint of PP researchers

- Flying!
 - 15% of E consumption for an average UK citizen
 - Particle Physicists fly a lot (esp. seniors)! Let's say, per year:
 - 8 European trips (eg use from London to Zurich): 8 x 148 Kg CO2: 1184 Kg CO2
 - 1 overseas trip (eg use from London to NYC): 986 Kg CO2
 - Total: 2170 Kg CO2: ~87 countries where the average citizen emits less CO2 in a year (incl. India, Morocco, Peru, Colombia)!
 - Using: <u>Guardian calculator</u>
 - A nearly carbon-neutral conference model




List of top CO₂ emitters


Forbes

	2018 CO2 Emissions	Global	Change Since
Country	in Billion Metric Tons	Share	Kyoto Protocol
China	9.43	27.8%	54.6%
U.S.	5.15	15.2%	-12.1%
India	2.48	7.3%	105.8%
Russia	1.55	4.6%	5.7%
Japan	1.15	3.4%	-10.1%
Germany	0.73	2.1%	-11.7%
South Korea	0.70	2.1%	34.1%
Iran	0.66	1.9%	57.7%
Saudi Arabia	0.57	1.7%	59.9%
Canada	0.55	1.6%	1.6%

Economicshelp.org



Emissions pathway



Sustainable HEP

28-30 June 2021 Zoom Europe/Zurich timezone

Effect your search

Q

Overview Timetable

Timetable

Call for Abstracts

Contribution List

Speaker List

Book of Abstracts

Participant List

Talk Recordings

ant recordings

Closing Statement

Closing Statement

Workshop "Sustainable HEP"

Closing Statement (status: 14th July 2021, 403 signatures)

On 28th—30th June 2021, the workshop "Sustainable High Energy Physics" took place by videoconferencing means with more than 350 registered participants from around 45 countries and five continents. The aim of this workshop was to initiate a community discussion on how to align the scientific operations within this particular subfield of physics with requirements of climate sustainability. Achieving the latter is a most pressing global issue for the present decade (as evidenced by the PPCC reports). The main focus of the workshop was on the scientific travel culture and the virtualisation of scientific exchange. The following topics were highlighted at this occasion:

- · characteristics of the climate crisis
- best practice examples on the virtualisation of scientific meetings
- challenges for research institutions to improve their climate sustainability
- · improvement of global inclusiveness in scientific exchange through virtualisation
- . domains of action for large scale experiments to improve their climate sustainability

We are organisers and participants of the workshop as well as members of the High Energy Physics community or related fields of physics. We understand that the climate impact of certain aspects of our field of research is a cause of concern and we assert that there is a need for determined action to align these with the goals of the Paris climate agreement and, more generally, with the needs of a sustainable society. Our aim is to trigger a discussion on how HEP can live up to its responsibility in the global transition to a sustainable and climate neutral world, while maintaining the high quality of research and international scientific exchange. In this context, we highlight increased inclusiveness as a crucial co-benefit of online formats.

We thus encourage members of our community to discuss and enable suitable implementations of sustainable development for our field. We stress that this is a call to develop a balanced and deliberated approach that brings together the needs of a global HEP community with the needs of climate sustainability. We call on research and funding institutions to adjust the general framework for research accordingly and to facilitate a transformation towards sustainable means. Consequently, we invite the formation of working groups to continue the discussions initiated at the workshop and to conduct further installations of the workshop on related topics of sustainability that deserve discussions in a broader setting.

Signatures

The following persons have signed the statement as individuals on their own behalf. Please note that institutions are mentioned merely to identify the signatories' current scientific affiliations. This statement does not (necessarily) reflect the opinions of these institutions.


sign here

workshop organisers:

Niklas Beisert (ETH Zürich)
Valerie Domcke (CERN/EPFL)
Astrid Eichhorn (CP3-0rigins, University of Southern Denmark)
Kal Schelmit (CFEN)

workshop participants:

Also: white paper for Australian
Astronomy: <u>"The imperative to reduce carbon emissions in astronomy"</u>



Source: Table 1.2, Final UK greenhouse gas emissions national statistics 1990-2019 Excel data tables Note: LULUCF is land use, land use change and forestry.

World Emissions Clock