# Parameter Exploration for Realistic Seeding

By Emma Yeats

#### Introduction/Recap

Continuing Rey's recent study on parameter optimization for the realistic seeder.

Two main results are discussed:

- a quick update on the bug found in ElCrecon
- my studies on varying parameters for realistic seeding and how they were changed by the bug

I will also show the updated optimal parameters.

All plots from the same sample: 10k single pions thrown with  $p \in [0,10]$  GeV/c and  $\eta \in [-4,4]$  (Brycecanyon, ACTS 21.1)

#### **Update on EICRecon Bug**

After a bug was found by Barak and Rey, EICRecon recently updated the pt calculation:



# Parameter Exploration

## Parameter Descriptions

Parameter	Description	ElCrecon default	Y.S. Lai's default	
bFieldInZ	z component of magnetic field	1.7 T	1.7 T	
rMax	Maximum r value to look for seeds	500 mm	440 mm	
rMin	Minimum r value to look for seeds	33 mm	33 mm	
zMin	Minimum z value to look for seeds	-800 mm	-1500 mm	
zMax	Maximum z value to look for seeds	800 mm	1700 mm	
beamPosX	Beam offset in x	0	0	
beamPosY	Beam offset in y	0	0	Parameters
deltaRMinTopSP	Min distance in r between middle and top SP in one seed	1 mm	50 mm	fixed based on
deltaRMinBottomSP	Min distance in r between middle and bottom SP in one seed	1 mm	50 mm	the geometry
deltaRMaxTopSP	Max distance in r between middle and top SP in one seed	400 mm	220 mm	the geometry
deltaRMaxBottomSP	Max distance in r between middle and top SP in one seed	400 mm	220 mm	
collisionRegionMin	Min z for primary vertex	-300 mm	-250 mm	
collisionRegionMax	Max z for primary vertex	300 mm	250 mm	J
cotThetaMax	Cotangent of max theta angle	16	16.54	
minPt	Min transverse momentum	100	100 MeV/cotThetaMax	
maxSeedsPerSpM	Max number of seeds a single middle space point can belong to - 1	1	0	
sigmaScattering	How many standard devs of scattering angles to consider	5	5	
radLengthPerSeed	Average radiation lengths of material on the length of a seed	0.1	0.1	
impactMax	Max transverse PCA allowed	20 mm	3 mm	
rMinMiddle	Min R for middle space point	20 mm	-	
rMaxMiddle	Max R for middle space point	400 mm	—	
bFieldMin	min B field	-	0.1 T	5

## Parameter Descriptions

Parameter	Description	ElCrecon default	Y.S. Lai's default	
bFieldInZ	z component of magnetic field	1.7 T	1.7 T	
rMax	Maximum r value to look for seeds	500 mm	440 mm	
rMin	Minimum r value to look for seeds	33 mm	33 mm	
zMin	Minimum z value to look for seeds	-800 mm	-1500 mm	
zMax	Maximum z value to look for seeds	800 mm	1700 mm	
beamPosX	Beam offset in x	0	0	
beamPosY	Beam offset in y	0	0	
deltaRMinTopSP	Min distance in r between middle and top SP in one seed	1 mm	50 mm	1
deltaRMinBottomSP	Min distance in r between middle and bottom SP in one seed	1 mm	50 mm	
deltaRMaxTopSP	Max distance in r between middle and top SP in one seed	400 mm	220 mm	
deltaRMaxBottomSP	Max distance in r between middle and top SP in one seed	400 mm	220 mm	
collisionRegionMin	Min z for primary vertex	-300 mm	-250 mm	Parameters
collisionRegionMax	Max z for primary vertex	300 mm	250 mm	Lstudiod
cotThetaMax	Cotangent of max theta angle	16	16.54	i studied
minPt	Min transverse momentum	100	100 MeV/cotThetaMax	
maxSeedsPerSpM	Max number of seeds a single middle space point can belong to - 1	1	0	
sigmaScattering	How many standard devs of scattering angles to consider	5	5	
radLengthPerSeed	Average radiation lengths of material on the length of a seed	0.1	0.1	
impactMax	Max transverse PCA allowed	20 mm	3 mm	
rMinMiddle	Min R for middle space point	20 mm	-	
rMaxMiddle	Max R for middle space point	400 mm		
bFieldMin	min B field	<u> </u>	0.1 T	6

#### Parameters Chosen For Discussion

Before this bug, I found one parameter in particular that was responsible for increasing the efficiency in the far eta regime (radLengthPerSeed). I will discuss it's behavior briefly just to explain the sudden efficiency jump you'll see in that region.

Several of the parameters' behavior did not change after the bug was fixed, like deltaRMinSP, deltaRMaxSP, cotThetaMax, and more. However some changed after the fix, including radLengthPerSeed and sigmaScattering, that once had a larger impact on the efficiency and now do not.

### radLengthPerSeed (pre-Bug)





#### Average radiation lengths of material on the length of a seed.

I initially varied this parameter from 0.01% to 1%. The number of reconstructed seeds and the efficiency both increase in the far eta regimes.

#### radLengthPerSeed (pre-Bug)





Above was calculated before the bug was fixed, to show where the sudden increases in efficiency come from. It is less sensitive post-Bug, and I found that this increase in efficiency now happens around ~1e-16.

#### radLengthPerSeed (post-Bug)

(we can see it no longer really has an effect, and we can revert to the default, radLengthPerSeed==0.1)



#### Now we look at the parameters that remained sensitive.

#### DeltaRMinTop/BottomSP (post-Bug)



We see that 10 mm is ideal, and also doesn't increase the number of reconstructed seeds too much.

#### deltaRMaxTop/BottomSP (post-Bug)



Maximum distance in r between Middle and Top/Bottom SP in one seed.

We see this should be set  $\geq$ 200 mm.



#### cotThetaMax (post-Bug)

3.63

10.02

16.54

27.29

15.4

5.70

3.46

2.10

2 3

4

3.5



27.29 is best, but I am still working to see why the full eta regime is not covered (future work). 100

50

-4 ... -2 .... 0 ... 2 ....

#### maxSeedsPerSpM (post-Bug)





The max number of seeds a single middle spacepoint can belong to (-1)

Efficiency is consistent, but the number of reconstructed seeds doubles after 0.

### **Updated Parameters**

Parameter	Description	ElCrecon default	Y.S. Lai's default	My New Default
bFieldInZ	z component of magnetic field	1.7 T	1.7 T	1.7 T
rMax	Maximum r value to look for seeds	500 mm	440 mm	440 mm
rMin	Minimum r value to look for seeds	33 mm	33 mm	33 mm
zMin	Minimum z value to look for seeds	-800 mm	-1500 mm	-1500 mm
zMax	Maximum z value to look for seeds	800 mm	1700 mm	1700 mm
beamPosX	Beam offset in x	0	0	0
beamPosY	Beam offset in y	0	0	0
deltaRMinTopSP	Min distance in r between middle and top SP in one seed	1 mm	50 mm	10 mm
leltaRMinBottomSP	Min distance in r between middle and bottom SP in one seed	1 mm	50 mm	10 mm
deltaRMaxTopSP	Max distance in r between middle and top SP in one seed	400 mm	220 mm	200 mm
leltaRMaxBottomSP	Max distance in r between middle and top SP in one seed	400 mm	220 mm	200 mm
collisionRegionMin	Min z for primary vertex	-300 mm	-250 mm	-250 mm
collisionRegionMax	Max z for primary vertex	300 mm	250 mm	250 mm
cotThetaMax	Cotangent of max theta angle	16	16.54	27.29
minPt	Min transverse momentum	100	100 MeV/cotThetaMax	100 MeV/cotThetaMax
maxSeedsPerSpM	Max number of seeds a single middle space point can belong to - 1	1	0	0
sigmaScattering	How many standard devs of scattering angles to consider	5	5	5
radLengthPerSeed	Average radiation lengths of material on the length of a seed	0.1	0.1	0.1 (mm)
impactMax	Max transverse PCA allowed	20 mm	3 mm	3 mm
rMinMiddle	Min R for middle space point	20 mm	-	-
rMaxMiddle	Max R for middle space point	400 mm	-	-
bFieldMin	min B field	-	0.1 T	0.1

#### Summary + Future Work

-Study produced an overall efficient setup! There are a few more parameters I can look at, especially after the recent units change in the pt calculation. But fixing the bug itself increased the efficiency by a ton.

#### -Including other resolution plots like d $\Theta$ , d $\Phi$ , vertexing position.







#### Momentum Resolutions in Greater Detail



### Figuring Out the Units

Pt is calculated by equating lorentz and centripetal forces.

$$m\frac{v^2}{R} = q_e \left( v \times B \right) \longrightarrow p[GeV/c] = 0.3 * R[m] * B[T]$$

#### This is where pt is calculated in EICRecon - two bugs here!



#### minPt



Parameter Values: 1,5,6,8,10 (default ==5). No effect on efficiency or #reconstructed seeds in the region varied, could increase values further move to backup

#### impactMax



#### bFieldMin



values=0,0.05,0.1,0.5,1,1.5,1.7 T ... seemed to have no effect at all on efficiency or reconstructed seeds.

#### Number of Missing Values

Number of missing values - the number of non-null entries in the generated branches can be larger than that in the reconstructed ones. These are cases in which no seed was reconstructed. Let's visualize this as the percentage of missing values



#### Varying rad length per seed after bField change



25







How many standard deviations in scattering angle are considered?

Overall we see that efficiency increases with increasing Sigma, with no effect on the number of reconstructed seeds.

This is important to keep in mind once a standard deviation is chosen for the detector later, that I have just kept my default at  $5\sigma$ . Efficiency will decrease if a smaller  $\sigma$  is chosen.

#### sigmaScattering

#### No longer influences efficiency much post-Bug



#### radLengthPerSeed (post-Bug)

