



RHIC Status

Kiel Hock

Last Week at RHIC



RHIC Status

- 111x111 physics running since 10/7
- STAR at 945 M minbias events. Goal is 1-2 B events.
- Various bunch configurations for sPHENIX detector commissioning (1x1, 6x6, 12x12, 56x56, 111x111)
- Vertical and longitudinal cooling enabled in blue and yellow.
- MVTX studies with single bunches in blue and/or yellow taking place post absorber install.
 - ▶ Reported MVTX background Wednesday. Decided to go ahead with absorber install Thursday.
 - ▶ Single beam studies in yellow began Friday.
 - ▶ Showed sensitivity to beam's momentum distribution and local orbit steering (via crossing angle).
 - ▶ Optimal configuration in yellow is $dR = -0.5$ mm, -8 mm at $yo8-dh14$, and a 1 mrad crossing angle.
 - ▶ Single beam study in blue yesterday showed the optimal configuration for yellow is also supported by blue.
 - ▶ Will start increasing the number of bunches to determine how it scales with bunch number.
 - ▶ Several weekly meetings since Friday to discuss the status of studies.
- Blue injection efficiency lower than yellow, being investigated.

Background Issues

sPHENIX

- Simulations show this can be caused by a single particle with initial trajectory along the longitudinal axis and inplane with the MVTX.
- Simulations show particles with high momentum amplitude could strike at the taper nearest the MVTX.
- Using a change in radius in the arcs in combination with a bump at yo8-th14 (incoming yellow side) the high amplitude particles. Change of angle from 2 to 1 mrad helps with distributing remaining losses.
- Need to reoptimize collisions and collimation.

STAR

- Au78 produced at sPHENIX gets deposited at STAR
- Likely will be helped with the radius change. Can look to install a bump exiting sPHENIX to see if it improves.

Additional studies

- gap cleaning off-tune
- squeeze of β^* at IP10
- unsqueeze of β^* at IP8
- octupoles
- 56 MHz
- revisit use of secondary collimation
- low intensity with heavy cooling