


Vetoing Efficiency of Incoherent Diffractive Vector Meson Production at the Second Interaction Region at the EIC

Jihee Kim (jkim11@bnl.gov)

2024/07/22

Lehigh University, Bethlehem, PA, USA


EIC 2nd Detector Motivation



- EIC Design
 - Two interaction points (IP-6 and IP-8)
 - Two interaction regions (IR-6 and IR-8)
- **Detector 1**, called **ePIC**, locates at **IP-6**
- Favors to accommodate the **second detector** and a **second interaction region**
- **A general-purpose collider detector to support full EIC program (complementarity)**
 - **Cross-checks & control of systematics**
 - Subdetector technologies
 - Magnetic field
 - **Broaden physics program (different physics focuses)**

EIC Interaction Regions

Requires specialized detectors integrated in the interaction region over 80 m



Crossing angle: **25 mrad**

IR-Design:
 $0.2 \text{ GeV} < p_T < 1.3 \text{ GeV}$


Same
Accelerator highlights
and challenges

Shared
Luminosity at both IRs

Center-of-mass energy
coverage

Different
Blind spots

Far-forward detector
acceptances




Crossing angle: **35 mrad**

IR-Design:
2nd “beam optics” focus
comes with challenges (ex. magnet
design) in accelerator machine


IR Concept – 2nd Focus in Far-Forward

- By adding additional magnet to focus beam ~ 45 m downstream from interaction point
- This is NOT detector design, but it is **machine design that detector can be benefit from**
- 2nd focus enables
 - Higher probability to **detect low p_T (< 250 MeV) particles**
 - Detects near-beam particles where get out beam envelop
- **Complementary to ePIC: exclusive, tagging, and diffractive physics analysis**




Physics Opportunities with 2nd Focus

- 2nd focus in IR8 greatly improves **forward acceptance**
- Complementarity with Detector 1 (ePIC) @ IR-6
- **Excellent low- p_T acceptance** for protons and light nuclei from exclusive reactions **at very low t**
- **Detection of target fragments** makes it possible
 - To veto breakup to study coherent process
 - To study final state when breakup occurs
- Coherent diffraction on heavy nuclei by vetoing breakups
- Rare isotopes detection and identification of heavy fragments



Physics Opportunities with 2nd Focus


- 2nd focus in IR8 greatly improves **forward acceptance**
- Complementarity with Detector 1 (ePIC) @ IR-6
- **Excellent low- p_T acceptance** for protons and light nuclei from exclusive reactions **at very low t**
- **Detection of target fragments** makes it possible
 - **To veto breakup to study coherent process**
 - To study final state when breakup occurs
- Coherent diffraction on heavy nuclei by vetoing breakups
- Rare isotopes detection and identification of heavy fragments
- **Today, show study on capabilities of separating coherent from incoherent diffractive events by tagging far-forward nuclear fragments**



Far-Forward Detector – Layout

Implemented in **proposed IR-8 Forward Hadron Lattice** and IR-6 detector configuration


pre-conceptual design




Subdetector	Detection
B0 spectrometer	Photons + charged particles
Off momentum detector	Charged particles
Zero degree calorimeter	Neutrons + photons
Roman Pot	Charged particles + Nuclear fragments

Far-Forward Detector – Acceptance


Zero Degree Calorimeter



- ZDC < 5 mrad uniformly ~100 %
- Exit window material impact on neutron acceptance



Roman Pot (@ 2nd focus)



ep DVCS
18x275 GeV²

- Windows on pots depending on the **beam optics (transverse beam size)**

$$\sigma_{x,y} = \sqrt{\epsilon_{x,y} \beta(z)_{x,y} + \left(D_{x,y} \frac{\Delta p}{p}\right)^2}$$

where
 ϵ : Emittance at z=0
 β : Beta function at z=RPSF
 D : Momentum dispersion at z=RPSF
 $\frac{\Delta p}{p}$: Momentum spread at z=0

© RPSF: Roman Pot at Secondary Focus





Approach – Vetoing Efficiency

- To understand **impact of 2nd focus on suppression of incoherent contribution** (background to coherent J/ψ production)
- Used **BeAGLE v1.03.02** ePb $18 \times 110 \text{ GeV}^2$ J/ψ production ($1 < Q^2 < 10$)
Incoherent events $ePb \rightarrow e' + J/\psi(ee/\mu\mu) + X$
- Applied **10σ safe distance cut** based on **eAu @ IR-8 Roman Pot at 2nd focus**
- **Tagged events for nuclear breakups tagging purpose**
 - ZDC Hcal: **any registered RAW hits**
 - RPSF: **one layer (closest to 2nd focus)** has registered RAW hits outside **10σ** safe distance
 - OMD: **two layers** (actual four layers as redundancy) have registered RAW hits
 - B0 Tracker: **at least two out of four layers** have registered RAW hits
 - B0 Ecal: **energy** of all hits greater than **100 MeV**
 - ZDC Ecal: **energy** of all hits greater than **100 MeV**

Nuclear Breakups Distribution


Nuclear Breakups at Final State	Number of Events
Only Neutrons	7.86 %
Only Protons	0.0001 %
Only Photons	3.45 %
Neutrons + Protons	3.18 %
Neutrons + Photons	45.41 %
Protons + Photons	1.85 %
Neutrons + Protons + Photons	38.25 %

About **95 %** of events have **neutrons**



Far-Forward Detector – Incoherent Veto


Reference from EIC YR p.352



At position of third diffractive minimum,
**rejection factor for incoherent event
better than 400:1 must be achievable
(0.0025 % inefficiency)**


Far-Forward Detector – Incoherent Veto

Reference from EIC YR p.352



At position of third diffractive minimum, rejection factor for incoherent event better than 400:1 must be achievable (0.0025 % inefficiency)

Veto inefficiency for incoherent events




- ZDC hcal tagged (neutrons)
- RPSF tagged (protons, nuclear fragments)
- OMD tagged (charged particles)
- B0 tracker tagged (charged particles)
- B0 ecal tagged (photons)
- ZDC ecal tagged (photons)

← Vetoing power > 10³

Found to be enough to suppress incoherent contribution at three minima
Vetoing efficiency is >> 99.99%

Simplified Beam Pipe Implementation


Proposed pre-conceptual design of IR-8 Forward Hadron Lattice




Detector Acceptance Comparison: ZDC

Single Neutron $E = 275$ GeV and $0 < \theta_{MC} < 10$ mrad


Generated



Accepted
without beam pipe/exit window



Accepted
with beam pipe/exit window




In total, about **99.96 % (97.2 % with beam pipe/exit window)** events were accepted.

Full acceptance in θ_{MC} changed from up to 5 mrad to 4.5 mrad


Detector Acceptance Comparison: RPSF

Single Proton $E = 275$ GeV and $0 < \theta_{MC} < 5$ mrad


Generated



Accepted
without beam pipe



Accepted
with beam pipe



In total, about **95.3 % (91.7 % with beam pipe)** events were accepted.
However, **full acceptance up to 4.2 mrad stays the same**

Vetoing Efficiency Comparison

Veto Selections	Surviving Events	
	Before Beam Pipe/Exit Window	After Beam Pipe/Exit Window
All events	997,820	998,161
Events with one scattered electron identified and $ \eta_{J/\psi} < 4$ and $1 < Q^2 < 10$	732,455 (100 %)	732,707 (100 %)
ZDC HCAL tagged	41,848 (5.71339 %)	42,476 (5.79713 %)
+ RPSF tagged	71 (0.00969343 %)	66 (0.00900769 %)
+ OMD tagged	71 (0.00969343 %)	64 (0.00873473 %)
+ B0 tracker tagged	30 (0.00409581 %)	30 (0.00409441 %)
+ B0 ecal tagged	17 (0.00232096 %)	19 (0.00259312 %)
+ ZDC ECAL tagged	4 (0.000546109 %)	10 (0.0013648 %)

Summary and Outlook

- To explore physics opportunities by taking full advantage of 2nd focus
 - Implemented **pre-conceptual design of IR-8 hadron beamline geometry** and its field configuration + **required far-forward detectors**
- Using BeAGLE incoherent events (ePb 18×110 GeV² J/ψ production), evaluated vetoing power by tagging nuclear fragments using far-forward detectors
 - **Found to be enough to suppress incoherent contribution at three diffractive minima**
 - Shown that it can **achieve vetoing efficiency of $\gg 99.99$ %** (i.e. vetoing power $\sim 10^3$)
- Possible physics cases with improved forward acceptance, in particular of low p_T
 - **Access low t** where one can probe large impact parameter b (pion cloud)
→ change of transverse spatial distribution of gluons or quarks with x allows to help us to **better understand mechanism of confinement**
 - Provide possibility for **constraining diffractive longitudinal structure function (F_L^D)**;
Reggeon and Pomeron → enables to **study two Reggeon and Pomeron contributions at the same machine** and may opens new opportunity to study separate from one contribution to another

Backup Slides

IP-8 Beam Parameters and 10σ Cut

From EIC CDR table 3.5 and Randy's eAu study

© RPSF: Roman Pot at Secondary Focus

eAu 18 GeV on 110 GeV	Momentum Dispersion ($D^{\text{secondary focus}}$)	Emittance X (ϵ_x^*) [mm]	Emittance Y (ϵ_y^*) [mm]	Beta function X ($\beta_x^{\text{secondary focus}}$) [mm]	Beta function Y ($\beta_y^{\text{secondary focus}}$) [mm]	Momentum spread ($\Delta p/p$)*
Old ep 18 on 275 GeV ²	0.382	43.2e-6	5.8e-6	2289.454596	4538.713168	6.2e-4
New ep 18 on 275 GeV ²	0.465446718	43.2e-6	5.8e-6	498.013008	3392.376638	6.2e-4
New eAu 18 on 110 GeV ²	0.467582853	43.2e-6	5.8e-6	565.292559	1870.555797	6.2e-4

$$\sigma_{x,y} = \sqrt{\epsilon_{x,y}\beta(z)_{x,y} + (D_{x,y}\frac{\Delta p}{p})^2}$$

where

ϵ : Emittance at z=0

β : Beta function at z=RPSF

D : Momentum dispersion at z=RPSF

$\frac{\Delta p}{p}$: Momentum spread at z=0

1σ calculation	$1\sigma_x$	$1\sigma_y$
ep β @ IR-8 RPSF (Old)	0.314867	0.1629770
ep β @ IR-8 RPSF (new)	0.146677	0.140271
eAu β @ IR-8 RPSF (new)	0.156271	0.104160

Nuclear Breakups Distribution

BeAGLE v1.01.01

Phys. Rev. D 104, 114030

produced particle	rate
only neutron	7.66%
only proton	0%
only photon	3.25%
neutron and proton	3.19 %
neutron and photon	44.24 %
proton and photon	2.27 %
neutron, proton and photon	39.39 %

TABLE II. Summary of particles produced in incoherent J/ψ production in BeAGLE.

BeAGLE v1.03.02

Nuclear Breakups at Final State	Number of Events
Only Neutrons	7.86 %
Only Protons	0.0001 %
Only Photons	3.45 %
Neutrons + Protons	3.18 %
Neutrons + Photons	45.41 %
Protons + Photons	1.85 %
Neutrons + Protons + Photons	38.25 %

About **95 %** of events have **neutrons**

Remaining (Non-Vetoed) Events

Phys. Rev. D 104, 114030


- Veto.1: no activity other than e^- and J/ψ in the main detector ($|\eta| < 4.0$ and $p_T > 100$ MeV/c);
- Veto.2: Veto.1 and no neutron in ZDC;
- Veto.3: Veto.2 and no proton in RP;
- Veto.4: Veto.3 and no proton in OMDs;
- Veto.5: Veto.4 and no proton in B0;
- Veto.6: Veto.5 and no photon in B0;
- Veto.7: Veto.6 and no photon with $E > 50$ MeV in ZDC.

Material	Survived Event Ratio			
	Without beam pipe	Beryllium	Aluminum	Stainless Steel
Total events	100 %	100 %	100%	100%
Veto.1	86.9%	86.9%	86.9%	86.9 %
Veto.2	5.81%	9.73%	9.85%	17.2%
Veto.3	5.81%	9.73 %	9.85%	17.2%
Veto.4	5.09%	8.77%	8.89%	15.73%
Veto.5	4.32%	6.22%	5.97%	10.18%
Veto.6	2.29%	3.32%	3.18%	5.68%
Veto.7 ($E_{\text{photon}} > 50$ MeV)	1.06%	2.05%	2.46%	5.58%
Veto.7 ($E_{\text{photon}} > 100$ MeV)	-	2.18%	-	-

TABLE III. Summary of the percentage of events surviving the different vetoing steps for incoherent events assuming no beam pipe and different beam pipe materials of beryllium, aluminum, and stainless steel.

Veto Selections	Surviving Events
	eAu β @ IR-8 RPSF
All events	997,820
Events with one scattered electron identified and $ \eta_{J/\psi} < 4$ and $1 < Q^2 < 10$	732,455 (100 %)
ZDC HCAL tagged	41,848 (5.71339 %)
+ RPSF tagged	71 (0.00969343 %)
+ OMD tagged	71 (0.00969343 %)
+ B0 tracker tagged	30 (0.00409581 %)
+ B0 ecal tagged	17 (0.00232096 %)
+ ZDC ECAL tagged	4 (0.000546109 %)

Hit Positions of Protons w/ & w/o Magnets



Neutrons* are separable from Protons

