

# Simulation Study of the Gen-II CALI prototype


Weibin Zhang

UC Riverside

2024-03-01

# Gen-II Prototype

- Beam test of Gen-I prototype in Hall D at JLab on Jan 23, 2023 [Instruments 2023, 7\(4\), 43](#)
- 40 channels in Gen-I prototype
- 300 channels in Gen-II prototype
- 4 hexagonal layers + 10 square layers





## Test Position




- $\gtrsim 7$  m away from the IP
- North of the beampipe,  $\sim 47$  cm away from the beampipe center ( $3.2 < \eta < 3.6$ )
- Roughly aligned in height
- Be parallel to the beampipe<sup>3</sup>

## Simulation




- DD4HEP framework
- 7 layers of hexagonal cell (7.9 cm<sup>2</sup>), 13 layers of square cell (22.5 cm<sup>2</sup>)
- 7 m away from the beam source,  $0 < \theta < 0.001$
- Hepmc3 file for  $\pi^0$  events, making sure the 2 photons hit the prototype

# Event Display: 40 GeV Photon




# Sampling Fraction



- $sf = 0.01$


# Event Energy

event energy (GeV)




5 GeV photon  
 $3.145/5 \sim 60\%$

event energy (GeV)



15 GeV photon  
 $12.1/15 \sim 80\%$

event energy (GeV)




30 GeV photon  
 $27.85/30 \sim 93\%$

- sf is energy dependent


# Hit Energy Distribution

hit energy (GeV)





# Layer Energy Distribution





# Reconstruction: Eicrecon


## Imaging topological cluster algorithm

```
app->Add(new J0mniFactoryGeneratorT<ImagingTopoCluster_factory>(
    "CALIImagingTopoClusters",
    {"CALIReCHits"},
    {"CALIImagingTopoClusters"},
    { 7 layers of hexagonal cell (7.9~cm2), 13 layers of square cell
      .neighbourLayersRange = 1,
      .localDistXY = {55*dd4hep::mm, 50*dd4hep::mm},
      // .layerDistEtaPhi = {0.9, 0.5},
      // .sectorDist = 10.0 * dd4hep::cm,
      .minClusterHitEdep = 10.0 * dd4hep::MeV,
      .minClusterCenterEdep = 1000.0 * dd4hep::MeV,
      .minClusterEdep4 = 3000.0 * dd4hep::MeV,
      .minClusterNhits = 1,
    },
    app // TODO: Remove me once fixed (ama/40GeV_prototype_event)
));
```

# Cluster




# $\pi^0$ : Photon Distributions




- The larger the opening angle, the larger the energy difference
- The lower energy photon can go to very low energy
- The higher the  $\pi^0$  energy, the smaller the minimum opening angle, the harder to distinguish the decay photons

# Event Display: 40 GeV $\pi^0$




# $\pi^0$ : Number of Clusters


$\Delta\theta = 6 - 7$  mrad




$\Delta\theta = 7 - 8$  mrad




$\Delta\theta = 8 - 9$  mrad




$\Delta\theta = 9 - 10$  mrad



$\Delta\theta = 10 - 11$  mrad




$\Delta\theta = 11 - 12$  mrad



# $\pi^0$ : Cluster Energy


$\Delta\theta = 6 - 7$  mrad

Cluster energy




$\Delta\theta = 7 - 8$  mrad

Cluster energy




$\Delta\theta = 8 - 9$  mrad

Cluster energy




$\Delta\theta = 9 - 10$  mrad

Cluster energy




$\Delta\theta = 10 - 11$  mrad

Cluster energy



$\Delta\theta = 11 - 12$  mrad


Cluster energy



# $\pi^0$ : Invariant Mass


$\Delta\theta = 6 - 7$  mrad

invariant mass




$\Delta\theta = 7 - 8$  mrad

invariant mass




$\Delta\theta = 8 - 9$  mrad

invariant mass




$\Delta\theta = 9 - 10$  mrad

invariant mass




$\Delta\theta = 10 - 11$  mrad

invariant mass



$\Delta\theta = 11 - 12$  mrad

invariant mass





# Summary


- Simulation of high energy photon and  $\pi^0$  with the Gen-II prototype
- Cluster reconstruction with the ImagingTopological algorithm
- Reconstruct the  $\pi^0$  events
- Next step: update the prototype and simulate the real beam conditions

# Backup

## Invariant Mass


$$m = \sqrt{(e_1 + e_2)^2 - \sum_{i=x,y,z} \left( e_1 \frac{i_1}{l_1} + e_2 \frac{i_2}{l_2} \right)^2}$$

## Separating Clusters In Square Cells



- Need a thin cluster: high `minClusterHitEdep`
- A large separation between the two decayed photons

## Separating Clusters



- Hexagon side:  $s = 1.74 \text{ cm}$
- Separation length:  $dx = 6 \times s = 10.44 \text{ cm}$