

Brief Overview of Λ_c Related Physics and Results

Joseph Bertaux

Purdue University


November 22, 2024



- 1 Physics introduction
- 2 Status of Λ_C analysis in the field
- 3 Analysis

- 1 Physics introduction
- 2 Status of Λ_C analysis in the field
- 3 Analysis

Standard Model of Elementary Particles



The standard model of particle physics. Image source: https://en.wikipedia.org/wiki/File:Standard_Model_of_Elementary_Particles.svg



Overview

- Proposed in 1964 as a solution to the Λ^{++} 's apparent violation of the Pauli exclusion principle (uuu)
- Needed another degree of freedom to distinguish 3 up quarks
 - 2 of which must have the same spin
- The additional property is color (red, green, blue)

Some takeaways

- No naturally occurring particles that have a non-zero color charge
 - Quarks are bound to color-neutral hadronic states (confinement)
- Gluons are self-interacting and this leads to an approximately distance-independent force ~ 1 GeV/fm
- QCD coupling α_S varies with energy scale
 - This allows perturbation theory to be applied at collider energies

- ① $q\bar{q}$ pairs are produced at high energy
- ② $q\bar{q}$ energy due to separation is sufficient to produce new $q\bar{q}$ pairs
 - This repeats until new $q\bar{q}$ pairs can be produced
- ③ The now lower energy quarks coalesce to form hadrons



- The prediction of the QGP state followed the discovery of the asymptotic freedom of the QCD coupling constant (1970s)
- This was first observed at RHIC in 2004
- Motivations for studying QGP:
 - State of the universe immediately following the Big Bang
 - Smallest specific viscosity (viscosity to entropy density)





Figure: (Adapted from [14])



- ① Collision:
 - $T \sim \sqrt{s_{NN}}$
 - $\mu_B \sim q$ - \bar{q} separation (small)
- ② Hadronization:
 - Energy loss from pair production
 - μ_B decreases (deconfinement)
- ③ Freeze-out
 - Too "cold" for q - \bar{q} production
 - μ_B increases (now separated, confined hadrons)

Figure: (Adapted from [16])

One important quantity is the nuclear modification factor,


$$R_{AB} = \frac{1}{\langle T_{AB} \rangle} \frac{dN_{AB}^x}{dp_T} \bigg/ \frac{d\sigma_{pp}^x}{dp_T} \quad (1)$$

This gives the production of x in collisions between species A , B relative to the yield in proton-proton collisions, where $\langle T_{AB} \rangle$ is the mean nuclear overlap function


$$\frac{1}{N_{AB}^{inel}} \frac{dN_{AB}^x}{dp_T} = \langle T_{AB} \rangle \frac{d\sigma_{pp}^x}{dp_T} \quad (2)$$

It gives the proper normalization for production in AB collisions considering there are more inelastic parton collisions there than in pp .


- Spin $\frac{1}{2}$, Isospin 0 family of baryons
 - Each contains one u quark and one d quark
 - (Recall isospin refers the u and d quark content of a hadron)
- A third quark is from a higher generation (s , c , b)



$$\Lambda^0, \tau \sim 10^{-10}\text{s}$$



$$\Lambda_c^+, \tau \sim 10^{-13}\text{s}$$



$$\Lambda_b^0, \tau \sim 10^{-12}\text{s}$$

- Theoretically, there is also be a Λ_t^+
 - But this will likely decay before it can hadronize

- 1 Physics introduction
- 2 Status of Λ_C analysis in the field
- 3 Analysis

Studying hadrons containing heavy quarks is preferable, as

- they are produced in initial hard scatterings on a timescale shorter than QGP formation, $\mathcal{O}(\text{fm}/c)$
- they have a lower annihilation rate than lighter quarks
- their mass is sufficient such that low- p_T hard-scattering processes can be treated perturbatively

Baryon/meson ratios are analyzed to better understand hadronization and coalescence; in this case,

- $\Lambda_c^+ \rightarrow pK^-\pi^+$
- $D_0 \rightarrow K^-\pi^+$

Many probes are done at CMS (5.02 TeV) or RHIC (200 GeV) energies; we have data for:

- Λ_c/D_0 ratio in CMS and ALICE conditions in pp
- Λ_c/D_0 ratio in RHIC conditions in AuAu

Both saw enhancement in this baryon/meson ratio, but RHIC saw this to 2.1σ significance over predictions using PYTHIA8.3 So, another measurement is needed:

- Λ_c/D_0 ratio in RHIC conditions in pp

This will help determine if there is further Λ_c/D_0 enhancement which can be attributed to the coalescence process

Figures from the STAR and CMS papers showing Λ_c/D_0 enhancement over the PHIA prediction




Figure: The STAR result (Figure 2 in [22])




Figure: The CMS result (Figure 7 in [18])


There are practical difficulties associated with Λ_c reconstruction

- 3-daughter decay (as opposed to 2-daughter decay)
 - The most probable decay branch has three daughter particles
 $\Lambda_c^+ \rightarrow pK^-\pi^+$
 - The additional daughter candidate means there is an order of magnitude higher background than when reconstructing 2-daughter decays
- Short lifetime
 - Time of flight $\mathcal{O}(10^{-13} \text{ s})$
- Need a high-resolution vertex detector
 - the MVTX (Maps-based VerTeX detector) from sPHENIX
 - the ITS-2 from ALICE

- 1 Physics introduction
- 2 Status of Λ_C analysis in the field
- 3 Analysis

- Methods:
 - ROOT's TMVA package (Multi-Variable Analysis)
 - BDTs (Boosted Decision Trees)
 - Each node of a decision tree evaluates an inequality which depends on a training variable
 - The truth value of the inequality determines which subtree is used for further decisions
 - For boosted trees, additional trees are added to minimize the residuals of the previous training epoch
 - Possibly use DNN (Deep Neural Network)
- Variables:
 - 2D decay length significance
 - 2D pointing angle
 - transverse momentum ratio of daughter tracks (p_T / σ_{p_T})

- 2D decay length significance
 - decay length / decay length error
 - Distinguishes secondary decays from primary vertex
- 2D pointing angle
 - angle between parent momentum and decay displacement
 - Identifies prompt decays
- “2D” meaning projection in x-y plane




For signal,

- Use Monte-Carlo generated simulation data with truth reconstruction
- Physical parameters of the reconstructed track are used in training when the track is a daughter of a Λ_c .

For background,

- Fit the mass spectrum of reconstructed Λ_c to a Gaussian distribution
- Choose data in the “sidebands” of this distribution ensures that reconstructed mass points from real data are unlikely to contain real Λ_c baryons



Plot showing sidebands from ([15]). The yield of a mass spectrum can be taken as the coefficient of the (normalized) signal probability distribution

- [1] S. Acharya and Acosta *et al.* (ALICE Collaboration).
 Λ_c^+ production in pp collisions at $\sqrt{s} = 7$ TeV and in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV.
Journal of High Energy Physics, 2018(4), April 2018.
- [2] A. *et al.* Andronic.
Heavy-flavour and quarkonium production in the LHC era: from proton–proton to heavy-ion collisions.
The European Physical Journal C, 76(3), February 2016.
- [3] Christian Bierlich, Smita Chakraborty, Nishita Desai, Leif Gellersen, Ilkka Helenius, Philip Ilten, Leif Lönnblad, Stephen Mrenna, Stefan Prestel, Christian T. Preuss, Torbjörn Sjöstrand, Peter Skands, Marius Uthmeim, and Rob Verheyen.
A comprehensive guide to the physics and usage of pythia 8.3, 2022.

- [4] Renè Brun and Fons Rademakers.
Root - an object-oriented data analysis framework.
1997.
- [5] Wit Busza, Krishna Rajagopal, and Wilke van der Schee.
Heavy ion collisions: The big picture and the big questions.
Annual Review of Nuclear and Particle Science, 68(1):339–376,
October 2018.
- [6] Debashis Ghoshal.
Current Perspectives in High Energy Physics.
Hindustan Book Agency, New Dehli, India, 2005.
- [7] David Griffiths.
Introduction to Elementary Particles.
WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim, 2008.

- [8] David J. Gross and Frank Wilczek.
Ultraviolet behavior of non-abelian gauge theories.
Phys. Rev. Lett., 30:1343–1346, Jun 1973.
- [9] Michael L. Miller, Klaus Reygers, Stephen J. Sanders, and Peter Steinberg.
Glauber modeling in high-energy nuclear collisions.
Annual Review of Nuclear and Particle Science, 57(1):205–243,
November 2007.
- [10] Kazuya Nagashima.
Energy loss of charm and bottom quarks in Quark-Gluon Plasma created in Au+Au collisions at 200 GeV.
PhD thesis, Hiroshima University, 2019.

- [11] H. David Politzer.
Reliable perturbative results for strong interactions?
Phys. Rev. Lett., 30:1346–1349, Jun 1973.
- [12] F. Reidt.
Upgrade of the alice its detector.
Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment, 1032:166632, June 2022.
- [13] Berndt Mueller *et al.*
The relativistic heavy ion collider status and future, 2013.
- [14] M. Arslanok *et al.*
Hot qcd white paper, 2023.

- [15] Soumik Chandra *et al.*
Production of prompt λ_c^+ baryons in proton-proton and lead-lead collisions at 5.02 TeV (2017/2018 data).
- [16] Yasuyuki Akiba *et al.*
The hot qcd white paper: Exploring the phases of qcd at rhic and the lhc, 2015.
- [17] I. Arsene *et al.* (BRAHMS Collaboration).
Quark-gluon plasma and color glass condensate at rhic? the perspective from the brahms experiment.
Nuclear Physics A, 757(1-2):1-27, August 2005.

- [18] A. Tumasyan *et al.* (CMS Collaboration).
Study of charm hadronization with prompt Λ_c^+ baryons in proton-proton and lead-lead collisions at $\sqrt{s_{NN}} = 5.02$ TeV.
Journal of High Energy Physics, 2024(1), January 2024.
- [19] K. Adcox *et al.* (PHENIX Collaboration).
Formation of dense partonic matter in relativistic nucleus–nucleus collisions at rhic: Experimental evaluation by the phenix collaboration.
Nuclear Physics A, 757(1–2):184–283, August 2005.
- [20] B.B.Back *et al.* (PHOBOS Collaboration).
The phobos perspective on discoveries at rhic.
Nuclear Physics A, 757(1–2):28–101, August 2005.

[21] J. Adams *et al.* (STAR Collaboration).

Experimental and theoretical challenges in the search for the quark–gluon plasma: The star collaboration’s critical assessment of the evidence from rhic collisions.

Nuclear Physics A, 757(1–2):102–183, August 2005.

[22] J. Adams *et al.* (STAR Collaboration).

First measurement of Λ_c baryon production in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV.

Physical Review Letters, 124(17), May 2020.