


# Tau benchmark in the Insert






Sebouh Paul  
UC Riverside  
10/22/2024

# $\tau$ benchmark as proxy for jets

- Simulate  $\tau$  with  $3.0 < \eta < 4.0$
- Allow them to decay in dd4hep
  - ~65% of  $\tau$  decays are hadronic
  - In analysis, only select events without a  $\nu_e$  nor a  $\nu_\mu$
- Truth “hadronic final state” four momentum,  $\mathbf{p}_{\text{hfs}} = \mathbf{p}_T - \mathbf{p}_{\text{VT}}$ 
  - Further require  $m_{\text{hfs}} > m_{\pi^\pm}$  to ensure that there is more than one hadron in the jet




# More events



# Energy reconstruction

- Sum of energies of all clusters in FEMC, insert and LFHCAL.
  - TODO include corrections for non-compensation of the insert
- Compared to the energy of the hadronic final state of the tau decay




# Theta reconstruction

- Direction determined by sum of momenta, assuming all particles are massless

$$\vec{p}_{\text{tot}} = \sum_{i \in \text{clusters}} E_i \frac{\vec{x}_i}{|\vec{x}_i|}$$

- This should improve when corrections are included




# Theta reconstruction

- Direction determined by sum of momenta, assuming all particles are massless

$$\vec{p}_{\text{tot}} = \sum_{i \in \text{clusters}} E_i \frac{\vec{x}_i}{|\vec{x}_i|}$$

- This should improve once energy corrections are included



# Summary

- A benchmark is being developed which uses tau decay as a source of jets in the insert
- Plans:
  - Distinguish between hadrons and photons (which come from  $\pi^0$  decay) in jets
  - Improve energy and theta reconstruction with corrections for the non-compensating nature of the detector
  - Increase statistics in order to determine  $\eta$  dependence of performance