


The Origin of Ultra-high Energy* Cosmic Rays




Glennys R. Farrar, New York University
BNL, Oct. 23, 2025

$\langle E \text{ per nucleon} \rangle \sim 2 \text{ EeV} \rightarrow \gamma > 10^9$

UHECR trivia

- Highest energy CR $> 200 \text{ EeV} \equiv 2 \times 10^{20} \text{ eV}$ — a single nucleus with the kinetic energy of the highest energy pro golf ball
- 100 EeV neutron: Relativistic time dilation ($\gamma = E/mc^2 \approx 10^{11}$) $\Rightarrow$ lifetime is a million years $\Rightarrow$ can travel across a galaxy without decaying
- Flux of CRs above 150 EeV $\approx 0.2 / \text{km}^2 / \text{millennium}$

It's been a long, hard search



Aeronautisches Gelände im Wiener Prater, von dem aus V. F. Hess in den Jahren 1911/12 seine ersten Freiballon-Forschungsfahrten unternommen hatte. (Courtesy of Heeresgeschichtliche Museum, Vienna)
<Ed> Contributed by R. Steinmaurer. See p. 17.





Hess on gondola in 1912 probably in test flight. The date and place is not clear at present.
<Ed> Contributed by R. Steinmaurer. See p. 17.

Hess: CRs
1911 or 1912

3




Linsley: 1st evt > 100 EeV
Volcano Ranch, NM ~1962



6

Telescope Array, Utah
Amaterasu ('23): 240 EeV



Pierre Auger Obs., Argentina
~50 evts > 100 EeV




1700 stations, 3000 km²




Fly's Eye Utah 1991
OMG: 320(250) EeV

How to deduce the mass and energy of a UHECR



- Depth of first interaction:
 - heavy nucleus: interacts quickly (starts high)
 - proton: 1st interaction is deep or shallow
- Shower development:
 - heavy nucleus: shower develops quickly
 - proton: more interactions needed to reach shower max
 - primary energy from integrated fluorescence emission
- Ground signal:
 - EM vs muon components $\Rightarrow$ nuclear mass
 - primary energy from total signal

The Hybrid Observation Method of Auger




$E^3 \times \text{Spectrum}$

Composition gets heavier with energy

$\langle X_{\text{max}} \rangle$

A single mass group dominates at each E

$\sigma(X_{\text{max}})$






Implications of the observations

- Composition is mixed, mass increases with E
- *Narrow range of masses at each E :*

Rigidity spectrum is narrow: $\langle R \equiv E/Z_e \rangle \approx 4$ EV

- Arrival directions: *Sources are abundant*
- No prominent sources for highest E events: *Sources are transients*



Binary Neutron Star Mergers

only (currently known) source candidate satisfying all criteria

- Universal rigidity spectrum explained:


1. Magnetic field is generated by *gravitational* dynamo 💕💕

2. Mass range of BNS is narrow 💕💕

Known BNS's: $M = 2.64 \pm 0.14$ (5%) $M_\odot$
($3.2 M_\odot \rightarrow 10\%$ (negligible) spread)

- UHECR energy injection rate promising:

- need UHECR energy per merger $\approx 10^{50}$ erg ($< 1\%$ of total energy emitted)



Binary Neutron Star Mergers


only (currently known) source candidate satisfying all criteria

- Universal rigidity spectrum explained:

1. Magnetic field is generated by *gravitational* dynamo 💕💕

2. Mass range of BNS is narrow 💕💕

Known BNS's: $M = 2.64 \pm 0.14$ (5%) $M_{\odot}$






Unique predictions enable definitive tests:

- EHE ν 's $\leftrightarrow$ gravitational waves
- Highest energy UHECRs: $Z > 26$
- BNS merger $\rightarrow$ initial B $\rightarrow$ predict spectrum

Questions we can answer now

- **Where in the merger ejecta does acceleration occur?**
(predict spectrum & cutoff)
- **What is the time profile for UHECR production?**
(coincidences between GW and EHE ν 's?)
- **Abundances of different UHECR nuclei?**
(r-process nucleosynthesis)



Outflow after merger (Kasen+2017)




Topics for today

- ✓ **UHECR acceleration occurs in the magnetized outflow**
→ Predicts rigidity cutoff R_{cut} and spectral shape
- ✓ **UHECR production occurs ~ 1 day after merger**
(coincidences between GW and EHE ν 's...)
- **Abundances of different UHECR nuclei**

Outflow after merger (Kasen+2017)




UHECR abundances



r-process abundances



Big Picture

time →




grav-driven dynamo → B
t = 150 ms, 500 km

**Gravitational Wave
Emitted**



cools to ~ 1 MeV
nucleosynthesis
~1 s: r ~ 10^10 cm

*r-process
nucleosynthesis*



B drops till synch losses
are subdominant


UHECR acceleration
~1 day: 10^14 cm

$$\tau_{\text{accel}} / \tau_{\text{synch-loss}} \sim 1$$

Big Picture

time

Expands at $v \sim 0.1 c$




inspiral
GRMHD dynamics
→ huge B


"Hypervnova"
0 - few sec

initialization of B
150ms, 500 km

Kiuchi+23 → B_{ini}




cools to ~ 1 MeV
nucleosynthesis
 ~ 1 s: $r \sim 10^{10}$ cm




UHE CR's
accelerated
 ~ 1 day

Gravitational Wave
Emitted





Nucleosynthesis:

- ▶ p, He in jet
- ▶ $Z, N \geq 20$ elsewhere
- ▶ r-process nuclei



Merging NS's produce "r-process" elements



Periodic table showing the cosmogenic origin of each element. The elements heavier than iron with origins in supernovae are typically those produced by the *r*-process, which is powered by supernova neutron bursts

r-process nucleosynthesis B²FH

REVIEWS OF MODERN PHYSICS

VOLUME 29, NUMBER 4

OCTOBER, 1957

Synthesis of the Elements in Stars*

E. MARGARET BURBIDGE, G. R. BURBIDGE, WILLIAM A. FOWLER, AND F. HOYLE


*Kellogg Radiation Laboratory, California Institute of Technology, and
Mount Wilson and Palomar Observatories, Carnegie Institution of Washington,
California Institute of Technology, Pasadena, California*

"It is the stars, The stars above us, govern our conditions";
(*King Lear*, Act IV, Scene 3)

but perhaps

"The fault, dear Brutus, is not in our stars, But in ourselves,"
(*Julius Caesar*, Act I, Scene 2)


r process.—The nuclear physics of this process demands that neutrons be added extremely rapidly, so that the total time-scale for the addition of a maximum of about 200 neutrons per iron nucleus is ~ 10 –100 sec.




Big Picture

time


Expands at $v \sim 0.1 c$




Gravitational Wave
Emitted




Magnetically dominated turbulence



$^* \sigma \approx U_{\text{mag}} / K_{\text{E}} \text{den}$



UHECRs produced
~1 day: 10^{14} cm




UHECRs mainly produced in the turbulent outflow (maybe some UHECRs also from jets)

- Composition: UHECRs have masses up to (at least) Fe, but jets only have p, He (Perego+22)

- Spectrum: $E^{-2.x}$ $\times$ function of (E/E_{cut})


UHECR data strongly favor cutoff: $\text{sech}[(E/E_{\text{cut}})^2]$



- Outflow: Magnetized turbulence $\rightarrow$ cutoff $\text{sech}[(E/E_{\text{cut}})^2]$ (Comisso, GRF, Muzio ApJL 2024)
- Jets: Diffusive Shock Acceleration cutoff: $\exp(-E/E_{\text{cut}})$??? (Protheroe+Stanev 1997)


Peak energy when acceleration = loss

- Initialize $B(r=500 \text{ km}) = 3.3 \cdot 10^{12} \text{ G}$; $L_{\text{coh}} \approx r/3$ (Kiuchi+2023)
- Homologous expansion: $B(r) = B_0 (r/r_0)^{-3/2}$ (Rosswog+2014)
- Maximum rigidity CRs at radius r : **Rigidity $R \equiv E/Z$**



- $R_{\text{max}}(r) = 0.65 B(r) L_{\text{coh}}(r)$ (turb. accel. CFM24)
- but require $\tau_{\text{synch-loss}}(A, Z, R, r) \approx \tau_{\text{accel}}(R, B)$
- **$\rightarrow R_{\text{cut}}(Z, A, r)$ is intersection**



$$\dot{E}_{\text{synch}} = \frac{4}{9} c \left(\frac{m_e r_0}{m_p} \right)^2 (\gamma \beta B)^2 \frac{Z^4}{A^2}$$



BNS initialization → no-free-parameter *predictions* of R_{cut}

rigidity $R \equiv E/Z$

- **Turbulent outflow:** $R_{\text{cut}} \{p, \text{He}, \text{O}, \text{Si}, \text{Fe}, \text{Te}\} \approx \{6.2, 9.4, 7.1, 6.3, 6.0, 5.9\}$ EV
 - **Auger fit:** $R_{\text{cut}} = 6.3$ EV (has factor-2 uncertainty)
 - **Jet:** some proton and/or He; $E_{\text{cut}} \approx 11$ EeV for proton, $E_{\text{cut}} \approx 35$ EeV for He
- Additional light component improves Auger fit by 5σ (Muzio+2019)



Tests of BNS merger scenario

- ✓ Successful prediction of R_{cut}
- ✓ Eroded r-process nuclei among UCRs ($A > 56$)
- Possible characteristic pattern of R_{cut} slightly varying with A , reflecting synchrotron-limited acceleration
- Flux of EHE neutrinos
- GW- ν coincidences for EHE ν 's ($E > \text{PeV}$) - time delay ≈ 1 day

Test 1: $E > 150 \text{ EeV}$ events are from r-process primaries



- $E = R Z_{\text{Te-Xe}} \approx 4.5 \text{ EV} \times (52-54) = 240 \text{ EeV}$
- $E_{\text{cut,Fe}} \approx 5 \text{ EV} \times 26 = 130 \text{ EeV} + 1\text{-}\sigma \text{ E uncertainty} \rightarrow 150 \text{ EeV}$
- ★ Auger should look closely at the ~ 10 events with $E > 130 \text{ EeV}$, to see if they favor $A > 56$.

Test 2: EHE Neutrino $\leftrightarrow$ Grav. Wave

- Neutrinos come from UHECRs:
 - spallation neutrons beta decay: $E_\nu \approx 10^{-3} (E_n \approx \mathbf{R}/2) \sim 2 \text{ PeV}$
 - photo-pion production: $E_\nu \approx (E/A)/20 \sim \mathbf{R}/40 \sim 100 \text{ PeV}$
- Time delay for ν is $>$ time for ejecta to expand to 10^{14} cm where UHECRs form:
 - $\langle v_{ej} \rangle = 0.1 c \rightarrow$ delay of $O(\text{day})$

To do: make better estimate of rate of observing GW-nu coincidences

very crude: 1/yr with Cosmic Explorer, Einstein telescope & ICGen2

Source candidates vs key constraints

	$n_s \gtrsim 10^{-3.5}$ Mpc ⁻³	energy injection	ordinary galaxy	Universal $R_{\max}$	Highest energy events
Powerful AGN	[X]	✓	X	X	X
Long GRBs	[X]	X	X	X	X
Tidal Disruption Events	?	?	✓	X	X
Accretion Shocks	?	?	[X]	X	X
BNS mergers	✓	[✓]	✓	✓	✓

(All can satisfy Hillas size > Larmor radius)

Summary: BNS mergers explain essential features of UHECRs

- Minimal source-to-source variation
- Spectrum from magnetized turbulence: $\sim E^{-2.1} \text{sech}[(E/E_{\text{cut}})^2]$
 - $R_{\text{cut}} \approx 6\text{-}9 \text{ EV}$ (observed: $R_{\text{cut}} = 6.3^{+6.3}_{-2.3} \text{ EV}$)
- First explanation for highest energy events: they originate as $A \sim 130$ (r-process peak at Tellurium/Xenon).

$$E = R Z_{\text{Te-Xe}} \approx 4.5 \text{ EV} \times (52\text{-}54) = 240 \text{ EeV}$$

$$E_{\text{OMG}} \approx 250 \pm 70 \text{ EeV}, \quad E_{\text{Amaterasu}} \approx 212 \pm 25 \text{ EeV}$$

Thank you!