

$$H_1(\phi, J) = v \cdot J + (2J)^{\frac{3}{2}} G_{3,N} \cos(3\Phi + \xi_{3,N})$$

$$\hat{G}_{k,n} = G_{k,n} \cdot e^{i\xi_{k,n}} = \frac{1}{48\pi} \oint S(s) \beta(s)^{\frac{3}{2}} e^{i[k \cdot \chi(s) - (kv-n)\theta(s)]} ds$$

6th Slow Extraction Workshop

October 6 - 9, 2025

Stony Brook University Wang Center

hosted by Brookhaven National Laboratory | <https://www.bnl.gov/sx-workshop/>

Local Organizing Committee

Christina Blas
Anna Petway
Bryan Callaghan
Lucy Lin
Eiad Hamwi (Cornell)

International Organizing Committee

Kevin Brown (BNL Chair)	Marco Pullia (CNAO)
Peter Forck (GSI 2024)	Elisabeth Renner (TU Wien)
Matthew Fraser (CERN)	Christian Schömers (HIT)
Vladimir Nagaslaev (Fermilab 2019 chair)	Hikaru Souda (Yamagata Univ.)
Ryotaro Muto (J-PARC)	Masahito Tomizawa (J-PARC 2022 chair)
Alessio Mereghetti (CNAO)	Francesco Velotti (CERN)
David Ondreka (GSI)	Jiancheng Yang (IMP/CAS)
Mauro Pivi (MedAustron 2024)	Ex-officio
Dale Prokopovich (MedAustron 2024)	Brennan Goddard (CERN, 2017 chair)

Photo used with permission from CERN. A prototype septum for the electron-positron-accumulator ring.