

Poincaré-friendly tracking with complex tune modulation

Steve Peggs (BNL)

Henry Lovelace III (BNL & U. Huddersfield),
Guillaume Robert-Demolaize (BNL),
Todd Satogata (JLab)

October 8, 2025



Goal: Construct a tune modulation “probe” element that can be added to a lattice to better evaluate transverse vulnerability to complex tune modulation.

Physical problem: Bad things can happen when a tune modulation spectrum *drive line* aligns with a resonance island *response line*: $Q_M \approx Q_I$

Complexity:

- Many resonances co-exist, weak & strong
- Polynomial chromaticities
- Multi-RF bucket structure (eg RHIC)
- Transverse chaos in (eg) H drives diffusion in V


Key subsidiary goal: re-enable Poincaré surface-of-section graphic insight.

Past

3 key tune domain parameters



- 1) Base tune Q_0 for small amplitude a particles
- 2) Resonance tune, eg $Q_R = 1/6$ when $Q(a_R) = 1/6$
- 3) Island tune Q_I



Near island center on turn t

$$a(t) \approx a_R + \hat{a} \sin(2\pi Q_I t)$$



$$\phi(t) \approx \phi_R + \hat{\phi} \cos(2\pi Q_I t)$$

Single tune modulation tune



Base tune modulation amplitude & tune are q and Q_M , so

$$Q_0(t) = Q_{00} + q \cos(2\pi Q_M t)$$



Generic theory & particular simulation agree ... plus structure


Tune domain dynamical zones




$(q, Q_M) = (0.001, 0.0052)$ below, while beam-beam tune shift parameter $\xi = 0.0042$ (Left) & 0.006 (Right)

Amplitude, a/σ

Amplitude, a/σ



Left: strong sidebands



Right: chaos


Present

Complex storage buckets



RHIC uses 2 RF systems in collision, helping IBS but ...


RHIC storage buckets with $(V_{28}, V_{197}) = (2.5, 0.3)$ MV 241225



Synchrotron tune versus $\hat{\delta}$




... causing Q_s to vary significantly with amplitude $\hat{\delta}$




Complex buckets require multiple harmonics $d_n(\hat{\delta})$ of $Q_s(\hat{\delta})$ to describe the turn-by-turn evolution of δ

$$\delta(\hat{\delta}, t) = \sum_{n=0}^{\infty} d_n(\hat{\delta}) \cos(2\pi n \cdot Q_s(\hat{\delta}) t)$$




Polynomial chromaticity has coefficients C_n where

$$\Delta Q(t) = \sum_{n=1}^{\infty} C_n \delta(t)^n$$



Putting this together delivers the q_n -harmonics

$$\Delta Q(\hat{\delta}, t) = \sum_{m=1}^{\infty} C_m \left(\sum_{n=0}^{\infty} d_n(\hat{\delta}) \cos(2\pi n \cdot Q_s(\hat{\delta}) \cdot t) \right)^m = \sum_{n=0}^{\infty} q_n(\hat{\delta}) \cos(2\pi n \cdot Q_s(\hat{\delta}) \cdot t)$$





Hénon said that “[a single sextupole lattice] exhibits ... all the typical properties of more complicated mappings and dynamical issues”.

Agreed!


We are using single sextupole (or octupole) lattices to begin debugging software that will soon have:

- 1) Poincaré-friendly tracking
- 2) Multi-sextupoles & octupoles (RHIC & HSR)
- 3) Transverse diffusion probes


One sextupole with $Q_0 = 0.15$



One sextupole, $Q_0 = 0.15$, 250603




One sextupole, $Q_0 = 0.15$, 250602




Many resonances are present, each with its own Q_I

Island tunes Q_I vs base tune Q_0




Vertical dashed lines at base tunes $Q_0 = (0.14, 0.15, 0.16)$

Island tune response spectra



Significant trouble is possible when one of these *response* spectral lines is close to an $n \cdot Q_s(\hat{\delta})$ *drive* line — for some values of $\hat{\delta}$ at a given Q_0


Synchrotron harmonics cross island tunes




Behavior parameterized by $\hat{\delta}$



$\delta_{\text{init}} = 0.00124$, $T_s = 152$, all q on 250603



$\delta_{\text{init}} = 0.00300$, $T_s = 263$, all q on 250603



Surfaces-of-section are plotted **once per synchrotron period** $T_s(\hat{\delta})$, which is **forced to be an integer**.



Crisp Poincaré surface-of-section plots enable visual and quantitative **tune modulation probes**.

The synchrotron period T_s is forced to be an integer.

Longitudinal motion is represented by **externally imposed tune modulation**.

The 30-second movie [2_bucket_NLC.mp4](#) shows a sequence of many such plots, for

$$0 < \hat{\delta} < 0.006$$

Future



Repeat the same on-momentum probe of RHIC & HSR, with many chromaticity sextupoles (& octupoles).

Add vertical motion and complex tune modulation.

Connect to the RHIC/HSR on-line model ?

Embedded H & V `tunemod` elements are being tested with realistic lattices in the new tracking code Sparks

Later, embed in BMAD, Xsuite, ... ?

Goal: develop generic tools to probe tune modulation resilience or vulnerability, independent of other longitudinal effects.

Finis