




Deuteron Tagging

Jan Vanek

University of New Hampshire

ePIC Collaboration Meeting, BNL

01/23/2026




OVERVIEW

- Extraction of deuteron and neutron reduced cross section from HepMC and EICRecon simulation output
 - True MC information, without any beam or detector effects
 - Simulation setup:
 - eD DIS events at $10 \times 130 \text{ GeV}^2$ (simulation by Alex Jentsch)
 - 500k events for tagged protons and 500k events for tagged neutrons
 - Goal:
 - Implementation of full TDIS framework on the new simulation sample
 - Reproduction results from A. Jentsch, Z. Tu, C. Weiss: [Phys. Rev. C 104, 065205](#)
- Today:
 - Analysis overview
 - New results
 - New EICRecon simulation sample with full statistics (fix from last report)
 - Spectator proton tagging using reconstructed information from off-momentum detector
 - Spectator neutron tagging in HepMC


PHYSICS MOTIVATION

- Main motivation: Understanding of internal structure of nucleons
- Historically was studied via Deep Inelastic Scattering (DIS) on protons
 - Relatively experimentally easy as we can accelerate electrons or protons
 - Structure functions and Parton Distribution Functions (PDFs) of proton reasonably well known
 - Despite long experimental efforts, still many open questions:
 - Spin structure of nucleons, Transverse and Generalized PDFs...
- We need also to need to measure **neutron internal structure** for full understanding of nucleon internal structure and QCD in general
 - Extraction of structure function F_2 of neutron
- Main challenges:
 - Neutrons are neutral and cannot be easily accelerated
 - Generally difficult to handle free neutrons



EXPERIMENTAL METHOD OVERVIEW

- Possible experimental solution: **Tagged DIS (TDIS)**
 - DIS on deuteron (d)
 - Measurement of the scattered electron and the spectator p or n
- **Challenges:**
 - Neutron is bound and not free, but we are interested in free n structure functions
 - Possible solution – pole extrapolation
 - Detection of the spectator p or n
 - Very difficult in fixed target experiments, as spectators have very low momentum and generally will stay in the target
 - Possible to measure at collider, such as the future Electron Ion Collider (EIC), as “target” is boosted
 - Still challenging, as spectators are very close to the hadron beam
- **Advantages:**
 - Straightforward, when spectators are successfully detected
 - Possible to measure p and cross-check with standard methods



A. Jentsch, Z. Tu, Ch. Weiss: [Phys. Rev. C 104, 065205.](#)

DEUTERON REDUCED CROSS SECTION

- Differential cross section on d can be written in terms of deuteron reduced cross ($\sigma_{red,d}$) section and photon flux:

$$\sigma_d = Flux(x, Q^2) \times \sigma_{red,d} \times \frac{dx}{2} dQ^2 \frac{d\phi_{e'}}{2\pi} [2(2\pi)^3]^{-1} \frac{d\alpha_p}{\alpha_p} \frac{dp_{T,p}^2}{2} d\phi_p$$

- Photon flux:

$$Flux(x, Q^2) = \frac{2\pi\alpha_{em}^2 y^2}{Q^4(1-\epsilon)x} = \frac{2\pi\alpha_{em}^2 [1+(1-y)^2]}{Q^4 x}$$

- Deuteron reduced cross section

$$\sigma_{red,d} = \frac{1}{Flux} \frac{d\sigma_d}{dx dQ^2 (d\phi_{e'}/2\pi) d\Gamma_p}, \text{ where } d\Gamma_p = [2(2\pi)^3]^{-1} \frac{d\alpha_p}{\alpha_p} \frac{dp_{T,p}^2}{2} d\phi_p$$

- Measured reduced cross section (integrated over $d\phi_p$)

$$\bar{\sigma}_{red,d} = \left(\frac{2.568}{L_{int}} \right) \frac{Q^4 x}{2\pi\alpha_{em}^2 [1+(1-y)^2]} \frac{[4(2\pi)^3] \alpha_p}{\Delta\alpha_p} \frac{dN}{\Delta x \Delta Q^2 \Delta p_{T,p}^2}$$

- $\Delta x, \Delta Q^2, \Delta p_{T,p}^2, \Delta\alpha_p$ are bin widths

- Scattered electron
- Spectator proton + struck deuteron
- Photon flux

A. Jentsch, Z. Tu, C. Weiss: [Phys. Rev. C 104, 065205.](#)
M. Strikman and C. Weiss: [Phys. Rev. C 97, 035209.](#)
C. Weiss and W. Cosyn: [Phys. Rev. C 102, 065204.](#)

- Full azimuthal coverage for electron
 - $(\Delta\phi_{e'}/2\pi) = 2\pi/2\pi = 1$
- Full azimuthal coverage for spectator
 - $\Delta\phi_p = 2\pi$
 - We are also integrating over full 2π

POLE EXTRAPOLATION METHOD

$$\sigma_{red,n} = \frac{\bar{\sigma}_{red,d}(x, Q^2)}{[2(2\pi)^3]S_d(p_{pT}, \alpha_p)[pole]}$$

- $\bar{\sigma}_{red,n}$ at the pole corresponds to a free n
 - $p_{pT}^2 \rightarrow -a_T^2$ which means negative (unphysical) p_{pT}^2
- Solution is to experimentally measure $\sigma_{red,n}$ as a function of p_{pT}^2 for small positive values and extrapolate to the pole
 - We are using DIS on bound n to extract F_2 of free n
- Same measurement can be done for p
 - Cross check with proton F_2 extracted with traditional method
 - Can be used to select optimal deuteron spectral function





- We can measure/calculate this
- External input from model/experiment

- Deuteron spectral function
 - $S_d(p_{pT}, \alpha_p)[pole] = \frac{R}{(p_{pT}^2 + a_T^2)^2}$
 - Position of pole
 - $a_T^2 = m_N^2 - \alpha_p(2 - \alpha_p)\frac{M_d^2}{4}$
 - Extrapolation done for $\alpha_p = 1$
 - Residue of spectral function
 - $R = 2\alpha_p^2 m_N \Gamma^2 (2 - \alpha_p)$
 - $\Gamma^2 = 0.007885 \text{ GeV}$

A. Jentsch, Z. Tu, C. Weiss: [Phys. Rev. C 104, 065205](#).
M. Strikman and C. Weiss: [Phys. Rev. C 97, 035209](#).
C. Weiss and W. Cosyn: [Phys. Rev. C 102, 065204](#).





TAGGED PROTON — BeAGLE

- (top) Deuteron reduced cross section in BeAGLE calculated using tagged protons
- (bottom) Neutron reduced cross section in BeAGLE calculated using tagged protons
 - Red point is extrapolated to pole
- Minor updates since last time shown:
 - Selecting only events with one proton in spectator region
 - Added $\eta > 5.5$ cut
 - To match and compare to cuts in EICRecon







TAGGED PROTON – BeAGLE

- (top) Deuteron reduced cross section in BeAGLE calculated using tagged protons
- (bottom) Neutron reduced cross section in BeAGLE calculated using tagged protons
 - Red point is extrapolated to pole
- Minor updates since last time shown:
 - Selecting only events with one proton in spectator region
 - Added $\eta > 5.5$ cut
 - To match and compare to cuts in EICRecon







TAGGED PROTON – EICRecon

- (top) Deuteron reduced cross section in EICRecon calculated using tagged protons
- (bottom) Neutron reduced cross section in EICRecon calculated using tagged protons
 - Red point is extrapolated to pole
- Updates since last time shown:
 - Reconstructed the full HepMC sample in EICRecon
 - True MC results from HepMC and EICRecon are now consistent







TAGGED PROTON – EICRecon

- (top) Deuteron reduced cross section in EICRecon calculated using tagged protons
- (bottom) Neutron reduced cross section in EICRecon calculated using tagged protons
 - Red point is extrapolated to pole
- Updates since last time shown:
 - Reconstructed the full HepMC sample in EICRecon
 - True MC results from HepMC and EICRecon are now consistent







TAGGED PROTON – EICRecon

- (top) Deuteron reduced cross section in EICRecon calculated using tagged protons
- (bottom) Neutron reduced cross section in EICRecon calculated using tagged protons
 - Red point is extrapolated to pole
- Updates since last time shown:
 - Reconstructed the full HepMC sample in EICRecon
 - True MC results from HepMC and EICRecon are now consistent
 - Added spectator RC information (full black circles)
 - Only spectator information is using RC information from off-momentum detector
 - Scattered electron info. is true MC for now







TAGGED PROTON – EICRecon

- (top) Deuteron reduced cross section in EICRecon calculated using tagged protons
- (bottom) Neutron reduced cross section in EICRecon calculated using tagged protons
 - Red point is extrapolated to pole
- Updates since last time shown:
 - Reconstructed the full HepMC sample in EICRecon
 - True MC results from HepMC and EICRecon are now consistent
 - Added spectator RC information (full black circles)
 - Only spectator information is using RC information from off-momentum detector
 - Scattered electron info. is true MC for now







TAGGED NEUTRON — BeAGLE

- (top) Deuteron reduced cross section in BeAGLE calculated using tagged neutrons
- (bottom) Proton reduced cross section in BeAGLE calculated using tagged neutrons
 - Red point is extrapolated to pole
- New results with tagged neutrons in BeAGLE
- Position of pole calculated using **proton mass**
 - $a_T^2 = m_N^2 - \alpha_p(2 - \alpha_p) \frac{M_d^2}{4}$
- Different result for proton reduced cross section than in original paper
 - A. Jentsch, Z. Tu, C. Weiss: [Phys. Rev. C 104, 065205](#).







TAGGED NEUTRON — BeAGLE

- (top) Deuteron reduced cross section in BeAGLE calculated using tagged neutrons
- (bottom) Proton reduced cross section in BeAGLE calculated using tagged neutrons
 - Red point is extrapolated to pole
- New results with tagged neutrons in BeAGLE
- Position of pole calculated using **proton mass**
 - $a_T^2 = m_N^2 - \alpha_p(2 - \alpha_p) \frac{M_d^2}{4}$
- Different result for proton reduced cross section than in original paper
 - A. Jentsch, Z. Tu, C. Weiss: [Phys. Rev. C 104, 065205](#).







TAGGED NEUTRON — BeAGLE

- (top) Deuteron reduced cross section in BeAGLE calculated using tagged neutrons
- (bottom) Proton reduced cross section in BeAGLE calculated using tagged neutrons
 - Red point is extrapolated to pole
- New results with tagged neutrons in BeAGLE
- Position of pole calculated using **neutron mass**
 - $a_T^2 = m_N^2 - \alpha_p(2 - \alpha_p) \frac{M_d^2}{4}$
- Nucleon mass used for calculation of a_T^2 has major influence on resulting nucleon reduced cross section



TAGGED NEUTRON — BeAGLE

- (top) Deuteron reduced cross section in BeAGLE calculated using tagged neutrons
- (bottom) Proton reduced cross section in BeAGLE calculated using tagged neutrons
 - Red point is extrapolated to pole
- New results with tagged neutrons in BeAGLE
- Position of pole calculated using **neutron mass**
 - $a_T^2 = m_N^2 - \alpha_p(2 - \alpha_p) \frac{M_d^2}{4}$
- Nucleon mass used for calculation of a_T^2 has major influence on resulting nucleon reduced cross section



SUMMARY AND OUTLOOK

- Fully implemented TDIS framework for tagged protons for true MC from both BeAGLE and EICRecon
 - True MC information from BeAGLE and EICRecon appear to be consistent
- Added RC level information for spectator protons in EICRecon
 - First iteration of adding RC level information only using RC level information from off-momentum detectors
- Started with implementation of framework for tagged neutrons
 - Open questions about correct reduced cross section calculation
 - After discussion with Alex – will/should use average nucleon mass in a_T^2 calculation
- To-do/outlook:
 - Update a_T^2 calculation using average nucleon mass
 - Add full RC level framework, including scattered electron to tagged protons
 - Finalize tagged neutron framework
 - Cross-check with original simulation used for results in original publication

A. Jentsch, Z. Tu, C. Weiss: [Phys. Rev. C 104, 065205](#).



BACKUP

VARIABLES

- Scattered electron

- $Q^2 = -q^2 = -(p_{e,beam} - p_{e,scat})^2$

- $x = \frac{Q^2}{P \cdot q}$

- $x_{nucleon} = \frac{Q^2}{2P_{nucl} \cdot q} = \frac{x}{2 - \alpha_p}$

- Kinematic variables

- $y = \frac{p_d \cdot q}{p_d \cdot p_{e,beam}}$

- $1 - \epsilon = \frac{y^2}{1 + (1 - y)^2}$

- Light-cone momentum fraction:

- $\alpha_p \equiv \frac{2p_p^+}{p_d^+} = \frac{2(E_p + p_{z,p})}{E_d + p_{z,d}}$

- Proton transfer momentum

- $p_{T,p} = \sqrt{p_{x,p}^2 + p_{y,p}^2}$

- Fine structure constant

- $\alpha_{em} = \frac{1}{137}$

- Luminosity and cross section

- $L_{int} = \frac{N_{events}}{\sigma_{tot}} = \frac{N_{events}}{4.5 \cdot 10^{-5} \text{ mb}}$

- $\text{mb} = 2.568 \text{ GeV}^{-2}$