

# Computer Algebra Algorithms for Harmonic Sums and Polylogarithms and their Generalizations

Jakob Ablinger  
RISC, J. Kepler University Linz, Austria

joint work with  
J. Blümlein (DESY) and C. Schneider (RISC)

14. May 2011

- The following examples arise in the context of 2- and 3-loop massive single scale Feynman diagrams with operator insertion.
- These are related to the QCD anomalous dimensions and massive operator matrix elements.
- At 2-loop order all respective calculations are finished:

M. Buza, Y. Matiounine, J. Smith, R. Migneron, W.L. van Neerven, Nucl. Phys. **B472** (1996) 611;

I. Bierenbaum, J. Blümlein, S. Klein, Nucl. Phys. **B780** (2007) 40;


I. Bierenbaum, J. Blümlein, S. Klein, C. Schneider, Nucl.Phys. **B803** (2008) 1;

and lead to representations in terms of harmonic sums.

# Example 1: All N-Results for 3-Loop Ladder Graphs

Joint work with J. Blümlein (DESY), C. Schneider (RISC)  
A. Hasselhuhn (DESY), S. Klein (RWTH)

Consider, e.g., the diagram



(containing three massive fermion propagators)



Around 1000 sums have to be calculated

## A typical sum

$$\sum_{j=0}^{N-2} \sum_{s=1}^{j+1} \sum_{r=0}^{N+s-j-2} \sum_{\sigma=0}^{\infty} \frac{-2(-1)^{s+r} \binom{j+1}{s} \binom{-j+N+s-2}{r} (N-j)! (s-1)! \sigma! S_1(r+2)}{(N-r)(r+1)(r+2)(-j+N+\sigma+1)(-j+N+\sigma+2)(-j+N+s+\sigma)!}$$

## A typical sum

$$\sum_{j=0}^{N-2} \sum_{s=1}^{j+1} \sum_{r=0}^{N+s-j-2} \sum_{\sigma=0}^{\infty} \frac{-2(-1)^{s+r} \binom{j+1}{s} \binom{-j+N+s-2}{r} (N-j)! (s-1)! \sigma! S_1(r+2)}{(N-r)(r+1)(r+2)(-j+N+\sigma+1)(-j+N+\sigma+2)(-j+N+s+\sigma)!}$$

$$= \frac{(2N^2 + 6N + 5) S_{-2}(N)^2}{2(N+1)(N+2)} + S_{-2,-1,2}(N) + S_{-2,1,-2}(N)$$

$$+ \dots$$

where, e.g.,

$$S_{-2,1,-2}(N) = \sum_{i=1}^N \frac{(-1)^i \sum_{j=1}^i \frac{\sum_{k=1}^j (-1)^k}{k^2}}{i^2}$$

Vermaseren 98; Blümlein/Kurth 98

## A typical sum

$$\begin{aligned} & \sum_{j=0}^{N-2} \sum_{s=1}^{j+1} \sum_{r=0}^{N+s-j-2} \sum_{\sigma=0}^{\infty} \frac{-2(-1)^{s+r} \binom{j+1}{s} \binom{-j+N+s-2}{r} (N-j)! (s-1)! \sigma! S_1(r+2)}{(N-r)(r+1)(r+2)(-j+N+\sigma+1)(-j+N+\sigma+2)(-j+N+s+\sigma)!} \\ &= \frac{(2N^2 + 6N + 5) S_{-2}(N)^2}{2(N+1)(N+2)} + S_{-2,-1,2}(N) + S_{-2,1,-2}(N) \\ &+ \dots - S_{2,1,1,1}(-1, 2, \frac{1}{2}, -1; N) + S_{2,1,1,1}(1, \frac{1}{2}, 1, 2; N) \\ &+ \dots \end{aligned}$$

where, e.g.,

145 S-sums occur

$$S_{2,1,1,1}(1, \frac{1}{2}, 1, 2; N) = \sum_{i=1}^N \frac{\sum_{j=1}^i \frac{\left(\frac{1}{2}\right)^j \sum_{k=1}^j \frac{\sum_{l=1}^k 2^l}{l}}{j}}{i^2}$$

S. Moch, P. Uwer, S. Weinzierl 02

# Generalized Harmonic Sums

For  $a_i \in \mathbb{N}$  and  $x_i \in \mathbb{R} \setminus \{0\}$  we define

$$S_{a_1, \dots, a_k}(x_1, \dots, x_k; n) = \sum_{n \geq i_1 \geq i_2 \geq \dots \geq i_k \geq 1} \frac{x_1^{i_1}}{i_1^{a_1}} \cdots \frac{x_k^{i_k}}{i_k^{a_k}}.$$

# Generalized Harmonic Sums

For  $a_i \in \mathbb{N}$  and  $x_i \in \mathbb{R} \setminus \{0\}$  we define

$$S_{a_1, \dots, a_k}(x_1, \dots, x_k; n) = \sum_{n \geq i_1 \geq i_2 \geq \dots \geq i_k \geq 1} \frac{x_1^{i_1}}{i_1^{a_1}} \cdots \frac{x_k^{i_k}}{i_k^{a_k}}.$$

## Example

$$S_{2,3,1}\left(2, \frac{1}{2}, -1; n\right) = \sum_{k=1}^n \frac{2^k \sum_{j+1}^k \frac{\left(\frac{1}{2}\right)^j \sum_{i=1}^j \frac{(-1)^i}{i}}{j^3}}{k^2}$$

$$S_{2,3,1}(-1, 1, -1; n) = \sum_{k=1}^n \frac{(-1)^k \sum_{j+1}^k \frac{\sum_{i=1}^j \frac{(-1)^i}{i}}{j^3}}{k^2} = S_{-2,3,-1}(n)$$



# Algebraic Relations

$$\begin{aligned} S_{a_1, a_1, a_2}(x_1, x_2, x_3; n) = \\ S_{a_2}(x_3; n) S_{a_1, a_1}(x_1, x_2; n) + S_{a_1, a_1 + a_2}(x_1, x_2 x_3; n) - S_{a_1}(x_1; n) \\ S_{a_2, a_1}(x_3, x_2; n) - S_{a_2, 2a_1}(x_3, x_1 x_2; n) + S_{a_2, a_1, a_1}(x_3, x_2, x_1; n), \end{aligned}$$

$$\begin{aligned} S_{a_1, a_1, a_2}(x_1, x_3, x_2; n) = \\ S_{a_2}(x_2; n) S_{a_1, a_1}(x_1, x_3; n) + S_{a_1, a_1 + a_2}(x_1, x_2 x_3; n) - S_{a_1}(x_1; n) S_{a_2, a_1}(x_2, x_3; n) \\ - S_{a_2, 2a_1}(x_2, x_1 x_3; n) + S_{a_2, a_1, a_1}(x_2, x_3, x_1; n), \end{aligned}$$

$$\begin{aligned} S_{a_1, a_1, a_2}(x_2, x_1, x_3; n) = \\ -S_{a_2}(x_3; n) S_{a_1, a_1}(x_1, x_2; n) + S_{a_1}(x_2; n) S_{a_1, a_2}(x_1, x_3; n) + S_{2a_1, a_2}(x_1 x_2, x_3; n) \\ - S_{a_1 + a_2, a_1}(x_1 x_3, x_2; n) + S_{a_2, a_1, a_1}(x_3, x_1, x_2; n) \end{aligned}$$

- integral representation

$$S_{1,2,1} \left( 2, \frac{1}{2}, 1; n \right) = \int_0^1 \frac{1}{x_1 - 1} \int_1^{x_1} \frac{1}{x_2} \int_0^{x_2} \frac{1}{x_3 - 2} \int_2^{x_3} \frac{x_4^n - 1}{x_4 - 1} dx_4 dx_3 dx_2 dx_1$$

- integral representation

$$S_{1,2,1} \left( 2, \frac{1}{2}, 1; n \right) = \int_0^1 \frac{1}{x_1 - 1} \int_1^{x_1} \frac{1}{x_2} \int_0^{x_2} \frac{1}{x_3 - 2} \int_2^{x_3} \frac{x_4^n - 1}{x_4 - 1} dx_4 dx_3 dx_2 dx_1$$

- differentiation

$$\frac{\partial S_{1,2,1} \left( 2, \frac{1}{2}, 1; n \right)}{\partial n} = \int_0^1 \frac{1}{x_1 - 1} \int_1^{x_1} \frac{1}{x_2} \int_0^{x_2} \frac{1}{x_3 - 2} \int_2^{x_3} \frac{x_4^n \log(x_4)}{x_4 - 1} dx_4 dx_3 dx_2 dx_1$$

- integral representation

$$S_{1,2,1} \left( 2, \frac{1}{2}, 1; n \right) = \int_0^1 \frac{1}{x_1 - 1} \int_1^{x_1} \frac{1}{x_2} \int_0^{x_2} \frac{1}{x_3 - 2} \int_2^{x_3} \frac{x_4^n - 1}{x_4 - 1} dx_4 dx_3 dx_2 dx_1$$

- differentiation

$$\frac{\partial S_{1,2,1} \left( 2, \frac{1}{2}, 1; n \right)}{\partial n} = \int_0^1 \frac{1}{x_1 - 1} \int_1^{x_1} \frac{1}{x_2} \int_0^{x_2} \frac{1}{x_3 - 2} \int_2^{x_3} \frac{x_4^n \log(x_4)}{x_4 - 1} dx_4 dx_3 dx_2 dx_1$$

- integration of right hand side

$$\begin{aligned} \frac{\partial S_{1,2,1} \left( 2, \frac{1}{2}, 1; n \right)}{\partial n} = & -2H_{0,0,1,2,1}(1) - 4H_{0,0,2,1,1}(1) - H_{0,1,0,2,1}(1) - H_{0,1,2,0,1}(1) - 2H_{0,2,0,1,1}(1) \\ & - H_{0,2,1,0,1}(1) + (H_2(1)(H_{0,1,2}(1) + H_{0,2,1}(1)) + 2H_{0,0,1,2}(1) + 2H_{0,0,2,1}(1) \\ & + H_{0,1,0,2}(1))S_1(2; n) + S_2(\infty) \left( \frac{3(H_{0,1,2}(1) + H_{0,2,1}(1))}{2} + S_{1,2} \left( 2, \frac{1}{2}; n \right) \right) \\ & - H_2(1)S_{1,2,1} \left( 2, \frac{1}{2}, 1; n \right) - S_{-1}(\infty)S_{1,2,1} \left( 2, \frac{1}{2}, 1; n \right) - S_{1,2,2} \left( 2, \frac{1}{2}, 1; n \right) \\ & - 2S_{1,3,1} \left( 2, \frac{1}{2}, 1; n \right) - S_{2,2,1} \left( 2, \frac{1}{2}, 1; n \right) \end{aligned}$$

# Number of Basic S-Sums

We consider

$$S_{a_1, \dots, a_k}(x_1, \dots, x_k; n)$$

with  $x_i \in \{1, -1, 1/2, -1/2, 2, -2\}$ .

# Number of Basic S-Sums

We consider

$$S_{a_1, \dots, a_k}(x_1, \dots, x_k; n)$$

with  $x_i \in \{1, -1, 1/2, -1/2, 2, -2\}$ .

Example (Harmonic Sums ( $x_i \in \{1, -1\}$ ))

w	Number of							
	All	$N_A$	$N_D$	$N_H$	$N_{AD}$	$N_{AH}$	$N_{DH}$	$N_{ADH}$
1	2	2	2	1	2	1	1	1
2	6	3	4	4	1	2	3	1
3	18	8	12	14	5	6	10	4
4	54	18	36	46	10	15	32	9
5	162	48	108	146	30	42	100	27
6	486	116	324	454	68	107	308	65
7	1458	312	972	1394	196	294	940	187
8	4374	810	2916	4246	498	780	2852	486

# Number of Basic Harmonic Sums

For  $w \geq 2$  we have

$$A(w) = 2 \cdot 3^{w-1}$$

$$N_A(w) = \frac{1}{w} \sum_{d|w} \mu\left(\frac{w}{d}\right) 3^d$$

$$N_D(w) = 4 \cdot 3^{w-2}$$

$$N_H(w) = 2 \cdot 3^{w-1} - 2^{w-1}$$

$$N_{AD}(w) = \frac{1}{w} \sum_{d|w} \mu\left(\frac{w}{d}\right) 3^d - \frac{1}{w-1} \sum_{d|w-1} \mu\left(\frac{w-1}{d}\right) 3^d$$

$$N_{AH}(w) = \frac{1}{w} \sum_{d|w} \mu\left(\frac{w}{d}\right) [3^d - 2^d]$$

$$N_{DH}(w) = 4 \cdot 3^{w-2} - 2^{w-2}$$

$$N_{ADH}(w) = \frac{1}{w} \sum_{d|w} \mu\left(\frac{w}{d}\right) [3^d - 2^d] - \frac{1}{w-1} \sum_{d|w-1} \mu\left(\frac{w-1}{d}\right) [3^d - 2^d]$$

# Number of Basic S-Sums

We consider

$$S_{a_1, \dots, a_k}(x_1, \dots, x_k; n)$$

with  $x_i \in \{1, -1, 1/2, -1/2, 2, -2\}$ .



# Number of Basic S-Sums

We consider

$$S_{a_1, \dots, a_k}(x_1, \dots, x_k; n)$$


with  $x_i \in \{1, -1, 1/2, -1/2, 2, -2\}$ .

## Example

Each of the indices  $\{1/2, -1/2, 2, -2\}$  is allowed to appear just once in each sum.

Weight	Number of			
	All	$N_A$	$N_D$	$N_{AD}$
1	6	6	6	6
2	38	23	32	17
3	222	120	184	97
4	1206	654	984	543
5	6150	3536	4944	2882
6	29718	18280	23568	14744


# Example 1: continued



C. Schneider's Sigma.m

Around 1000 sums are calculated containing in total 533  $S$ -sums

# Example 1: continued



C. Schneider's Sigma.m

Around 1000 sums are calculated containing in total 533  $S$ -sums



HarmonicSum.m

After elimination the following sums remain:

$$S_{-4}(N), S_{-3}(N), S_{-2}(N), S_1(N), S_2(N), S_3(N), S_4(N), S_{-3,1}(N), \\ S_{-2,1}(N), S_{2,-2}(N), S_{2,1}(N), S_{3,1}(N), S_{-2,1,1}(N), S_{2,1,1}(N)$$

# Asymptotic Expansion of Harmonic Sums

We say that the function  $f : \mathbb{R} \rightarrow \mathbb{R}$  is expanded in an asymptotic series

$$f(x) \sim \sum_{k=0}^{\infty} \frac{a_k}{x^k}, \quad x \rightarrow \infty,$$

where  $a_k$  are constants, if for all  $K \geq 0$

$$R_K(x) = f(x) - \sum_{k=0}^K \frac{a_k}{x^k} = o\left(\frac{1}{x^K}\right), \quad x \rightarrow \infty.$$

Why do we need these expansions of harmonic sums?

E.g.,

- for limits of the form

$$\lim_{n \rightarrow \infty} n \left( S_2(n) - \zeta_2 - S_{2,2}(n) + \frac{7}{10} \zeta_2^2 \right)$$

- for the approximation of the values of analytic continued harmonic sums at the complex plane

$$S_{2,-3}(-20 + 10i)$$

$$S_{-1,3}(n) = (-1)^n \int_0^1 x^n \frac{H_{1,0,0}(x)}{1+x} dx + \text{const}$$

$\varphi(x)$ 

$$S_{-1,3}(n) = (-1)^n \int_0^1 x^n \overbrace{\frac{H_{1,0,0}(x)}{1+x}}^{\varphi(x)} dx + \text{const}$$

$$\varphi(x) \rightarrow \varphi(1-x) = \sum_{k=0}^{\infty} a_k x^k$$

$$S_{-1,3}(n) = (-1)^n \int_0^1 x^n \overbrace{\frac{H_{1,0,0}(x)}{1+x}} dx + \text{const}$$



$$\varphi(x) \longrightarrow \varphi(1-x) = \sum_{k=0}^{\infty} a_k x^k$$

$$S_{-1,3}(n) = (-1)^n \underbrace{\int_0^1 x^n \frac{\overbrace{H_{1,0,0}(x)}}{1+x} dx}_{+const}$$
$$\sum_{k=0}^{\infty} \frac{a_{k+1} k!}{n(n+1)\dots(n+k)}$$

$$\varphi(x) \longrightarrow \varphi(1-x) = \sum_{k=0}^{\infty} a_k x^k$$

$$S_{-1,3}(n) = (-1)^n \underbrace{\int_0^1 x^n \frac{\overbrace{H_{1,0,0}(x)}}{1+x} dx}_{+const}$$
$$\sum_{k=0}^{\infty} \frac{a_{k+1} k!}{n(n+1)\dots(n+k)} = \sum_{k=1}^{\infty} \frac{b_k}{n^k}$$

$$\varphi(x) \longrightarrow \varphi(1-x) = \sum_{k=0}^{\infty} a_k x^k$$

$$S_{-1,3}(n) = (-1)^n \underbrace{\int_0^1 x^n \frac{\overbrace{H_{1,0,0}(x)}}{1+x} dx}_{+const}$$

$$\sum_{k=0}^{\infty} \frac{a_{k+1} k!}{n(n+1)\dots(n+k)} = \sum_{k=1}^{\infty} \frac{b_k}{n^k}$$

$$b_1 = a_1$$

$$b_k = \sum_{l=0}^{k-2} (-1)^l S'_{k-l} a_{k-l} (k-l)!$$

$$\varphi(x) \rightarrow \varphi(1-x) = \sum_{k=0}^{\infty} a_k x^k$$

$$S_{-1,3}(n) = (-1)^n \underbrace{\int_0^1 x^n \frac{H_{1,0,0}(x)}{1+x} dx}_{+const}$$

$$\sum_{k=0}^{\infty} \frac{a_{k+1} k!}{n(n+1)\dots(n+k)} = \sum_{k=1}^{\infty} \frac{b_k}{n^k}$$

$$b_1 = a_1$$

$$b_k = \sum_{l=0}^{k-2} (-1)^l S'_{k-l} a_{k-l} (k-l)!$$

$$S_{-1,3}(n) \sim (-1)^n \left( -\frac{1}{4n^3} + \frac{5}{8n^4} - \frac{5}{8n^5} - \frac{5}{16n^6} \right) + \frac{3 \ln 2 z^3}{4}$$

$$+ (-1)^n \left( \frac{1}{2n} - \frac{1}{4n^2} + \frac{1}{8n^4} - \frac{1}{4n^6} \right) z^3 - \frac{19 z^2}{40}$$

# Example:

In[1]:= **SExpansion**[S[-1, 3, n], n, 10]

Out[1]=

$$(-1)^n \left( -\frac{1}{4n^3} + \frac{5}{8n^4} - \frac{5}{8n^5} - \frac{5}{16n^6} + \frac{31}{24n^7} + \frac{133}{96n^8} - \frac{169}{24n^9} - \frac{163}{16n^{10}} \right) + \frac{3 \ln 2 z^3}{4} + (-1)^n \left( \frac{1}{2n} - \frac{1}{4n^2} + \frac{1}{8n^4} - \frac{1}{4n^6} + \frac{17}{16n^8} - \frac{31}{4n^{10}} \right) z^3 - \frac{19 z^2}{40}$$


In[2]:= **GetApproximation**[S[-1,3,n], {-2.5, 2}]

Out[2]= -0.795096 - 0.105476 i


In[3]:= **HLimit**[n\*(S[2, n] - z^2 - S[2, 2, n] + 7\*z^2/10), n]

Out[3]= -1+z^2

We consider another diagram:




We consider another diagram:



↓  
Sigma ...

$$\begin{aligned}
 I_3 = & \frac{C_3}{(N+1)(N+2)(N+3)} \left\{ \frac{1}{6} S_1^3 + \frac{N^2 + 12N + 16}{2(N+1)(N+2)} S_1^2 + \frac{4(2N+3)}{(N+1)^2(N+2)} S_1 \right. \\
 & + \frac{8(2N+3)}{(N+1)^3(N+2)} + 2 \left[ -2^{N+3} + 3 - (-1)^N \right] \zeta_3 - (-1)^N S_{-3} + \left[ \frac{3N^2 + 40N + 56}{2(N+1)(N+2)} - \frac{1}{2} S_1 \right] S_2 \\
 & - \frac{3N+17}{3} S_3 - 2(-1)^N S_{-2,1} - (N+3) S_{2,1} + 2^{N+4} S_{1,2} \left( \frac{1}{2}, 1; N \right) \\
 & \left. + 2^{N+3} S_{1,1,1} \left( \frac{1}{2}, 1, 1; N \right) \right\} + O(\epsilon)
 \end{aligned}$$

We consider another diagram:



↓  
Sigma ...

$$\begin{aligned}
 I_3 = & \frac{C_3}{(N+1)(N+2)(N+3)} \left\{ \frac{1}{6} S_1^3 + \frac{N^2 + 12N + 16}{2(N+1)(N+2)} S_1^2 + \frac{4(2N+3)}{(N+1)^2(N+2)} S_1 \right. \\
 & + \frac{8(2N+3)}{(N+1)^3(N+2)} + 2 \left[ -2^{N+3} + 3 - (-1)^N \right] \zeta_3 - (-1)^N S_{-3} + \left[ \frac{3N^2 + 40N + 56}{2(N+1)(N+2)} - \frac{1}{2} S_1 \right] S_2 \\
 & - \frac{3N+17}{3} S_3 - 2(-1)^N S_{-2,1} - (N+3) S_{2,1} + 2^{N+4} S_{1,2} \left( \frac{1}{2}, 1; N \right) \\
 & \left. + 2^{N+3} S_{1,1,1} \left( \frac{1}{2}, 1, 1; N \right) \right\} + O(\epsilon)
 \end{aligned}$$

remaining sums:

$$S_1(n), S_2(n), S_3(n), S_{-3}(n), S_{2,1}(n), S_{-2,1}(n), S_{1,2} \left( \frac{1}{2}, 1; n \right), S_{1,1,1} \left( \frac{1}{2}, 1, 1; n \right)$$



# Asymptotic Expansion: Basic Idea

- again we start from the integral representation

$$S_{1,2,1}\left(\frac{1}{3}, \frac{1}{2}, 1; n\right) = \int_0^{\frac{1}{6}} \frac{1}{x_1 - \frac{1}{6}} \int_{\frac{1}{6}}^{x_1} \frac{1}{x_2} \int_0^{x_2} \frac{1}{x_3 - \frac{1}{3}} \int_{\frac{1}{3}}^{x_3} \frac{x_4^n - 1}{x_4 - 1} dx_4 dx_3 dx_2 dx_1$$

# Asymptotic Expansion: Basic Idea

- again we start from the integral representation

$$S_{1,2,1}\left(\frac{1}{3}, \frac{1}{2}, 1; n\right) = \int_0^{\frac{1}{6}} \frac{1}{x_1 - \frac{1}{6}} \int_{\frac{1}{6}}^{x_1} \frac{1}{x_2} \int_0^{x_2} \frac{1}{x_3 - \frac{1}{3}} \int_{\frac{1}{3}}^{x_3} \frac{x_4^n - 1}{x_4 - 1} dx_4 dx_3 dx_2 dx_1$$

- integration by parts leads to integrals of depth one:

$$\begin{aligned} S_{1,2,1}\left(\frac{1}{3}, \frac{1}{2}, 1; n\right) &= -\left(H_{\frac{1}{3}, 0, \frac{1}{6}}\left(\frac{1}{6}\right) - H_{\frac{1}{3}}\left(\frac{1}{6}\right)H_{0, \frac{1}{6}}\left(\frac{1}{6}\right)\right) \int_{\frac{1}{6}}^{\frac{1}{3}} \frac{x^n - 1}{x - 1} dx + H_{0, \frac{1}{6}}\left(\frac{1}{6}\right) \int_0^{\frac{1}{6}} \frac{x^n - 1}{x - 1} H_{\frac{1}{3}}(x) dx \\ &\quad - \int_0^{\frac{1}{6}} \frac{x^n - 1}{x - 1} H_{\frac{1}{3}, 0, \frac{1}{6}}(x) dx \end{aligned}$$

# Asymptotic Expansion: Basic Idea

- again we start from the integral representation

$$S_{1,2,1}\left(\frac{1}{3}, \frac{1}{2}, 1; n\right) = \int_0^{\frac{1}{6}} \frac{1}{x_1 - \frac{1}{6}} \int_{\frac{1}{6}}^{x_1} \frac{1}{x_2} \int_0^{x_2} \frac{1}{x_3 - \frac{1}{3}} \int_{\frac{1}{3}}^{x_3} \frac{x_4^n - 1}{x_4 - 1} dx_4 dx_3 dx_2 dx_1$$

- integration by parts leads to integrals of depth one:

$$\begin{aligned} S_{1,2,1}\left(\frac{1}{3}, \frac{1}{2}, 1; n\right) &= -\left(H_{\frac{1}{3}, 0, \frac{1}{6}}\left(\frac{1}{6}\right) - H_{\frac{1}{3}}\left(\frac{1}{6}\right)H_{0, \frac{1}{6}}\left(\frac{1}{6}\right)\right) \int_{\frac{1}{6}}^{\frac{1}{3}} \frac{x^n - 1}{x - 1} dx + H_{0, \frac{1}{6}}\left(\frac{1}{6}\right) \int_0^{\frac{1}{6}} \frac{x^n - 1}{x - 1} H_{\frac{1}{3}}(x) dx \\ &\quad - \int_0^{\frac{1}{6}} \frac{x^n - 1}{x - 1} H_{\frac{1}{3}, 0, \frac{1}{6}}(x) dx \end{aligned}$$

- as before for harmonic sums we can expand these integrals and get

$$\begin{aligned} \int_{\frac{1}{6}}^{\frac{1}{3}} \frac{x^n - 1}{x - 1} dx &\sim -H_1\left(\frac{1}{6}\right) + H_1\left(\frac{1}{3}\right) + 3^{-n} \left( -\frac{15}{n^5} + \frac{33}{8n^4} - \frac{3}{2n^3} + \frac{3}{4n^2} - 2^{-n} \left( -\frac{4074}{3125n^5} + \frac{366}{625n^4} \right. \right. \\ &\quad \left. \left. - \frac{42}{125n^3} + \frac{6}{25n^2} - \frac{1}{5n} \right) - \frac{1}{2n} \right) \\ \int_0^{\frac{1}{6}} \frac{x^n - 1}{x - 1} H_{\frac{1}{3}}(x) dx &\sim H_{1, \frac{1}{3}}\left(\frac{1}{6}\right) + 6^{-n} \left( \log(2) \left( -\frac{4074}{3125n^5} + \frac{366}{625n^4} - \frac{42}{125n^3} + \frac{6}{25n^2} - \frac{1}{5n} \right) - \frac{10834}{625n^5} \right. \\ &\quad \left. + \frac{456}{125n^4} - \frac{22}{25n^3} + \frac{1}{5n^2} \right) \end{aligned}$$

$$\begin{aligned}
S_{1,2,1} \left( \frac{1}{3}, \frac{1}{2}, 1; n \right) &\sim \\
S_{1,2,1} \left( \frac{1}{3}, \frac{1}{2}, 1; \infty \right) &+ 3^{-n} \left( 2^{-n} \left( L(n) \left( \frac{42H_{0,2}(1)}{125n^3} - \frac{6H_{0,2}(1)}{25n^2} + \frac{H_{0,2}(1)}{5n} + \log(2)^2 \left( \frac{21}{125n^3} - \frac{3}{25n^2} + \frac{1}{10n} \right) \right. \right. \right. \\
&- \frac{21\zeta_2}{125n^3} + \frac{1}{5n^3} + \frac{3\zeta_2}{25n^2} - \frac{\zeta_2}{10n} \left. \right) + \log(2) \left( -\frac{42H_{0,2}(1)}{125n^3} + \frac{6H_{0,2}(1)}{25n^2} - \frac{H_{0,2}(1)}{5n} \right) + \frac{42H_2(1)H_{0,2}(1)}{125n^3} \\
&- \frac{2461H_{0,2}(1)}{1500n^3} + \frac{126H_{0,0,2}(1)}{125n^3} - \frac{42(H_{0,0,2}(1) + H_{0,2,2}(1))}{125n^3} - \frac{42(H_{0,0,2}(1) + H_{0,6,2}(1))}{125n^3} - \frac{42(H_{0,0,2}(1) + H_{2,0,2}(1))}{125n^3} \\
&- \frac{42(H_{0,0,2}(1) + H_{6,0,2}(1))}{125n^3} - \frac{42H_{6,2,0}(1)}{125n^3} - \frac{6H_2(1)H_{0,2}(1)}{25n^2} + \frac{27H_{0,2}(1)}{50n^2} - \frac{18H_{0,0,2}(1)}{25n^2} + \frac{6(H_{0,0,2}(1) + H_{0,2,2}(1))}{25n^2} \\
&+ \frac{6(H_{0,0,2}(1) + H_{0,6,2}(1))}{25n^2} + \frac{6(H_{0,0,2}(1) + H_{2,0,2}(1))}{25n^2} + \frac{6(H_{0,0,2}(1) + H_{6,0,2}(1))}{25n^2} + \frac{6H_{6,2,0}(1)}{25n^2} \\
&+ \frac{H_2(1)H_{0,2}(1)}{5n} + \frac{3H_{0,0,2}(1)}{5n} - \frac{H_{0,0,2}(1) + H_{0,2,2}(1)}{5n} - \frac{H_{0,0,2}(1) + H_{0,6,2}(1)}{5n} - \frac{H_{0,0,2}(1) + H_{2,0,2}(1)}{5n} \\
&- \frac{H_{0,0,2}(1) + H_{6,0,2}(1)}{5n} - \frac{H_{6,2,0}(1)}{5n} + \log(2)^3 \left( -\frac{7}{125n^3} + \frac{1}{25n^2} - \frac{1}{30n} \right) + \log(2)^2 \left( \frac{27}{100n^2} - \frac{2461}{3000n^3} \right) \\
&+ \frac{2461\zeta_2}{3000n^3} + \frac{63\zeta_3}{250n^3} - \frac{27\zeta_2}{100n^2} - \frac{9\zeta_3}{50n^2} + \frac{3\zeta_3}{20n} \left. \right) - \frac{3H_2(1)H_{0,2}(1)}{2n^3} - \frac{9H_{0,0,2}(1)}{2n^3} + \frac{3(H_{0,0,2}(1) + H_{0,2,2}(1))}{2n^3} \\
&+ \frac{3(H_{0,0,2}(1) + H_{2,0,2}(1))}{2n^3} + \frac{3H_2(1)H_{0,2}(1)}{4n^2} + \frac{9H_{0,0,2}(1)}{4n^2} - \frac{3(H_{0,0,2}(1) + H_{0,2,2}(1))}{4n^2} - \frac{3(H_{0,0,2}(1) + H_{2,0,2}(1))}{4n^2} \\
&- \frac{H_2(1)H_{0,2}(1)}{2n} - \frac{3H_{0,0,2}(1)}{2n} + \frac{H_{0,0,2}(1) + H_{0,2,2}(1)}{2n} + \frac{H_{0,0,2}(1) + H_{2,0,2}(1)}{2n} \left. \right)
\end{aligned}$$

# Cyclotomic Harmonic Sums

$$S_{\{a_1, b_1, c_1\}, \dots, \{a_l, b_l, c_l\}}(s_1, \dots, s_l; n) = \sum_{k_1=1}^n \frac{s_1^{k_1}}{(a_1 k_1 + b_1)^{c_1}} S_{\{a_2, b_2, c_2\}; \dots; \{a_l, b_l, c_l\}}(s_2, \dots, s_l; k_1),$$

with  $S_\emptyset = 1$  and  $N, a_i, c_i \in \mathbb{N}_+, b_i \in \mathbb{N}, s_i = \pm 1, a_i > b_i$ .

# Cyclotomic Harmonic Sums

$$S_{\{a_1, b_1, c_1\}, \dots, \{a_l, b_l, c_l\}}(s_1, \dots, s_l; n) = \sum_{k_1=1}^n \frac{s_1^{k_1}}{(a_1 k_1 + b_1)^{c_1}} S_{\{a_2, b_2, c_2\}; \dots; \{a_l, b_l, c_l\}}(s_2, \dots, s_l; k_1),$$

with  $S_\emptyset = 1$  and  $N, a_i, c_i \in \mathbb{N}_+, b_i \in \mathbb{N}, s_i = \pm 1, a_i > b_i$ .

- algebraic relations

# Cyclotomic Harmonic Sums

$$S_{\{a_1, b_1, c_1\}, \dots, \{a_l, b_l, c_l\}}(s_1, \dots, s_l; n) = \sum_{k_1=1}^n \frac{s_1^{k_1}}{(a_1 k_1 + b_1)^{c_1}} S_{\{a_2, b_2, c_2\}; \dots; \{a_l, b_l, c_l\}}(s_2, \dots, s_l; k_1),$$

with  $S_\emptyset = 1$  and  $N, a_i, c_i \in \mathbb{N}_+, b_i \in \mathbb{N}, s_i = \pm 1, a_i > b_i$ .

- algebraic relations
- differential relations

# Cyclotomic Harmonic Sums

$$S_{\{a_1, b_1, c_1\}, \dots, \{a_l, b_l, c_l\}}(s_1, \dots, s_l; n) = \sum_{k_1=1}^n \frac{s_1^{k_1}}{(a_1 k_1 + b_1)^{c_1}} S_{\{a_2, b_2, c_2\}; \dots; \{a_l, b_l, c_l\}}(s_2, \dots, s_l; k_1),$$

with  $S_\emptyset = 1$  and  $N, a_i, c_i \in \mathbb{N}_+, b_i \in \mathbb{N}, s_i = \pm 1, a_i > b_i$ .

- algebraic relations
- differential relations
- multiple argument relations



# Cyclotomic Harmonic Sums

$$S_{\{a_1, b_1, c_1\}, \dots, \{a_l, b_l, c_l\}}(s_1, \dots, s_l; n) = \sum_{k_1=1}^n \frac{s_1^{k_1}}{(a_1 k_1 + b_1)^{c_1}} S_{\{a_2, b_2, c_2\}; \dots; \{a_l, b_l, c_l\}}(s_2, \dots, s_l; k_1),$$

with  $S_\emptyset = 1$  and  $N, a_i, c_i \in \mathbb{N}_+, b_i \in \mathbb{N}, s_i = \pm 1, a_i > b_i$ .

- algebraic relations
- differential relations
- multiple argument relations
- asymptotic expansion


# Cyclotomic Harmonic Sums

$$S_{\{a_1, b_1, c_1\}, \dots, \{a_l, b_l, c_l\}}(s_1, \dots, s_l; n) = \sum_{k_1=1}^n \frac{s_1^{k_1}}{(a_1 k_1 + b_1)^{c_1}} S_{\{a_2, b_2, c_2\}; \dots; \{a_l, b_l, c_l\}}(s_2, \dots, s_l; k_1),$$

with  $S_\emptyset = 1$  and  $N, a_i, c_i \in \mathbb{N}_+, b_i \in \mathbb{N}, s_i = \pm 1, a_i > b_i$ .

- algebraic relations
- differential relations
- multiple argument relations
- asymptotic expansion
- allowing  $s_i \in \mathbb{R}_+$  leads to Cyclotomic S-Sums.

# Harmonic Sums and their Generalizations



# The Package HarmonicSums

The package `HarmonicSums` offers functions to

- find algebraic and structural relations of harmonic sums and their generalizations

# The Package HarmonicSums

The package `HarmonicSums` offers functions to

- find algebraic and structural relations of harmonic sums and their generalizations
- compute the Mellin transform of harmonic sums and their generalizations, this leads to harmonic polylogarithms and their generalizations

# The Package HarmonicSums

The package `HarmonicSums` offers functions to

- find algebraic and structural relations of harmonic sums and their generalizations
- compute the Mellin transform of harmonic sums and their generalizations, this leads to harmonic polylogarithms and their generalizations
- find algebraic and structural relations of harmonic polylogarithms and their generalizations

# The Package HarmonicSums

The package `HarmonicSums` offers functions to

- find algebraic and structural relations of harmonic sums and their generalizations
- compute the Mellin transform of harmonic sums and their generalizations, this leads to harmonic polylogarithms and their generalizations
- find algebraic and structural relations of harmonic polylogarithms and their generalizations
- apply algebraic and structural relations to harmonic sums

# The Package HarmonicSums

The package `HarmonicSums` offers functions to

- find algebraic and structural relations of harmonic sums and their generalizations
- compute the Mellin transform of harmonic sums and their generalizations, this leads to harmonic polylogarithms and their generalizations
- find algebraic and structural relations of harmonic polylogarithms and their generalizations
- apply algebraic and structural relations to harmonic sums
- calculate the asymptotic expansion of harmonic sums and their generalizations













# The Package HarmonicSums

The package `HarmonicSums` offers functions to

- find algebraic and structural relations of harmonic sums and their generalizations
- compute the Mellin transform of harmonic sums and their generalizations, this leads to harmonic polylogarithms and their generalizations
- find algebraic and structural relations of harmonic polylogarithms and their generalizations
- apply algebraic and structural relations to harmonic sums
- calculate the asymptotic expansion of harmonic sums and their generalizations
- perform several other tasks not mentioned in this talk

#### Related work:

-  J. Ablinger, J. Blümlein, and C. Schneider.  
Harmonic sums and polylogarithms generated by cyclotomic polynomials.  
2011.
-  J. Ablinger, J. Blümlein, and C. Schneider.  
In preparation.  
2011.
-  J. Blümlein.  
Algebraic relations between harmonic sums and associated quantities.  
*Comput. Phys. Commun.*, 159:19 [arXiv:hep-ph/0311046], 2004.
-  J. Blümlein.  
Structural relations of harmonic sums and mellin transforms at weight  $w=6$ .  
*Clay Math. J.*, e-Print: arXiv:0901.0837 [math-ph]:in print, 2009.
-  J. Blümlein.  
Structural relations of harmonic sums and mellin transforms up to weight  $w=5$ .  
*Comput. Phys. Commun.* 180:2218–2249, 180:2218–2249, 2009.
-  J. Blümlein, D.J. Broadhurst, and J.A.M. Vermaseren.  
The multiple zeta value data mine.  
*Comput. Phys. Commun.*, 181:582–625, 2010.
-  J. Blümlein, M. Kauer, S. Klein, and C. Schneider.  
From moments to functions in quantum chromodynamics.  
*Comput. Phys. Commun.*, e-Print: arXiv:0902.4091 [hep-ph], 2009.
-  M. Hoffman.  
Quasi-shuffle products.  
*J. Algebraic Combin.*, 11:49–68, 2000.
-  E. Remiddi and J.A.M. Vermaseren.  
Harmonic polylogarithms.  
*Int. J. Mod. Phys., A* 15:725 [arXiv:hep-ph/9905237], 2000.
-  J.A.M. Vermaseren.  
Harmonic sums, Mellin transforms and integrals.  
*Int. J. Mod. Phys., A* 14:2037–2076, 1999.