

Light Higgs bosons in SUSY cascades


Oscar Stål



Brookhaven Forum

20.10 2011

SM Higgs Boson limits



- Higgs with SM-like couplings still allowed for $M_H = 115-150 \text{ GeV}$



Light Higgs bosons I: MSSM

- LEP -> Lower limit on Higgs mass
- SM-like Higgs: $M_H \gtrsim 114 \text{ GeV}$
- SUSY -> two Higgs doublets
-> 3 neutral Higgs states H_i
- CP-conserving MSSM: H_i, A^0

$$\frac{C(H_i VV)}{C(H_i VV)_{\text{SM}}} \propto \xi_i \quad \sum_{i=1}^2 \xi_i^2 = 1$$

- Lightest Higgs mass

$$90 \text{ GeV} \lesssim M_h \lesssim 130 \text{ GeV}$$



Light Higgs bosons II: CP-violating MSSM

- Relaxing CP conservation, all 3 neutral Higgs bosons mix

$$\sum_{i=1}^3 \xi_i^2 = 1$$


- Benchmark scenario “CPX” shows hole in excluded region

$$M_{H_1} \sim 35\text{--}45 \text{ GeV}$$

- Such light H_1 opens for

$$H_2 \rightarrow H_1 H_1$$

-> Complicated 4f final states from SM-like Higgs



Light Higgs bosons III: The NMSSM

- The NMSSM extends the MSSM, solves μ -problem

$$W_{\text{MSSM}}^{(2)} \sim \mu \hat{H}_2 \cdot \hat{H}_1 \rightarrow \lambda \hat{S} \hat{H}_2 \cdot \hat{H}_1 + \frac{1}{3} \kappa \hat{S}^3$$

- Additional singlet \rightarrow 3 CP-even, 2 CP-odd Higgs states $\sum_{i=1}^3 \xi_i^2 = 1$

- If lightest Higgs is singlet dominated \rightarrow Avoid LEP bound

- Another interesting possibility is very light A_1 : $M_{A_1} < 2m_b$
 H_1 explains largest LEP excess: “Ideal” Higgs scenarios



Dermisek, Gunion

- Difficult to discover light A_1/H_1 at LHC, and also “ H_{SM} ” since $H_{\text{SM}} \rightarrow \phi\phi \rightarrow 4b, 4\tau, 2b2\tau$
 $\phi = A_1, H_1$

MSSM -> NMSSM – Light CP-even Higgs bosons

- The lightest NMSSM Higgs may be much lighter than for the corresponding MSSM scenario...

Constr. Mh-max



F. Mahmoudi, J. Rathsman, OS, L. Zeune, [1012.4490]

... provided the H_1 coupling to Z bosons is suppressed by a large singlet component $\xi_1^2 \ll 1$

- Reduced production in standard channels at LHC

Higgs bosons in SUSY cascades

$$\tilde{q} \rightarrow q\tilde{\chi}_i^0 \rightarrow q\tilde{\chi}_j^0 H_k \rightarrow \dots$$

- Alternative to standard production ($gg \rightarrow H$, VBF, etc.)
 - Strong cross sections
 - Hard scale for $H \rightarrow b\bar{b}$
 - Trigger on hard jet(s) + high MET


- CMS MC analysis of “LM5”

$$M_{H_1} \sim 116 \text{ GeV}$$

$$5\sigma \text{ with } 1.5 \text{ fb}^{-1} \text{ (14 TeV)}$$



- Boosting the bb analysis

[Kribs, Roy, Martin, Spannowsky,, \[0912.4731\], \[1006.1656\]](#)
[Gori, Schwaller, Wagner \[1103.4138\]](#)



Cascades for light Higgs scenarios

- Higgs modes can be **dominant** when kinematically accessible
Full 1-loop corrections to $\tilde{\chi}_i^0 \rightarrow \tilde{\chi}_j^0 H_k$ decays in CPV MSSM



Fowler, Weiglein, [0909.5165]

Bandyopadhyay, [1008.3339]

Kraml, Porod, [hep-ph/0507055]

Choi, Miller, Zerwas, [hep-ph/0407209]

Cheung, Hou, [0809.1122]

- Also in the NMSSM the relevant decays have been studied at the level of BR

Phenomenological LHC analysis

OS, G. Weiglein, [1108.0595]

- Starting point: NMSSM “P4” benchmark (light singlet H_1)...

A. Djouadi et al, [0801.4321]

... slightly modified to allow for varying lightest Higgs mass

Higgs sector parameters					
λ	0.6		κ	0.12	
$\tan \beta$	2.6		μ_{eff}	-200	GeV
A_λ	-510	GeV	A_κ	0 - 300	GeV


Gaugino masses					
M_1	300	GeV	M_2	600	GeV
M_3	1000	GeV			

Trilinear couplings					
$A_t = A_b = A_\tau = 0$ GeV					

Soft scalar mass					
$M_{\text{SUSY}} = 750$ GeV, 1 TeV					

Modified P4 Higgs spectrum

NMSSMTools 2.3.5






■ Example working point

$$M_{H_1} = 40 \text{ GeV}$$

$$M_{H_2} = 116 \text{ GeV}$$




$$M_{A_1} = 184 \text{ GeV}$$

$$M_{H^{\pm}} \simeq 570 \text{ GeV} (\simeq M_{H_3} \simeq M_{A_2})$$



Neutralino sector

$\tilde{\chi}_1^0$	97.6 GeV	\tilde{S}, \tilde{H}
$\tilde{\chi}_2^0$	227 GeV	\tilde{H}
$\tilde{\chi}_3^0$	228 GeV	\tilde{H}
$\tilde{\chi}_4^0$	304 GeV	\tilde{B}
$\tilde{\chi}_5^0$	616 GeV	\tilde{W}



LHC analysis: MC simulation strategy

- SUSY-QCD matrix elements:

MadGraph/MadEvent

$$pp \rightarrow \tilde{g}\tilde{g}, \tilde{q}\tilde{q}, \tilde{g}\tilde{q}, \tilde{q}\tilde{\bar{q}}$$

- NMSSM resonance decays (external BR)

Pythia

$$\tilde{q} \rightarrow q\chi_i^0 \rightarrow q\chi_1^0 H_k \rightarrow q\chi_1^0 b\bar{b},$$

$$n_j \geq 1, \quad n_b \geq 2,$$

$$\tilde{g} \rightarrow g\tilde{q} \rightarrow gq\chi_i^0 \rightarrow gq\chi_1^0 H_k \rightarrow gq\chi_1^0 b\bar{b},$$

$$n_j \geq 2, \quad n_b \geq 2$$

cascades may include charginos, W, Z, ...

- ISR, FSR, Fragmentation, MPI

Pythia

- Energy Smearing, fast 'ATLAS' detector simulation

Delphes

$$P(\text{btag}|b) = 60\% \quad P(\text{btag}|c) = 10\% \quad P(\text{btag}|q) = 1\%$$





Squark/Gluino production

- Production of colored sparticles unchanged wrt MSSM

$$pp \rightarrow \tilde{g}\tilde{g}, \tilde{q}\tilde{q}, \tilde{g}\tilde{q}, \tilde{q}\tilde{\bar{q}}$$

- (S)QCD corrections sizable -> NLO





W. Beenakker, R. Höpker, M. Spira, P. M. Zerwas, [hep-ph/9610490]
T. Plehn, Prospino

Masses (GeV)		σ_{LO} (pb)					σ_{NLO} (pb)				
$M_{\tilde{g}}$	$M_{\tilde{q}}$	$\tilde{g}\tilde{g}$	$\tilde{q}\tilde{q}$	$\tilde{g}\tilde{q}$	$\tilde{q}\tilde{\bar{q}}$	Σ	$\tilde{g}\tilde{g}$	$\tilde{q}\tilde{q}$	$\tilde{g}\tilde{q}$	$\tilde{q}\tilde{\bar{q}}$	Σ
$\sqrt{s} = 7 \text{ TeV}$											
750	750	0.03	0.23	0.25	0.05	0.56	0.07	0.27	0.39	0.08	0.82
1000	750	0.002	0.19	0.06	0.05	0.31	0.006	0.21	0.10	0.07	0.39 
1000	1000	0.001	0.03	0.02	0.004	0.06	0.005	0.04	0.04	0.006	0.08 
$\sqrt{s} = 14 \text{ TeV}$											
750	750	1.18	1.67	5.20	1.06	9.11	2.21	2.06	6.78	1.53	12.6
1000	750	0.15	1.41	1.86	0.96	4.38	0.32	1.59	2.44	1.34	5.69 
1000	1000	0.14	0.42	0.87	0.18	1.61	0.31	0.51	1.19	0.26	2.27 
1500	1500	0.01	0.04	0.05	0.01	0.10	0.01	0.05	0.07	0.02	0.15

Kinematic distributions

$$M_{\text{SUSY}} = 750 \text{ GeV}$$

All histograms normalized to unity



Results for LHC@7 TeV

- Cuts-based analysis to suppress SM tt background
- ΔR cut selects b-jet pairs from H_1
SUSY BG suppressed by N_b



$$N_{\text{jets}} \geq 4 \quad N_b \geq 2$$

$$p_T(\text{jet1}) > 250 \text{ GeV}$$

$$p_T(\text{jet2}) > 100 \text{ GeV}$$

$$E_T^{\text{miss}} > 150 \text{ GeV}$$

$$\min \Delta R(b\bar{b}) < 1.5$$



Outlook: LHC@14 TeV

- Statistics improve greatly
- Clear signal also for 1 TeV squark case



$$N_{\text{jets}} \geq 4 \quad N_b \geq 2$$

$$p_T(\text{jet1}) > 250 \text{ GeV}$$

$$p_T(\text{jet2}) > 100 \text{ GeV}$$

$$E_T^{\text{miss}} > 200 \text{ GeV}$$

$$\min \Delta R(b\bar{b}) < 1.2$$



Conclusions

- Light Higgs bosons ($M_H < M_Z$) still compatible with constraints in extended models
- The NMSSM is an appealing scenario where this can be realized
- One method to search for these difficult light Higgses at the LHC is through production in SUSY cascades
- We have performed MC study of the light NMSSM Higgs case:
 - > Still meaningful to do Higgs searches in SUSY cascades at 7 TeV with squarks and/or gluinos close to present bounds
 - > Prospects at 14 TeV promising for much higher masses