

$t\bar{t}$ Forward-Backward Asymmetry from Loops of New Strongly Coupled Quarks

Thomas McElmurry
BNL → University of Rochester

H. Davoudiasl, T.M., A. Soni, arXiv:1108.1173

Brookhaven Forum
Upton, New York
2011-10-19

Introduction

Forward-backward asymmetry

- ▶ Choose Forward and Backward directions along beam axis.
- ▶ Define $t\bar{t}$ events as F or B depending on whether the t is found in the F or B hemisphere.
- ▶ Asymmetry:

$$A \equiv \frac{N_F - N_B}{N_F + N_B}$$

- ▶ To see an asymmetry, one needs an asymmetric initial state:
 - ▶ At a pp collider, one proton is as good as the other.
 - ▶ At a $p\bar{p}$ collider, gluon and sea quark PDFs are the same in p as in \bar{p} .
 - ▶ We need a $u\bar{u}$ or $d\bar{d}$ initial state.

Forward-backward asymmetry

at the Tevatron

- ▶ The Tevatron is a good place to measure the asymmetry.
 - ▶ It's a $p\bar{p}$ collider.
 - ▶ Protons go clockwise (\equiv Forward).
 - ▶ Antiprotons go anticlockwise (\equiv Backward).
 - ▶ $t\bar{t}$ production is dominated by the $q\bar{q}$ channel.
- ▶ The value of A depends on the frame of reference.
 - ▶ Lab frame? Center of mass?
 - ▶ We'll study the asymmetry in the $t\bar{t}$ rest frame: $A^{t\bar{t}}$.

Forward-backward asymmetry

in the Standard Model

- ▶ At leading order, $A = 0$ due to charge-conjugation symmetry
- ▶ NLO QCD permits an asymmetry due to interference between C -even and C -odd amplitudes
 - ▶ Virtual corrections give a positive asymmetry
 - ▶ Real radiation gives a negative asymmetry
 - ▶ Net result: $A^{t\bar{t}} \approx 0.05$. [Kühn, Rodrigo 1998]

Tevatron results

[CDF, Phys. Rev. D83, 112003 (2011); D0, arXiv:1107.4995]

- ▶ CDF measures $A^{t\bar{t}} = 0.158 \pm 0.075 \dots$
 - ▶ ... with a significant dependence on invariant mass:
 - ▶ $A^{t\bar{t}}(M_{t\bar{t}} < 450 \text{ GeV}) = -0.116 \pm 0.153$
 - ▶ $A^{t\bar{t}}(M_{t\bar{t}} > 450 \text{ GeV}) = 0.475 \pm 0.114$
 - ▶ ... and on separation in rapidity:
 - ▶ $A^{t\bar{t}}(|\Delta y_{t\bar{t}}| < 1) = 0.026 \pm 0.118$
 - ▶ $A^{t\bar{t}}(|\Delta y_{t\bar{t}}| > 1) = 0.611 \pm 0.256$
- ▶ D0 measures $A^{t\bar{t}} = 0.196 \pm 0.065 \dots$
 - ▶ ... without the strong enhancement at high $M_{t\bar{t}}$ and $|\Delta y_{t\bar{t}}|$.

The theorists' challenge

How to explain the observed asymmetry without changing the total cross section or violating other constraints?

- ▶ Various ideas:
 - ▶ Flavor-changing Z' ...
 - ▶ New scalar in the t -channel ...
 - ▶ Axigluons ...
- ▶ A new idea: $A^{t\bar{t}}$ arises from loop diagrams involving new heavy quarks.

A model

- ▶ New heavy quarks t' and b' , with mass m_Q
- ▶ New scalars ϕ^0, ϕ^\pm , with mass m_ϕ
- ▶ Flavor-changing interactions $\phi \bar{q} Q$

$$\mathcal{L} \supset \lambda_{ut'} \phi^0 \bar{u} t' + \lambda_{ub'} \phi^+ \bar{u} b' + \lambda_{dt'} \phi^- \bar{d} t' + \lambda_{db'} \phi^0 \bar{d} b' \\ + \lambda_{tt'} \phi^0 \bar{t} t' + \lambda_{tb'} \phi^+ \bar{t} b' + \text{H.C.}$$

Some of these Yukawa couplings will need to be strong.

A model

- ▶ New heavy quarks t' and b' , with mass m_Q
- ▶ New scalars ϕ^0, ϕ^\pm , with mass m_ϕ
- ▶ Flavor-changing interactions $\phi \bar{q} Q$


$$\mathcal{L} \supset \lambda_u (\phi^0 \bar{u} t' + \phi^+ \bar{u} b') + \lambda_d (\phi^- \bar{d} t' + \phi^0 \bar{d} b') \\ + \lambda_t (\phi^0 \bar{t} t' + \phi^+ \bar{t} b') + \text{H.C.}$$

Some of these Yukawa couplings will need to be strong.

New diagrams

Boxes!

- ▶ The new particles appear in box diagrams ...




- ▶ ... which interfere with the LO diagram and produce an asymmetry.

New diagrams

Triangles!

- ▶ We also have vertex corrections ...



- ▶ ... analogous to SM diagrams involving the Higgs.

[Stange, Willenbrock 1993]

- ▶ These diagrams do not produce an asymmetry, but do affect the total cross section.

Computation

- ▶ So we have some loops to compute.
- ▶ Evaluate the traces by hand . . .
- ▶ . . . and check using **QGRAF** and **FORM**.
- ▶ Cancel UV singularities analytically.
- ▶ Integrate numerically over Feynman parameters and phase space.

Asymmetry


Can we match the CDF measurement?

For what values of λ_u , λ_d , λ_t , m_Q , m_ϕ do we pass these six criteria?

- ▶ Agreement with the CDF inclusive asymmetry, within 2σ
- ▶ and with the high and low $M_{t\bar{t}}$ bins, within 2σ
- ▶ and with the high and low $|\Delta y_{t\bar{t}}|$ bins, within 2σ
- ▶ and with the SM total cross section, within 30%

Asymmetry

Can we match the CDF measurement?




Cross-hatched: $\lambda_U = 1$, $\lambda_D = 3$, $\lambda_t = 5$

Shaded: $\lambda_U = 1$, $\lambda_D = 2$, $\lambda_t = 6$

Hatched: $\lambda_U = 0$, $\lambda_D = 3.5$, $\lambda_t = 4.5$

Asymmetry

Can we match the CDF measurement? Yes.



Cross-hatched: $\lambda_U = 1, \lambda_D = 3, \lambda_t = 5$

Shaded: $\lambda_U = 1, \lambda_D = 2, \lambda_t = 6$


Hatched: $\lambda_U = 0, \lambda_D = 3.5, \lambda_t = 4.5$

Predictions and constraints

- ▶ This model predicts new heavy quarks t' and b' .
 - ▶ Experimental searches rule out e.g. $t' \rightarrow Wb$ with $m_{t'} \lesssim 450$ GeV. [CMS 2011]
 - ▶ These bounds do not necessarily apply to our model, since we can have decay modes such as $t' \rightarrow t\phi \rightarrow Wbq\bar{q}$.
- ▶ We can produce same-sign top pairs: $uu \rightarrow tt$.
 - ▶ We estimate that current constraints [CMS, JHEP 1108, 005 (2011)] do not exclude our model if $m_Q \gtrsim 400$ GeV.
 - ▶ We can also evade the bounds by choosing couplings such that λ_u is small.

Predictions and constraints

- ▶ Our model allows exotic single-top production.



- ▶ This can be suppressed by $\lambda_u \rightarrow 0$.
- ▶ Or, if $\lambda_{ut'} \approx -\lambda_{ub'}$, the two diagrams cancel.

Discussion

- ▶ Strongly coupled heavy quarks appearing in loops can produce an $A^{t\bar{t}}$ consistent with the CDF results.
- ▶ The D0 results can also be accommodated with smaller couplings.
- ▶ Various predictions and possibilities for the LHC:
 - ▶ Discovery of t', b'
 - ▶ Decay modes such as $t' \rightarrow t\phi$
 - ▶ Single t', b' production
 - ▶ Forward-backward asymmetry in $t'\bar{t}'$ and $b'\bar{b}'$ production