Neutrinos II

Mary Bishai Brookhaven National Laboratory

MeV Neutrinos from Reactors
Exercises
Daya Bay

GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments

GeV-PeV Neutrinos

Big Bang Neutrinos

ν Applications

Conclusions

Neutrinos II

Experimental Survey
Physics Summer School 2018

Mary Bishai
Brookhaven National Laboratory

July 12^{th}, 2018

Sources of Neutrinos

Neutrinos II

Mary Bishai Brookhaven National Laboratory

MeV Neutrinos from Reactors

Exercises

Daya Bay

GeV Neutrinos from Accelerators

Exercises

Current Experiments

Future Experiments


GeV-PeV Neutrinos

Big Bang Neutrinos

ν Applications

Conclusions

Big Bang



10^{-4} eV

$300/\mathrm{cm}^3$


Reactors



few MeV

$10^{21}/\mathrm{GW}_{th}/\mathrm{s}$


Sun



0.1-14 MeV

$10^{10}/\mathrm{cm}^2/\mathrm{s}$


SuperNova



$\sim$ 10 MeV

$10^{9}/\mathrm{cm}^2/\mathrm{s}$

Atmosphere



$\sim$ 1 GeV

$\mathrm{few}/\mathrm{cm}^2/\mathrm{s}$


Accelerators



1-20 GeV

$10^5/\mathrm{cm}^2/\mathrm{s}$ (at 1km)

Extragalactic



TeV-PeV

varies

Neutrinos II

Mary Bishai Brookhaven National Laboratory

MeV Neutrinos from Reactors
Exercises
Daya Bay

GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments

GeV-PeV Neutrinos

Big Bang Neutrinos

ν Applications

Conclusions

Neutrino Experiments

- **MeV scale Neutrinos: The Daya Bay Reactor Experiment**
- **GeV scale Neutrinos: The T2K, NOνA and DUNE experiments**
- **TeV-PeV scale Neutrinos: The IceCUBE Experiment NEW RESULT TODAY**
- **Big Bang Neutrinos: The PTOLEMY Experiment**

ν: A Truly Elusive Particle!

Neutrinos II

Mary Bishai Brookhaven National Laboratory

MeV Neutrinos from Reactors
Exercises
Daya Bay

GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments

GeV-PeV Neutrinos

Big Bang Neutrinos

ν Applications

Conclusions

Reines and Cowan were the first to estimate the interaction strength of neutrinos. **The cross-section is $\sigma \sim 10^{-43}\mathrm{cm}^2$ per nucleon (N = n or p).**

$$\nu \text{ mean free path} = \frac{1}{\sigma \times \text{number of nucleons per cm}^3}$$

ν Exercise: What is the mean free path of a neutrino in lead?

(use Table of atomic and nuclear properties)

ν: A Truly Elusive Particle!

Neutrinos II

Mary Bishai Brookhaven National Laboratory

MeV Neutrinos from Reactors
Exercises
Daya Bay

GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments

GeV-PeV Neutrinos

Big Bang Neutrinos

ν Applications

Conclusions

Reines and Cowan were the first to estimate the interaction strength of neutrinos. The cross-section is $\sigma \sim 10^{-43}\mathrm{cm}^2$ per nucleon (N = n or p).

$$\nu \text{ mean free path} = \frac{1}{\sigma \times \text{number of nucleons per cm}^3}$$

ν Exercise: What is the mean free path of a neutrino in lead?

(use Table of atomic and nuclear properties)

$$= \frac{1}{10^{-43}\mathrm{cm}^2 \times 11.4\mathrm{g/cm}^3 \times 6.02 \times 10^{23}\mathrm{nucleons/g}}$$

$$\approx 1.5 \times 10^{16} m$$

How many light years is that? How does it compare to the distance from the sun to the moon?

ν: A Truly Elusive Particle!

Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions

Reines and Cowan were the first to estimate the interaction strength of neutrinos. The cross-section is $\sigma \sim 10^{-43}\mathrm{cm}^2$ per nucleon (N = n or p).

$$\nu \text{ mean free path} = \frac{1}{\sigma \times \text{number of nucleons per cm}^3}$$

ν Exercise: What is the mean free path of a neutrino in lead?

(use Table of atomic and nuclear properties)

$$= \frac{1}{10^{-43}\mathrm{cm}^2 \times 11.4\mathrm{g/cm}^3 \times 6.02 \times 10^{23}\mathrm{nucleons/g}}$$

$$\approx 1.5 \times 10^{16} m$$

How many light years is that? How does it compare to the distance from the sun to the moon?

$$= \text{1.6 LIGHT YEARS OF LEAD}$$

$$= 100,000 \text{ distance earth to sun}$$

A proton has a mean free path of 10cm in lead

Reactor power and neutrinos


Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions

ν Exercise:

The following table shows the breakdown of energy released per fission from ^{235}U:

Fission fragment	Energy (MeV)
Fission products	175
$\langle 2.44 \rangle$ neutrons	5
γ from fission	7
γs and βs from *beta* decay	13
$\langle 6 \rangle$ neutrinos	10
Total	210

5% of a reactor's power is in neutrinos !



How many neutrinos are emitted per second from a 1 Gigawatt (thermal) reactor?


Reactor power and neutrinos

Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions

ν Exercise:

The following table shows the breakdown of energy released per fission from ^{235}U:

Fission fragment	Energy (MeV)
Fission products	175
$\langle 2.44 \rangle$ neutrons	5
γ from fission	7
γs and βs from *beta* decay	13
$\langle 6 \rangle$ neutrinos	10
Total	210



5% of a reactor's power is in neutrinos !

How many neutrinos are emitted per second from a 1 Gigawatt (thermal) reactor?

$$
\begin{aligned}
1 \times 10^9 \text{ Joules/sec} &= 6.242 \times 10^{18} \text{ GeV/sec} \\
&= 3 \times 10^{19} \text{ fissions/sec} \\
&\sim 2 \times 10^{20} \ \nu\text{/sec} \\
&= 1.6 \times 10^{13} / \text{m}^2 / \text{sec at 1 km}
\end{aligned}
$$

Reactor Power and Neutrinos

Neutrinos II

Mary Bishai Brookhaven National Laboratory

MeV Neutrinos from Reactors
Exercises
Daya Bay

GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments

GeV-PeV Neutrinos

Big Bang Neutrinos

ν Applications

Conclusions

ν Exercise:

Using the rate of neutrinos emitted from a reactor ($= 2 \times 10^{20}$/sec/GW) and the average cross-section of the inverse beta decay process ($\bar{\nu}_e + p \rightarrow e^+ + n$) is $\sigma = 10^{-43}\mathrm{cm}^2$/proton, what is the rate of neutrino interactions per day in a detector containing 100 tons of scintillator (CH_2) located 1km from a 1GW reactor? Note that the IBD process only happens on free protons (H)

Reactor Power and Neutrinos

Neutrinos II

Mary Bishai Brookhaven National Laboratory

MeV Neutrinos from Reactors
Exercises
Daya Bay

GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments

GeV-PeV Neutrinos

Big Bang Neutrinos

ν Applications

Conclusions

ν Exercise:

Using the rate of neutrinos emitted from a reactor ($= 2 \times 10^{20}$/sec/GW) and the average cross-section of the inverse beta decay process ($\bar{\nu}_e + p \rightarrow e^+ + n$) is $\sigma = 10^{-43}\mathrm{cm}^2$/proton, what is the rate of neutrino interactions per day in a detector containing 100 tons of scintillator (CH_2) located 1km from a 1GW reactor? Note that the IBD process only happens on free protons (H)

 # interactions/day = flux (ν/cm^2/day) $\times$ σ (cm^2/p) $\times$ protons/Nucleons $\times$ Nucleons/gram $\times$ 10^8 g/100tons

Reactor Power and Neutrinos

Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions

ν Exercise:

Using the rate of neutrinos emitted from a reactor ($= 2 \times 10^{20}$/sec/GW) and the average cross-section of the inverse beta decay process ($\bar{\nu}_e + p \rightarrow e^+ + n$) is $\sigma = 10^{-43}\mathrm{cm}^2$/proton, what is the rate of neutrino interactions per day in a detector containing 100 tons of scintillator (CH_2) located 1km from a 1GW reactor? Note that the IBD process only happens on free protons (H)

 # interactions/day = flux (ν/cm^2/day) $\times$ σ (cm^2/p) $\times$ protons/Nucleons $\times$ Nucleons/gram $\times$ 10^8 g/100tons


 # interactions/day = 118

Precision ν expt: need 1 GW nuclear reactor ($1B) + 100's tons

More Reactor $\bar{\nu}_e$: Measuring Oscillation Parameters

Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions

$$P_{\bar{\nu}_e \to \bar{\nu}_e} \approx 1 - \sin^2 2\theta_{13} \sin^2(\Delta m^2_{31} L/4E) - \cos^4 \theta_{13} \sin^2 2\theta_{12} \sin^2(\Delta m^2_{21} L/4E)$$



The Daya Bay Reactor Complex

Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrino from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Application
Conclusions





Reactor Specs:

Located 55km north-east of Hong Kong.

Initially: 2 cores at Daya Bay site + 2 cores at Ling Ao site = 11.6 GW_{th}

By 2011: 2 more cores at Ling Ao II site = 17.4 GW_{th} $\Rightarrow$ top five worldwide


1 $GW_{th} = 2 \times 10^{20} \bar{\nu}_e$/second

Deploy multiple near and far detectors

Reactor power uncertainties $< 0.1\%$

The Daya Bay Collaboration : 231 Collaborators

Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions



Asia (21)
Beijing Normal Univ., CNG, CIAE, Dongguan Polytechnic, ECUST, IHEP, Nanjing Univ., Nankai Univ., NCEPU, Shandong Univ., Shanghai Jiao Tong Univ., Shenzhen Univ., Tsinghua Univ., USTC, Xian Jiaotong Univ., Zhongshan Univ., Chinese Univ. of Hong Kong, Univ. of Hong Kong, National Chiao Tung Univ., National Taiwan Univ., National United Univ.

Europe (2)
Charles University, JINR Dubna


North America (17)
Brookhaven Natl Lab, CalTech, Illinois Institute of Technology, Iowa State, Lawrence Berkeley Natl Lab, Princeton, Rensselaer Polytechnic, Siena College, UC Berkeley, UCLA, Univ. of Cincinnati, Univ. of Houston, UIUC, Univ. of Wisconsin, Virginia Tech, William & Mary, Yale

South America (1)
Catholic Univ. of Chile

Detecting Neutrinos from the Daya Bay Reactors

Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions

The active target in each detector is liquid scintillator loaded with 0.1% Gd




- $\bar{\nu_e} + p \rightarrow n + e^+$
- $e^+ + e^- \rightarrow \gamma\gamma$ (2X 0.511 MeV $+ T_{e^+}$, prompt)
- $n + p \rightarrow D + \gamma$ (2.2 MeV, $\tau \sim 180\mu s$). OR
- $n + Gd \rightarrow Gd* \rightarrow Gd + \gamma$'s (8 MeV, $\tau \sim 28\mu s$).

$\Rightarrow$ delayed co-incidence of e^+ conversion and n-capture ($>$ 6 MeV)

with a specfic energy signature

The Daya Bay Experimental Apparatus

Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions




- **Multiple "identical" detectors at each site.**
- **Manual and multiple automated calibration systems per detector.**
- **Thick water shield to reduce cosmogenic and radiation bkgds.**

	DYB	LA	Far
Event rates/20T/day	840	740	90

Daya Bay Measurement of Non-zero θ_{13}

Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions




First to discover non-zero θ_{13} (2012) and currently most precise result:

$$\sin^2 2\theta_{13} = 0.084 \pm 0.005$$

Measured and predicted neutrino flux at DYB

Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions

—Predicted —Predicted with best fit value of θ_{13}




ν detectors can now monitor reactors!

CP Violation in PMNS (leptons) and CKM (quarks)

Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions

In 3-flavor mixing the degree of CP violation is determined by the Jarlskog invariant:

$$J_{CP}^{\rm PMNS} \equiv \frac{1}{8} \sin 2\theta_{12} \sin 2\theta_{13} \sin 2\theta_{23} \cos\theta_{13} \sin\delta_{\rm CP}.$$



(JHEP 11 (2014) 052, arXiv:1409.5439)

Given the current best-fit values of the ν mixing angles :

$$J_{CP}^{\rm PMNS} \approx 3 \times 10^{-2} \sin\delta_{\rm CP}.$$

For CKM (mixing among the 3 quark generations):

$$J_{CP}^{\rm CKM} \approx 3 \times 10^{-5},$$

despite the large value of $\delta_{CP}^{\rm CKM} \approx 70^\circ$.

$\nu_\mu \rightarrow \nu_e$ Oscillations in the 3-flavor ν SM

Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions

In the ν 3-flavor model matter/anti-matter asymmetries in neutrinos are best probed using $\nu_\mu/\bar{\nu}_\mu \rightarrow \nu_e/\bar{\nu}_e$ oscillations (or vice versa). With terms up to second order in $\alpha \equiv \Delta m^2_{21}/\Delta m^2_{31} = 0.03$ and $\sin^2\theta_{13} = 0.02$, (M. Freund. Phys. Rev. D 64, 053003):

$$P(\nu_\mu \rightarrow \nu_e) \cong P(\nu_e \rightarrow \nu_\mu) \cong \underbrace{P_0}_{\theta_{13}} + \underbrace{P_{\sin\delta}}_{\text{CP violating}} + \underbrace{P_{\cos\delta}}_{\text{CP conserving}} + \underbrace{P_3}_{\text{solar oscillation}}$$

where **for oscillations in vacuum:**

$$P_0 = \sin^2\theta_{23}\sin^2 2\theta_{13}\sin^2(\Delta),$$

$$P_{\sin\delta} = \alpha\, 8 J_{cp} \sin^3(\Delta),$$

$$P_{\cos\delta} = \alpha\, 8 J_{cp} \cot\delta_{CP} \cos\Delta \sin^2(\Delta),$$

$$P_3 = \alpha^2 \cos^2\theta_{23}\sin^2 2\theta_{12}\sin^2(\Delta),$$

where $\Delta = 1.27\Delta m^2_{31}(eV^2)L(km)/E(GeV)$

For $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$, $\underbrace{P_{\sin\delta} \rightarrow -P_{\sin\delta}}_{\text{CP asymmetry}}$,

$\nu_\mu \rightarrow \nu_e$ Oscillations in the 3-flavor ν SM

Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions

In the ν 3-flavor model matter/anti-matter asymmetries in neutrinos are best probed using $\nu_\mu/\bar{\nu}_\mu \rightarrow \nu_e/\bar{\nu}_e$ oscillations (or vice versa). With terms up to second order in $\alpha \equiv \Delta m^2_{21}/\Delta m^2_{31} = 0.03$ and $\sin^2\theta_{13} = 0.02$, (M. Freund. Phys. Rev. D 64, 053003):

$$P(\nu_\mu \rightarrow \nu_e) \cong P(\nu_e \rightarrow \nu_\mu) \cong \underbrace{P_0}_{\theta_{13}} + \underbrace{P_{\sin\delta}}_{\text{CP violating}} + \underbrace{P_{\cos\delta}}_{\text{CP conserving}} + \underbrace{P_3}_{\text{solar oscillation}}$$

where **for oscillations in matter with constant density:**

$$P_0 = \sin^2\theta_{23}\frac{\sin^2 2\theta_{13}}{(A-1)^2}\sin^2[(A-1)\Delta],$$

$$P_{\sin\delta} = \alpha\frac{8J_{cp}}{A(1-A)}\sin\Delta\sin(A\Delta)\sin[(1-A)\Delta],$$

$$P_{\cos\delta} = \alpha\frac{8J_{cp}\cot\delta_{\rm CP}}{A(1-A)}\cos\Delta\sin(A\Delta)\sin[(1-A)\Delta],$$

$$P_3 = \alpha^2\cos^2\theta_{23}\frac{\sin^2 2\theta_{12}}{A^2}\sin^2(A\Delta),$$

where $\Delta = 1.27\Delta m^2_{31}(eV^2)L(km)/E(GeV)$ and $A = \sqrt{2}G_F N_e 2E/\Delta m^2_{31}$.

For $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$, $\underbrace{P_{\sin\delta} \rightarrow -P_{\sin\delta}}_{\text{CP asymmetry}}$, $\underbrace{A \rightarrow -A}_{\text{matter asymmetry}}$

Osc. vs L/E

$\sin^2 2\theta_{13} = 0.09, \sin^2 \theta_{23} = 0.5, \Delta m^2_{31} = \pm 2.4 \times 10^{-3} \mathrm{eV}^2$

Neutrinos II

Mary Bishai Brookhaven National Laboratory

MeV Neutrinos from Reactors
Exercises
Daya Bay

GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments

GeV-PeV Neutrinos

Big Bang Neutrinos

ν Applications


Conclusions

ν Exercise: Use ROOT and reproduce the plots shown below

The $\nu_\mu \rightarrow \nu_e$ oscillation probability maxima occur at

$$\frac{L\ (\mathrm{km})}{E_n(\mathrm{GeV})} = \left(\frac{\pi}{2}\right) \frac{(2n-1)}{1.27 \times \Delta m^2_{31}(\mathrm{eV}^2)} \approx (2n-1) \times \frac{515\ \mathrm{km}}{\mathrm{GeV}}$$

Oscillations in vacuum - different terms ($\delta_{CP} = 0$)



Osc. vs L/E

$\sin^2 2\theta_{13} = 0.09, \sin^2 \theta_{23} = 0.5, \Delta m^2_{31} = \pm 2.4 \times 10^{-3} \mathrm{eV}^2$


Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions

ν Exercise: Use ROOT and reproduce the plots shown below

The $\nu_\mu \rightarrow \nu_e$ oscillation probability maxima occur at

$$\frac{L\ (\mathrm{km})}{E_n(\mathrm{GeV})} = \left(\frac{\pi}{2}\right) \frac{(2n-1)}{1.27 \times \Delta m^2_{31}(\mathrm{eV}^2)} \approx (2n-1) \times \frac{515\ \mathrm{km}}{\mathrm{GeV}}$$

Impact of δ_{CP} on oscillations in vacuum, $\Delta m^2_{31} > 0$ (NH)



Osc. vs L/E

$\sin^2 2\theta_{13} = 0.09, \sin^2 \theta_{23} = 0.5, \Delta m^2_{31} = \pm 2.4 \times 10^{-3} \mathrm{eV}^2$


Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions

ν Exercise: Use ROOT and reproduce the plots shown below

The $\nu_\mu \rightarrow \nu_e$ oscillation probability maxima occur at

$$\frac{L\ (\mathrm{km})}{E_n(\mathrm{GeV})} = \left(\frac{\pi}{2}\right) \frac{(2n-1)}{1.27 \times \Delta m^2_{31}(\mathrm{eV}^2)} \approx (2n-1) \times \frac{515\ \mathrm{km}}{\mathrm{GeV}}$$

Impact of δ_{CP} on oscillations in vacuum, $\Delta m^2_{31} < 0$ (IH)



Osc. vs L/E

$\sin^2 2\theta_{13} = 0.09, \sin^2 \theta_{23} = 0.5, \Delta m^2_{31} = \pm 2.4 \times 10^{-3} \mathrm{eV}^2$


Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions

ν Exercise: Use ROOT and reproduce the plots shown below

The $\nu_\mu \rightarrow \nu_e$ oscillation probability maxima occur at

$$\frac{L\ (\mathrm{km})}{E_n(\mathrm{GeV})} = \left(\frac{\pi}{2}\right) \frac{(2n-1)}{1.27 \times \Delta m^2_{31}(\mathrm{eV}^2)} \approx (2n-1) \times \frac{515\ \mathrm{km}}{\mathrm{GeV}}$$

Impact of matter effect on ν_μ oscillations ($\delta_{\mathrm{CP}} = 0$)



Osc. vs L/E

$\sin^2 2\theta_{13} = 0.09, \sin^2 \theta_{23} = 0.5, \Delta m^2_{31} = \pm 2.4 \times 10^{-3} \mathrm{eV}^2$


Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions

ν Exercise: Use ROOT and reproduce the plots shown below

The $\nu_\mu \rightarrow \nu_e$ oscillation probability maxima occur at

$$\frac{L\ (\mathrm{km})}{E_n(\mathrm{GeV})} = \left(\frac{\pi}{2}\right) \frac{(2n-1)}{1.27 \times \Delta m^2_{31}(\mathrm{eV}^2)} \approx (2n-1) \times \frac{515\ \mathrm{km}}{\mathrm{GeV}}$$

Impact of matter effect on $\bar{\nu}_\mu$ oscillations ($\delta_{\mathrm{CP}} = 0$)



Expected Appearance Signal Event Rates

Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions

ν Exercise: **The total number of electron neutrino appearance events expected for a given exposure from a muon neutrino source as a function of baseline is given as**

$$N_{\nu_e}^{\text{appear}}(L) = \int \Phi^{\nu_\mu}(E_\nu, L) \times P^{\nu_\mu \to \nu_e}(E_\nu, L) \times \sigma^{\nu_e}(E_\nu) dE_\nu$$

Assume the neutrino source produces a flux that is constant in energy and using only the dominant term in the probability(no matter effect)

$$\begin{aligned} \Phi^{\nu_\mu}(E_\nu, L) &\approx \frac{C}{L^2}, \quad C = \text{number of } \nu_\mu/\text{m}^2/\text{GeV}/\text{sec at 1 km} \\ P^{\nu_\mu \to \nu_e}(E_\nu, L) &\approx \underbrace{\sin^2\theta_{23} \sin^2 2\theta_{13} \sin^2(1.27\Delta m^2_{31} L/E_\nu)}_{P_0} \\ \sigma^{\nu_e}(E_\nu) &= 0.7 \times 10^{-42}(\text{m}^2/\text{GeV}/N) \times E_\nu, \quad E_\nu > 1 \text{ GeV} \end{aligned}$$

Prove that the rate of ν_e appearing integrated over a constant range of L/E is independent of baseline!

Expected Appearance Signal Event Rates

Neutrinos II

Mary Bishai Brookhaven National Laboratory

MeV Neutrinos from Reactors
Exercises
Daya Bay

GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments

GeV-PeV Neutrinos

Big Bang Neutrinos

ν Applications

Conclusions

$$N_{\nu_e}^{\text{appear}}(L) \propto \text{constant term} \times \int \frac{\sin^2(ax)}{x^3} dx,$$

$$x \equiv L/E_\nu, \; a \equiv 1.27 \Delta m^2_{31} \; \text{GeV}/(\text{eV}^2.\text{km})$$

ν Exercise:

$C \approx 1 \times 10^{17}$ $\nu_\mu/\text{m}^2/\text{GeV}/\text{yr}$ at 1 km (from 1MW accelerator)

$\sin^2 2\theta_{13} = 0.084$, $\sin^2 \theta_{23} = 0.5$, $\Delta m^2_{31} = 2.4 \times 10^{-3} \text{eV}^2$

Calculate the rate of ν_e events observed per kton of detector integrating over the region $x = 100$ km/GeV to 2000 km/GeV. Use ROOT to do the integral!

Expected Appearance Signal Event Rates

Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions

$$N_{\nu_e}^{\text{appear}}(L) \propto \text{constant term} \times \int \frac{\sin^2(ax)}{x^3} dx,$$

$$x \equiv L/E_\nu, \; a \equiv 1.27 \Delta m^2_{31} \; \text{GeV}/(\text{eV}^2.\text{km})$$

ν Exercise:

$C \approx 1 \times 10^{17} \; \nu_\mu/\text{m}^2/\text{GeV}/\text{yr}$ at 1 km (from 1MW accelerator)
$\sin^2 2\theta_{13} = 0.084$, $\sin^2 \theta_{23} = 0.5$, $\Delta m^2_{31} = 2.4 \times 10^{-3} \text{eV}^2$

Calculate the rate of ν_e events observed per kton of detector integrating over the region $x = 100$ km/GeV to 2000 km/GeV. Use ROOT to do the integral!

$$N_{\nu_e}^{\text{appear}}(L) \approx (2 \times 10^6 \text{events/kton/yr}) \cdot (\text{km/GeV})^2 \int_{x_0}^{x_1} \frac{\sin^2(ax)}{x^3} dx,$$

$$N_{\nu_e}^{\text{appear}}(L) \sim \mathcal{O}(20 - 30) \text{ events/kton/yr}$$

Neutrinos from Accelerators

Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions

To produce neutrinos from accelerators

$p^+ + A \rightarrow \pi^\pm + X, \quad \pi^\pm \rightarrow \mu^\pm + \nu_\mu/\bar{\nu}_\mu$

where A = Carbon (Graphite), Berillyium, Tungsten, X is other particles

ν Exercise: The Main Injector accelerator at Fermilab produces 4.86×10^{13} 120 GeV protons in a 10 microsecond pulse every 1.33 seconds to the NuMI beamline. What is the average power of the proton beam delivered in megawatts?

Neutrinos from Accelerators

Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions


To produce neutrinos from accelerators

$p^+ + A \rightarrow \pi^\pm + X, \quad \pi^\pm \rightarrow \mu^\pm + \nu_\mu/\bar{\nu}_\mu$

where A = Carbon (Graphite), Berillyium, Tungsten, X is other particles

ν Exercise: The Main Injector accelerator at Fermilab produces 4.86×10^{13} 120 GeV protons in a 10 microsecond pulse every 1.33 seconds to the NuMI beamline. What is the average power of the proton beam delivered in megawatts?


Power = 120 GeV $\times$ 4.86 10^{13} protons $\times$ 1.6 10^{-10} Joules/GeV $\times$ 1/1.33s = 702 kW




Neutrinos from Accelerators

Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions

The result of a FLUKA simulation of pion production from 120 GeV protons is shown below







ν Exercise: What fraction of 6 GeV pions on average will decay before reaching the end of an evacuated pipe 200m (675m) long? The π^+ rest mass and lifetime are 140 MeV and 26 ns

Neutrinos from Accelerators

Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions

The result of a FLUKA simulation of pion production from 120 GeV protons is shown below






ν Exercise: What fraction of 6 GeV pions on average will decay before reaching the end of an evacuated pipe 200m (675m) long? The π^+ rest mass and lifetime are 140 MeV and 26 ns


6 GeV π^+ lifetime: $\tau = \gamma\tau_0 = \frac{E}{m_0 c^2} \times 26\text{ns} = 1.1\text{ns}$, $c\tau = 334$ m

Neutrinos from Accelerators

Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions

The result of a FLUKA simulation of pion production from 120 GeV protons is shown below





ν Exercise: What fraction of 6 GeV pions on average will decay before reaching the end of an evacuated pipe 200m (675m) long? The π^+ rest mass and lifetime are 140 MeV and 26 ns


6 GeV π^+ lifetime: $\tau = \gamma\tau_0 = \frac{E}{m_0 c^2} \times 26\text{ns} = 1.1\text{ns}$, $c\tau = 334$ m

$F_{decays} = (1 - \exp^{-l/c\tau}) = 0.45(0.87)$


Neutrinos at the Main Injector

The longest baseline accel. ν expt in operation with highest power (700kW)

Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions



Fermi Natl. Lab., IL **Soudan Underground Lab, MN**







NuMI Horn 2 inner conductor
Radial field, $B \propto 1/r$

3T at 200 kA

Making Neutrinos and Anti-Neutrinos


Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions





Making Neutrinos and Anti-Neutrinos

Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions



The NOνA Experiment

Neutrinos II

Mary Bishai Brookhaven National Laboratory

MeV Neutrinos from Reactors
Exercises
Daya Bay


GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments


GeV-PeV Neutrinos


Big Bang Neutrinos

ν Applications

Conclusions







Neutrino Events in NOνA

Neutrinos II

Mary Bishai Brookhaven National Laboratory

MeV Neutrinos from Reactors

Exercises

Daya Bay

GeV Neutrinos from Accelerators

Exercises

Current Experiments


Future Experiments

GeV-PeV Neutrinos

Big Bang Neutrinos

ν Applications

Conclusions



Neutrino Events in NOνA

Neutrinos II

Mary Bishai Brookhaven National Laboratory

MeV Neutrinos from Reactors

Exercises

Daya Bay

GeV Neutrinos from Accelerators

Exercises

Current Experiments


Future Experiments

GeV-PeV Neutrinos

Big Bang Neutrinos

ν Applications


Conclusions




Latest results from NOνA

Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions


$\nu_\mu \rightarrow \nu_\mu$




$\bar{\nu}_\mu \rightarrow \bar{\nu}_\mu$



$\nu_\mu \rightarrow \nu_e$



$\bar{\nu}_\mu \rightarrow \bar{\nu}_e$





Off-axis high intensity ν_μ beams: T2K

Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions


First proposed for BNL E-889 (1995): A narrow beam of ν can be achieved by going off-axis to the π beam. Better S:B at oscillation max.


Signal at $\sin^2 2\theta_{13} = 0.1$:





SuperKamiokande





INGRID ND

T2K first results announced in March 2011

T2K beam ν_e Candidate Event 2010

Neutrinos II

Mary Bishai Brookhaven National Laboratory

MeV Neutrinos from Reactors
Exercises
Daya Bay


GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments

GeV-PeV Neutrinos

Big Bang Neutrinos

ν Applications

Conclusions



Item	Event	T2K cut
Date (JST)	2010 May 12th 21:3:22	
Ring, PID	1-Ring electron-like	OK
Momentum	378 MeV	>100
N_{dcy}	0	0
$\cos(\theta_{\nu e})$	0.55 (57 degree)	N/A
$Mass$	0.13 MeV	<105
E_{rec}	496 MeV	<1250

T2K: First Observation of $\nu_\mu \rightarrow \nu_e$ APPEARANCE

Neutrinos II

Mary Bishai Brookhaven National Laboratory

MeV Neutrinos from Reactors
Exercises
Daya Bay


GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments

GeV-PeV Neutrinos

Big Bang Neutrinos

ν Applications

Conclusions



In 2014 T2K observes conversion of ν_μ to ν_e (atmospheric oscillation scale) with an amplitude of $\sin^2 2\theta_{13} = 0.140^{+0.038}_{-0.032}$.

2016 Breakthrough Prize in Fundamental Physics

Neutrinos II

Mary Bishai Brookhaven National Laboratory

MeV Neutrinos from Reactors
Exercises
Daya Bay

GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments

GeV-PeV Neutrinos

Big Bang Neutrinos

ν Applications

Conclusions



The 2016 Breakthrough Prize in Fundamental Physics awarded to 7 leaders and 1370 members of 5 experiments investigating neutrino oscillation: Daya Bay (China); KamLAND (Japan); K2K / T2K (Japan); Sudbury Neutrino Observatory (Canada); and Super-Kamiokande (Japan)

The Deep Underground Neutrino Experiment

Neutrinos II

Mary Bishai Brookhaven National Laboratory

MeV Neutrinos from Reactors

Exercises

Daya Bay

GeV Neutrinos from Accelerators

Exercises

Current Experiments


Future Experiments

GeV-PeV Neutrinos

Big Bang Neutrinos

ν Applications

Conclusions



- **A very long baseline experiment: 1300km from Fermilab in Batavia, IL to the Sanford Underground Research Facility (former Homestake Gold Mine) in Lead, SD.**
- **A highly capable near detector at Fermilab.**
- **A very deep (1.5 km underground) far detector: massive 40-kton Liquid Argon Time-Projection-Chamber with state-of-the-art instrumentation.**
- **High intensity *tunable* wide-band neutrino beam from LBNF produced from the 1-2MW upgraded Main Injecctor accelerator at Fermilab.**

The DUNE Scientific Collaboration


Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions

As of Jan 2018:

60 % non-US

1061 collaborators from 175 institutions in 31 nations

Armenia, Brazil, Bulgaria, Canada, CERN, Chile, China, Colombia, Czech Republic, Finland, France, Greece, India, Iran, Italy, Japan, Madagascar, Mexico, Netherlands, Paraguay, Peru, Poland, Romania, Russia, South Korea, Spain, Sweden, Switzerland, Turkey, UK, Ukraine, USA





Scientific Objectives of DUNE

Neutrinos II

Mary Bishai Brookhaven National Laboratory

MeV Neutrinos from Reactors

Exercises

Daya Bay

GeV Neutrinos from Accelerators

Exercises

Current Experiments


Future Experiments

GeV-PeV Neutrinos

Big Bang Neutrinos

ν Applications

Conclusions



1. **precision measurements of the parameters that govern $\nu_\mu \rightarrow \nu_e$ oscillations; this includes precision measurement of the third mixing angle θ_{13}, measurement of the charge-parity (CP) violating phase δ_{CP}, and determination of the neutrino mass ordering (the sign of $\Delta m^2_{31} = m^2_3 - m^2_1$), the so-called mass hierarchy**
2. **precision measurements of the mixing angle θ_{23}, including the determination of the octant in which this angle lies, and the value of the mass difference, $|\Delta m^2_{32}|$, in $\nu_\mu \rightarrow \nu_{e,\mu}$ oscillations**

Scientific Objectives of DUNE

Neutrinos II

Mary Bishai Brookhaven National Laboratory

MeV Neutrinos from Reactors

Exercises

Daya Bay

GeV Neutrinos from Accelerators

Exercises

Current Experiments


Future Experiments


GeV-PeV Neutrinos

Big Bang Neutrinos

ν Applications

Conclusions






3 **search for proton decay, yielding significant improvement in the current limits on the partial lifetime of the proton (τ/BR) in one or more important candidate decay modes, e.g., $p \rightarrow K^+\overline{\nu}$**

4 **detection and measurement of the neutrino flux from a core-collapse supernova within our galaxy, should one occur during the lifetime of DUNE**

The Sanford Underground Research Facility

Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions



Experimental facility operated by the state of South Dakota. LUX (dark matter) and Majorana ($0\nu - 2\beta$) demonstrator operational expts at 4850-ft level. Chosen as site of G2 dark matter experiment

The DUNE Far Detector

Neutrinos II

Mary Bishai Brookhaven National Laboratory

MeV Neutrinos from Reactors
Exercises
Daya Bay

GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments

GeV-PeV Neutrinos


Big Bang Neutrinos

ν Applications

Conclusions

A large cryogenic liquid Argon detector located a mile underground in the former Homestake Mine with a mass of at least 40 kilo-tons is used to image neutrino interactions with unprecedented precision:

Single Phase LArTPC





The DUNE prototype wireplane

The DUNE Far Detector

Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions

Dual Phase LArTPC


4.) Charge **collection** on a **2D anode readout** (symmetric unipolar signals with two orthogonal views)


3.) Charge multiplication in the holes of the **Large Electron Multiplier (LEM)**



2.) Drift electrons are efficiently emitted into the gas phase

1.) Ionization electrons drift towards the liquid argon surface





The DUNE Far Detector

Neutrinos II

Mary Bishai Brookhaven National Laboratory

MeV Neutrinos from Reactors
Exercises
Daya Bay

GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments


GeV-PeV Neutrinos

Big Bang Neutrinos

ν Applications

Conclusions

The 40-kton (fiducial) detector is constructed of four modules with a total mass of 17.4 kton each.



External (Internal) Dimensions
19.1m (16.9m) W x 18.0m (15.8m) H x 66.0m (63.8m) L

Reconstructed Neutrino Interactions in a LArTPC

Neutrinos II

Mary Bishai Brookhaven National Laboratory

MeV Neutrinos from Reactors
Exercises
Daya Bay


GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments


GeV-PeV Neutrinos

Big Bang Neutrinos

ν Applications

Conclusions







DUNE Event Spectra

Exposure: 150 kT.MW.yr (equal $\nu/\bar{\nu}$) 1MW.yr = 1×10^{21} p.o.t at 120 GeV. ($\sin^2 2\theta_{13} = 0.085$, $\sin^2 \theta_{23} = 0.45$, $\delta m^2_{31} = 2.46 \times 10^{-3}$ eV^2)


Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions




8329 ν_μ, 192 $\bar{\nu}_\mu$, 72 NC, 29 ν_τ



2420 $\bar{\nu}_\mu$, 791 ν_μ, 33 NC, 13 ν_τ



930 ν_e, 5 $\bar{\nu}_e$, 204 ν_e^{beam}, 17 NC, 19 ν_τ, 3 ν_μ




154 $\bar{\nu}_e$,32 ν_e, 98 ν_e^{beam}, 7 NC, 8 ν_τ, 1 ν_μ

Simultaneous fit to all four samples. Richness of spectral information in both ν_μ and $\bar{\nu}_\mu$ $\Rightarrow$ explicit demonstration of CPV


DUNE Event Spectra

Exposure: 150 kT.MW.yr (equal $\nu/\bar{\nu}$) 1MW.yr = 1×10^{21} p.o.t at 120 GeV. ($\sin^2 2\theta_{13} = 0.085$, $\sin^2 \theta_{23} = 0.45$, $\delta m^2_{31} = 2.46 \times 10^{-3}$ eV^2)


Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions




8329 ν_μ, 192 $\bar{\nu}_\mu$, 72 NC, 29 ν_τ



2420 $\bar{\nu}_\mu$, 791 ν_μ, 33 NC, 13 ν_τ



1171 ν_e,3 $\bar{\nu}_e$, 204 ν_e^{beam}, 17 NC, 19 ν_τ, 3 ν_μ



94 $\bar{\nu}_e$,39 ν_e, 98 ν_e^{Beam}, 7 NC, 8 ν_τ, 1 ν_μ

Simultaneous fit to all four samples. Richness of spectral information in both ν_μ and $\bar{\nu}_\mu$ $\Rightarrow$ explicit demonstration of CPV


Possible Supernova Signature in DUNE

Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions


Liquid argon is particularly sensitive to the ν_e component of a supernova neutrino burst:

$$\nu_e + {}^{40}\mathrm{Ar} \rightarrow e^- + {}^{40}\mathrm{K}^*, \tag{1}$$

Expected time-dependent signal in 40 kton of liquid argon for a Supernova at 10 kpc:



Time distribution



Energy spectrum (time integrated)

LBNF/DUNE Schedule

Neutrinos II

Mary Bishai Brookhaven National Laboratory

MeV Neutrinos from Reactors

Exercises

Daya Bay

GeV Neutrinos from Accelerators

Exercises

Current Experiments

Future Experiments

GeV-PeV Neutrinos

Big Bang Neutrinos

ν Applications

Conclusions

- **2017: Cavern excavation begins**
- **2018: DUNE prototypes (single & dual phase) operational in test beam at CERN**
- **2019: Technical design review (beam and far detectors) by US-DOE and international funding energies. Conceptual design for near detector ready.**
- **2021: First 10kton FD module (single phase) installation**
- **2023: Second FD module (single or dual phase) installation**
- **2024: Data taking (non-beam) starts with 20 kton operational**
- **2026: First beam operations at 1.2 MW**

The IceCUBE Experiment

Neutrinos II

Mary Bishai Brookhaven National Laboratory

MeV Neutrinos from Reactors

Exercises

Daya Bay

GeV Neutrinos from Accelerators

Exercises

Current Experiments


Future Experiments

GeV-PeV Neutrinos

Big Bang Neutrinos

ν Applications

Conclusions



The IceCUBE Experiment

Neutrinos II

Mary Bishai Brookhaven National Laboratory

MeV Neutrinos from Reactors
Exercises
Daya Bay


GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments

GeV-PeV Neutrinos

Big Bang Neutrinos

ν Applications

Conclusions



The Highest Energy Neutrinos (Gamma Ray Bursts)

Neutrinos II

Mary Bishai Brookhaven National Laboratory

MeV Neutrinos from Reactors

Exercises Daya Bay

GeV Neutrinos from Accelerators


Exercises Current Experiments Future Experiments

GeV-PeV Neutrinos

Big Bang Neutrinos

ν Applications

Conclusions



The Highest Energy Neutrinos (Gamma Ray Bursts)

Neutrinos II

Mary Bishai Brookhaven National Laboratory

MeV Neutrinos from Reactors
Exercises
Daya Bay

GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments


GeV-PeV Neutrinos

Big Bang Neutrinos

ν Applications

Conclusions

Neutrino events with energies $>$ PeV ($10^{15} eV$)



The Highest Energy Neutrinos (Gamma Ray Bursts)

Neutrinos II

Mary Bishai Brookhaven National Laboratory

MeV Neutrinos from Reactors

Exercises

Daya Bay

GeV Neutrinos from Accelerators

Exercises

Current Experiments


Future Experiments

GeV-PeV Neutrinos

Big Bang Neutrinos

ν Applications

Conclusions



NEW RESULT FROM ICECUBE: July 12,2018

Neutrinos II

Mary Bishai Brookhaven National Laboratory

MeV Neutrinos from Reactors
Exercises
Daya Bay

GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments


GeV-PeV Neutrinos

Big Bang Neutrinos

ν Applications

Conclusions

Sep 22, 2017 a very high energy muon neutrino is recorded in IceCUBE:



NEW RESULT FROM ICECUBE: July 12,2018

Neutrinos II

Mary Bishai Brookhaven National Laboratory

MeV Neutrinos from Reactors
Exercises
Daya Bay

GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments


GeV-PeV Neutrinos

Big Bang Neutrinos

ν Applications

Conclusions

Sep 22, 2017 a very high energy neutrino is recorded in IceCUBE:



NEW RESULT FROM ICECUBE: July 12,2018

Neutrinos II

Mary Bishai Brookhaven National Laboratory

MeV Neutrinos from Reactors
Exercises
Daya Bay

GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments


GeV-PeV Neutrinos

Big Bang Neutrinos

ν Applications

Conclusions


Fermi-LAT (space telescope) also sees a flare up of very high energy gamma rays from a source near Orion



NEW RESULT FROM ICECUBE: July 12,2018

Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions

Eventually many observatories see the burst in optical, γs, X-rays and radio:



A new dawn for multi-messengar astronomy

PTOLEMY: Detecting Big Bang Neutrinos

Neutrinos II

Mary Bishai Brookhaven National Laboratory

MeV Neutrinos from Reactors

Exercises

Daya Bay

GeV Neutrinos from Accelerators

Exercises

Current Experiments

Future Experiments


GeV-PeV Neutrinos

Big Bang Neutrinos

ν Applications

Conclusions

Princeton Tritium Observatory for Light, Early-Universe, Massive-Neutrino Yield



How to detect Big Bang Neutrinos

Neutrinos II

Mary Bishai Brookhaven National Laboratory

MeV Neutrinos from Reactors
Exercises
Daya Bay

GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments

GeV-PeV Neutrinos

Big Bang Neutrinos

ν Applications

Conclusions

From paper by Steven Weinberg in 1962 (Phys. Rev. 128:3 1457]. Detect capture of BB neutrinos on a beta decaying nucleus:





Figure 1: Emitted electron density of states vs kinetic energy for neutrino capture on beta decaying nuclei. The spike at $Q + 2m$ is the CNB signal

Experimental Concept


Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions




Many techincal challenges!!!

Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions

The biggest nearly insurmountable problem for relic neutrino detection using capture on tritium is to provide a large enough surface area to hold at least 100 grams of weakly bound atomic tritium!



Ultra-modern materials science needed: Use tritium trapped in very thin layers of graphene:




Practical Applications of Technologies for ν Experiments


Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions

Synergies and Applications - Examples


Cyclotrons for neutrino physics (and industrial applications)



KEN K2600 SUPERCONDUCTING RING CYCLOTRON



Neutrino detectors for reactor monitoring and non-proliferation





remote discovery of undeclared nuclear reactors with large detectors at km scale




US Short-Baseline Experiment

reactor antineutrino studies at short baselines

Multi-MW Accelerators Driving Thorium Reactors

Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions

First proposed by Carlo Rubbia in 1995 (1984 Nobel Prize winner)



Global energy resources in ZetaJoules

Resource	*Type*	*Yearly consumption (1999) ZJ*	*Resources ZJ*	*Consumed until 1999 (ZJ)*
Oil	Conventional	0.13	12.08	4.85
	Unconventional	0.01	20.35	0.29
	Total oil	0.14	32.42	5.14
Natural gas	Conventional	0.08	16.56	2.35
	Unconventional	0.00	33.23	0.03
	Total gas	0.08	49.79	2.38
Coal	Total coal	0.09	199.67	5.99
Total Fossils		0.31	281.88	13.51
Uranium	Thermal reactors	0.04	5.41 (2'000, sw)	
	Breeder	0	324 (120'000, sw)	
Thorium			1'300'000	

sw: including sea water
1 ZJ (ZetaJoule)= 10^3 EJ(ExaJoule)= 10^{21} J(Joule)

Requires proton accelerators with powers of 10 MW. Currently neutrino and neutron experiments are driving the technology of high power MW class proton beams.






Figure 1. Schematic representation of Energy Amplifier proposed by Rubbia [4]


Neutrinos and Earth's Geology

Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises
Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions

Plate Tectonics,
Convection,
Geodynamo








Does heat from radioactive decay drive the Earth's engine?

Neutrinos and Earth's Geology

Neutrinos II
Mary Bishai Brookhaven National Laboratory
MeV Neutrinos from Reactors
Exercises Daya Bay
GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments
GeV-PeV Neutrinos
Big Bang Neutrinos
ν Applications
Conclusions

Signal of $\bar{\nu}_e$ from radioactive decays of U/TH in the earth observed in the BOREXINO solar neutrino experiment:



Summary

Neutrinos II

Mary Bishai Brookhaven National Laboratory

MeV Neutrinos from Reactors
Exercises
Daya Bay

GeV Neutrinos from Accelerators
Exercises
Current Experiments
Future Experiments

GeV-PeV Neutrinos

Big Bang Neutrinos

ν Applications

Conclusions

- **Neutrinos have been at the forefront of fundamental discoveries in particle physics for decades.**
- **Discoveries of neutrino properties like the very small mass, large almost maximal mixing, are the *ONLY direct evidence for physics beyond the Standard Model of particle physics, and new hidden symmetries.***
- **Neutrinos straddle the fields of nuclear physics, particle astrophysics, cosmology and high energy particle physics. Thus, they provide a unique probe to test for consistency in our picture of the Universe from the development of the Big Bang, the mechanics of Supernova explosions, the chemistry of stars, the geology of the earth, and the nuclear physics of reactors.**
- **Studying the properties of neutrinos with energies varying from the very cold (Big Bang ν) to the PeV scale requires a huge diversity of experiments, each with its own unique technical challenges.**



THANK YOU

Click for Neutrino rap!!