Entanglement at collider energies

Stefan Floerchinger (Heidelberg U.)

Initial Stages 2019 conference, Columbia University, New York City 27 June 2019







What is entanglement and why is it interesting?

• Can quantum-mechanical description of physical reality be considered complete? [Einstein, Podolsky, Rosen (1935), Bohm (1951)]

$$\psi = \frac{1}{\sqrt{2}} \left(|\uparrow\rangle_A |\downarrow\rangle_B - |\downarrow\rangle_A |\uparrow\rangle_B \right)$$
$$= \frac{1}{\sqrt{2}} \left(|\rightarrow\rangle_A |\leftrightarrow\rangle_B - |\leftrightarrow\rangle_A |\rightarrow\rangle_B \right)$$

• Bertlemann's socks and the nature of reality [Bell (1980)]



Bell's inequalities and Bell tests

[John Stewart Bell (1966)]

• most popular version [Clauser, Horne, Shimony, Holt (1969)]

 $S = |E(a, b) - E(a, b') + E(a', b) + E(a', b')| \le 2$

holds for local hidden variable theories

expectation value of product of two observables

 $E(a,b) = \langle A(a)B(b)\rangle$

with possible values $A = \pm 1$, $B = \pm 1$.

- depending on measurement settings a, a' and b, b' respectively
- quantum mechanical bound is $S \le 2\sqrt{2}$
- experimental values $2 < S \le 2\sqrt{2}$ rule out local hidden variables
- one measurement setting but at different times [Leggett, Garg (1985)]

Entanglement at collider energies

[..., Elze (1996), Kovner, Lublinsky (2015), Kharzeev & Levin (2017), Berges, Floerchinger & Venugopalan (2017), Shuryak & Zahed (2017), Kovner, Lublinsky, Serino (2018), Baker & Kharzeev (2018), Tu, Kharzeev & Ullrich (2019), Armesto, Dominguez, Kovner, Lublinsky, Skokov (2019), ...]

- entanglement of quantum fields instead of particles
- entanglement on sub-nucleonic scales
- $\bullet\,$ entanglement in non-Abelian gauge theory / color / confinement
- discussions in mathematical physics [e. g. Witten (2018)]
- connections to black holes and holography [Ryu & Takayanagi (2006)]
- thermalization in closed quantum systems

Entropy in quantum theory

[John von Neumann (1932)]

 $S = -\mathsf{Tr}\{\rho \ln \rho\}$

- \bullet based on the quantum density operator ρ
- \bullet for pure states $\rho = |\psi\rangle \langle \psi|$ one has S=0
- for mixed states $ho = \sum_j p_j |j\rangle \langle j|$ one has $S = -\sum_j p_j \ln p_j > 0$
- unitary time evolution conserves entropy

 $-\mathrm{Tr}\{(U\rho U^{\dagger})\ln(U\rho U^{\dagger})\} = -\mathrm{Tr}\{\rho\ln\rho\} \qquad \rightarrow \qquad S = \mathrm{const.}$

• global characterization of quantum state

Local dissipation, entropy and entanglement

• local dissipation = local entropy production

 $-\nabla_{\mu}s^{\mu}(x) \ge 0$

• relativistic fluid dynamics in Navier-Stokes approximation

$$-\nabla_{\mu}s^{\mu} = \frac{1}{T} \left[2\eta\sigma_{\mu\nu}\sigma^{\mu\nu} + \zeta(\nabla_{\rho}u^{\rho})^2 \right]$$

- can not be density of global von-Neumann entropy for closed system
- kinetic theory for weakly coupled (quasi-) particles [Boltzmann (1890)]

$$s^{\mu}(x) = -\int \frac{d^3p}{p^0} \left\{ p^{\mu} f(x,p) \ln f(x,p) \right\}$$

- how to go beyond weak coupling / quasiparticles?
- local dissipation = *entanglement generation*
- $s^{\mu}(x)$ must (most likely) be seen as entanglement entropy current

The thermal model puzzle

- elementary particle collision experiments such as $e^+ e^-$ collisions show some thermal-like features
- particle multiplicities well described by thermal model



[Becattini, Casterina, Milov & Satz, EPJC 66, 377 (2010)]

- conventional thermalization by collisions unlikely
- more thermal-like features difficult to understand in PyTHIA [Fischer, Sjöstrand (2017)]
- alternative explanations needed

$QCD \ strings$



- particle production from QCD strings
- Lund string model (e. g. PYTHIA)
- different regions in a string are entangled
- \bullet subinterval A is described by reduced density matrix

 $\rho_A = \mathsf{Tr}_B\{\rho\}$

- reduced density matrix is of mixed state form
- could this lead to thermal-like effects?

Entropy and entanglement

• consider a split of a quantum system into two A + B



 $\bullet\,$ reduced density operator for system A

 $\rho_A = \mathsf{Tr}_B\{\rho\}$

• entropy associated with subsystem A: entanglement entropy

$$S_A = -\mathsf{Tr}_A\{\rho_A \ln \rho_A\}$$

- globally pure state S = 0 can be locally mixed $S_A > 0$
- coherent information $I_{B \mid A} = S_A S$ can be positive

$Microscopic \ model$

 \bullet QCD in 1+1 dimensions described by 't Hooft model

$$\mathscr{L} = -ar{\psi}_i \gamma^\mu (\partial_\mu - ig \mathbf{A}_\mu) \psi_i - m_i ar{\psi}_i \psi_i - rac{1}{2} \mathrm{tr} \, \mathbf{F}_{\mu
u} \mathbf{F}^{\mu
u}$$

- fermionic fields ψ_i with sums over flavor species $i=1,\ldots,N_f$
- SU (N_c) gauge fields ${f A}_\mu$ with field strength tensor ${f F}_{\mu
 u}$
- gluons are not dynamical in two dimensions
- \bullet gauge coupling g has dimension of mass
- non-trivial, interacting theory, cannot be solved exactly
- spectrum of excitations known for $N_c \rightarrow \infty$ with $g^2 N_c$ fixed ['t Hooft (1974)]

Schwinger model

• QED in 1+1 dimension

$$\mathscr{L} = -\bar{\psi}_i \gamma^\mu (\partial_\mu - iqA_\mu)\psi_i - m_i \bar{\psi}_i \psi_i - \frac{1}{4} F_{\mu\nu} F^{\mu\nu}$$

- geometric confinement
- U(1) charge related to string tension $q = \sqrt{2\sigma}$
- for single fermion one can bosonize theory exactly [Coleman, Jackiw, Susskind (1975)]

$$S = \int d^2x \sqrt{g} \left\{ -\frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - \frac{1}{2} M^2 \phi^2 - \frac{m \, q \, e^\gamma}{2\pi^{3/2}} \cos\left(2\sqrt{\pi}\phi + \theta\right) \right\}$$

- Schwinger bosons are dipoles $\phi \sim \bar{\psi} \psi$
- scalar mass related to U(1) charge by $M=q/\sqrt{\pi}=\sqrt{2\sigma/\pi}$
- massless Schwinger model m = 0 leads to free bosonic theory

Expanding string solution 1



- external quark-anti-quark pair on trajectories $z = \pm t$
- coordinates: Bjorken time $\tau = \sqrt{t^2 z^2}$, rapidity $\eta = \operatorname{arctanh}(z/t)$
- metric $ds^2 = -d\tau^2 + \tau^2 d\eta^2$
- $\bullet\,$ symmetry with respect to longitudinal boosts $\eta \to \eta + \Delta \eta$

Expanding string solution 2

 $\bullet\,$ Schwinger boson field depends only on τ

$$\bar{\phi} = \bar{\phi}(\tau)$$

• equation of motion

$$\partial_{\tau}^2 \bar{\phi} + \frac{1}{\tau} \partial_{\tau} \bar{\phi} + M^2 \bar{\phi} = 0.$$

• Gauss law: electric field $E = q\phi/\sqrt{\pi}$ must approach the U(1) charge of the external quarks $E \to q_e$ for $\tau \to 0_+$

$$\bar{\phi}(\tau) \to \frac{\sqrt{\pi}q_{\mathsf{e}}}{q} \qquad (\tau \to 0_+)$$

• solution of equation of motion [Loshaj, Kharzeev (2011)]

$$\bar{\phi}(\tau) = \frac{\sqrt{\pi}q_{\mathsf{e}}}{q} J_0(M\tau)$$

Gaussian states

- theories with quadratic action often have Gaussian density matrix
- fully characterized by field expectation values

 $\bar{\phi}(x) = \langle \phi(x) \rangle, \qquad \bar{\pi}(x) = \langle \pi(x) \rangle$

and connected two-point correlation functions, e. g.

$$\langle \phi(x)\phi(y)\rangle_c = \langle \phi(x)\phi(y)\rangle - \bar{\phi}(x)\bar{\phi}(y)$$

• if ρ is Gaussian, also reduced density matrix ρ_A is Gaussian

Entanglement entropy for Gaussian state

• entanglement entropy of Gaussian state in region A [Berges, Floerchinger, Venugopalan, JHEP 1804 (2018) 145]

$$S_A = \frac{1}{2} \operatorname{Tr}_A \left\{ D \ln(D^2) \right\}$$

- operator trace over region A only
- matrix of correlation functions

$$D(x,y) = \begin{pmatrix} -i\langle\phi(x)\pi(y)\rangle_c & i\langle\phi(x)\phi(y)\rangle_c \\ -i\langle\pi(x)\pi(y)\rangle_c & i\langle\pi(x)\phi(y)\rangle_c \end{pmatrix}$$

- \bullet involves connected correlation functions of field $\phi(x)$ and canonically conjugate momentum field $\pi(x)$
- expectation value $\bar{\phi}$ does not appear explicitly
- coherent states and vacuum have equal entanglement entropy S_A

Rapidity interval



- $\bullet\,$ consider rapidity interval $(-\Delta\eta/2,\Delta\eta/2)$ at fixed Bjorken time τ
- entanglement entropy does not change by unitary time evolution with endpoints kept fixed
- can be evaluated equivalently in interval $\Delta z = 2\tau \sinh(\Delta \eta/2)$ at fixed time $t = \tau \cosh(\Delta \eta/2)$
- need to solve eigenvalue problem with correct boundary conditions

Bosonized massless Schwinger model

- entanglement entropy understood numerically for free massive scalars [Casini, Huerta (2009)]
- entanglement entropy density $dS/d\Delta\eta$ for bosonized massless Schwinger model ($M=\frac{q}{\sqrt{\pi}})$



Conformal limit

• For M au o 0 one has conformal field theory limit [Holzhey, Larsen, Wilczek (1994)]

$$S(\Delta z) = rac{c}{3} \ln{(\Delta z/\epsilon)} + {
m constant}$$

with small length ϵ acting as UV cutoff.

Here this implies

$$S(\tau, \Delta \eta) = \frac{c}{3} \ln \left(2\tau \sinh(\Delta \eta/2)/\epsilon \right) + \text{constant}$$

- Conformal charge c = 1 for free massless scalars or Dirac fermions.
- Additive constant not universal but entropy density is

$$\begin{split} \frac{\partial}{\partial \Delta \eta} S(\tau, \Delta \eta) = & \frac{c}{6} \mathsf{coth}(\Delta \eta / 2) \\ \to & \frac{c}{6} \qquad (\Delta \eta \gg 1) \end{split}$$

• Entropy becomes extensive in $\Delta \eta$!

Universal entanglement entropy density

• for very early times "Hubble" expansion rate dominates over masses and interactions

$$H = \frac{1}{\tau} \gg M = \frac{q}{\sqrt{\pi}}, m$$

- theory dominated by free, massless fermions
- universal entanglement entropy density

$$\frac{dS}{d\Delta\eta} = \frac{c}{6}$$

with conformal charge c

• for QCD in 1+1 D (gluons not dynamical, no transverse excitations)

 $c = N_c \times N_f$

• from fluctuating transverse coordinates (Nambu-Goto action)

$$c = N_c \times N_f + 2 \approx 9 + 2 = 11$$

Modular or entanglement Hamiltonian 1



- conformal field theory
- $\bullet\,$ hypersurface Σ with boundary on the intersection of two light cones
- reduced density matrix [Casini, Huerta, Myers (2011), Arias, Blanco, Casini, Huerta (2017), see also Candelas, Dowker (1979)]

$$\rho_A = \frac{1}{Z_A} e^{-K}, \qquad Z_A = \operatorname{Tr} e^{-K}$$

• modular or entanglement Hamiltonian K

Modular or entanglement Hamiltonian 2

• modular or entanglement Hamiltonian is local expression

$$K = \int_{\Sigma} d\Sigma_{\mu} \, \xi_{\nu}(x) \, T^{\mu\nu}(x).$$

- energy-momentum tensor $T^{\mu\nu}(x)$ of excitations
- vector field

$$\begin{split} \xi^{\mu}(x) &= \frac{2\pi}{(q-p)^2} [(q-x)^{\mu}(x-p)(q-p) \\ &\quad + (x-p)^{\mu}(q-x)(q-p) - (q-p)^{\mu}(x-p)(q-x)] \end{split}$$

end point of future light cone q, starting point of past light cone p• inverse temperature and fluid velocity

$$\xi^{\mu}(x) = \beta^{\mu}(x) = \frac{u^{\mu}(x)}{T(x)}$$

Modular or entanglement Hamiltonian 3



• for $\Delta\eta \rightarrow \infty$: fluid velocity in τ -direction, τ -dependent temperature

$$T(\tau) = \frac{\hbar}{2\pi\tau}$$

- Entanglement between different rapidity intervals alone leads to local thermal density matrix at very early times !
- Hawking-Unruh temperature in Rindler wedge $T(x) = \hbar c/(2\pi x)$

Physics picture

- coherent state vacuum at early time contains entangled pairs of quasi-particles with opposite wave numbers
- on finite rapidity interval $(-\Delta\eta/2,\Delta\eta/2)$ in- and out-flux of quasi-particles with thermal distribution via boundaries
- \bullet technically limits $\Delta\eta \to \infty$ and $M\tau \to 0$ do not commute
 - $\Delta\eta \to \infty$ for any finite $M\tau$ gives pure state
 - $M\tau \to 0$ for any finite $\Delta \eta$ gives thermal state with $T=1/(2\pi\tau)$

Particle production in massive Schwinger model

[ongoing work with Lara Kuhn, Jürgen Berges]



- for expanding strings
- $\bullet\,$ asymptotic particle number depends on $g\sim m/q$
- \bullet exponential suppression for large fermion mass $g\gg 1$

$$\frac{N}{\Delta \eta} \sim e^{-0.55\frac{m}{q} + 7.48\frac{q}{m} + \dots} = e^{-0.55\frac{m}{\sqrt{2\sigma}} + 7.48\frac{\sqrt{2\sigma}}{m} + \dots}$$

Conclusions

- entanglement at colliders is fascinating emerging topic of research
- experimental proof for entanglement needs Bell test
- entanglement entropy useful to describe local thermalization
- rapidity intervals in an expanding string are entangled
- at very early times theory effectively conformal

$$\frac{1}{\tau} \gg m, q$$

reduced density matrix is of locally thermal form with temperature

$$T = \frac{\hbar}{2\pi\tau}$$

• entanglement important to understand early thermalization

Backup

Wigner distribution and entanglement

- Classical field approximation usually based on non-negative Wigner representation of density matrix
- leads for many observables to classical statistical description
- can nevertheless show entanglement and pass Bell test for "improper" variables where Weyl transform of operator has values outside of its spectrum [Revzen, Mello, Mann, Johansen (2005)]
- Bell test violation also possible for negative Wigner distribution [Bell (1986)]

Transverse coordinates

- so far dynamics strictly confined to 1+1 dimensions
- transverse coordinates may fluctuate, can be described by Nambu-Goto action $(h_{\mu\nu} = \partial_{\mu} X^m \partial_{\nu} X_m)$

$$S_{\rm NG} = \int d^2x \sqrt{-\det h_{\mu\nu}} \{-\sigma + \ldots\}$$
$$\approx \int d^2x \sqrt{g} \left\{ -\sigma - \frac{\sigma}{2} g^{\mu\nu} \partial_{\mu} X^i \partial_{\nu} X^i + \ldots \right\}$$

• two additional, massless, bosonic degrees of freedom corresponding to transverse coordinates X^i with i = 1, 2

Temperature and entanglement entropy

- for conformal fields, entanglement entropy has also been calculated at non-zero temperature.
- for static interval of length L [Korepin (2004); Calabrese, Cardy (2004)]

$$S(T, l) = \frac{c}{3} \ln \left(\frac{1}{\pi T \epsilon} \sinh(\pi L T) \right) + \text{const}$$

• compare this to our result in expanding geometry

$$S(\tau,\Delta\eta) = \frac{c}{3}\ln\left(\frac{2\tau}{\epsilon}\sinh(\Delta\eta/2)\right) + {\rm const}$$

• expressions agree for $L = \tau \Delta \eta$ (with metric $ds^2 = -d\tau^2 + \tau^2 d\eta^2$) and time-dependent temperature

$$T = \frac{1}{2\pi\tau}$$

Rapidity distribution



[open (filled) symbols: e⁺e⁻ (pp), Grosse-Oetringhaus & Reygers (2010)]

- rapidity distribution $dN/d\eta$ has plateau around midrapidity
- only logarithmic dependence on collision energy

Experimental access to entanglement?

- could longitudinal entanglement be tested experimentally?
- \bullet unfortunately entropy density $dS/d\eta$ not straight-forward to access
- measured in e^+e^- is the number of charged particles per unit rapidity $dN_{\rm ch}/d\eta$ (rapidity defined with respect to the thrust axis)
- $\bullet\,$ typical values for collision energies $\sqrt{s}=14-206~{\rm GeV}$ in the range

 $dN_{\rm ch}/d\eta \approx 2-4$

• entropy per particle S/N can be estimated for a hadron resonance gas in thermal equilibrium $S/N_{\rm ch}=7.2$ would give

 $dS/d\eta \approx 14 - 28$

• this is an upper bound: correlations beyond one-particle functions would lead to reduced entropy

Entanglement and QCD physics

- how strongly entangled is the nuclear wave function?
- what is the entropy of quasi-free partons and can it be understood as a result of entanglement? [Kharzeev, Levin (2017)]
- does saturation at small Bjorken-x have an entropic meaning?
- entanglement entropy and entropy production in the color glass condensate [Kovner, Lublinsky (2015); Kovner, Lublinsky, Serino (2018)]
- could entanglement entropy help for a non-perturbative extension of the parton model?
- entropy of perturbative and non-perturbative Pomeron descriptions [Shuryak, Zahed (2017)]