

# Initial Stages 2019



## Report of Contributions

Contribution ID: 5

Type: **Oral**

## Hydrodynamic simulations of relativistic nuclear collisions with nucleon substructure: combined analysis of p+Pb and Pb+Pb collision systems at 5.02 TeV

*Wednesday, 26 June 2019 17:30 (20 minutes)*

Simulations of relativistic heavy-ion collisions based on viscous hydrodynamics provide an accurate description of the bulk observables measured at RHIC and LHC beam energies, including identified particle yields, mean  $p_T$  and multiparticle correlations. The success of the hydrodynamic framework, however, is naturally expected to break down in the dilute limit where discrete particle degrees of freedom dominate.

It was thus surprising when the multiparticle correlations measured in high-multiplicity proton-lead collisions were found to be similar in magnitude to those observed in lead-lead collisions. The observation suggests that hydrodynamic behavior could be manifest in small droplets of quark-gluon plasma (QGP), and that flow might develop at length scales smaller than a proton.

In this work, we assume the existence of hydrodynamic flow in small collision systems and evaluate the likelihood of our assertion using Bayesian inference. Specifically, we model the dynamics of proton-lead and lead-lead collisions at 5.02 TeV using QGP initial conditions with parametric nucleon substructure, a pre-equilibrium free-streaming stage, event-by-event viscous hydrodynamics with shear and bulk coupling, and a microscopic hadronic afterburner to simulate the dynamics of the collision below the QGP transition temperature.

The model is evaluated on a scaffolding of parameter points, and emulators are trained to interpolate the model predictions at intermediate regions of parameter space. Markov chain Monte Carlo importance sampling is then used to explore the Bayesian posterior probability distribution as a function of the model input parameters.

We use the resulting posterior distribution to sample preferred regions of parameter space and evaluate the performance of the model with optimally chosen parameter values. This semi-exhaustive model validation enables us to comment on the implied viability of hydrodynamics in small collision systems subject to the approximations of the chosen framework. We also present marginalized posterior distributions for each model input parameter, e.g. nucleon substructure degrees of freedom, which demonstrate the constraining power of global statistical analysis and reveal new insight into nuclear matter at extreme temperatures and densities.

**Primary author:** BASS, Steffen (Duke)

**Co-authors:** MORELAND, Scott (Duke University); Dr BERNHARD, Jonah (Duke University)

**Presenter:** BASS, Steffen (Duke)

**Session Classification:** Parallel: Collectivity in small systems 2

**Track Classification:** Collectivity in small systems

Contribution ID: 9

Type: **Oral**

## New heavy flavor program for the future Electron Ion Collider

*Tuesday, 25 June 2019 16:21 (1 minute)*

The proposed high-luminosity high-energy Electron Ion Collider (EIC) will provide one of the cleanest environments to precisely determine the nuclear parton distribution functions (nPDFs) in a wide  $x$ - $Q^2$  phase space. Heavy flavor production at the EIC can access up to the confinement boundary, which allows us to directly study nPDFs, quark/gluon fragmentation processes, and energy loss within the poorly constrained high Bjorken- $x$  region. The group at Los Alamos National Laboratory propose to develop a new experimental and theoretical physics program to study the heavy flavor products, flavor tagged jets and heavy flavor hadron-jet correlations in the nucleon/nucleus going direction at the future EIC. The proposed measurements will provide a unique path to explore the flavor dependent fragmentation functions and energy loss in heavy nucleus, which can constrain the initial state effects for previous and ongoing heavy ion measurements at the Relativistic Heavy Ion Collider (RHIC) and the Large Hadron Collider (LHC). Details of the proposed physics program will be discussed in this presentation.

**Primary authors:** Dr LI, Xuan (Los Alamos National Lab); Dr DURHAM, Matt (Los Alamos National Lab); Dr DA SILVA, Cesar (Los Alamos National Lab)

**Presenter:** Dr LI, Xuan (Los Alamos National Lab)

**Session Classification:** Posters

**Track Classification:** Future facilities

Contribution ID: 10

Type: **Poster**

## Small-x calculations with a Biased Ensemble

*Tuesday, 25 June 2019 16:13 (1 minute)*

Computing observables in high-energy collisions requires a functional average over the configurations of small-x gluons in the wave functions of the colliding hadrons. We discuss a method for performing biased averages, for example due to a multiplicity or centrality bias, where the gluon distributions of the hadrons are modified from their unbiased average. We consider specifically potential effects due to a bias on the correlator of two Wilson lines, i.e. the dipole scattering amplitude, and on azimuthal angular correlations of gluons at high transverse momentum (the “glasma graphs”).

**Primary authors:** Mr KAPILEVICH, Gary; Prof. DUMITRU, Adrian (Professor at Baruch college; Professor at CUNY GC; fellow at BNL)

**Presenter:** Mr KAPILEVICH, Gary

**Session Classification:** Posters

**Track Classification:** High pT probes of the initial state

Contribution ID: 12

Type: **Oral**

## Anisotropic hydrodynamics with a realistic collisional kernel

*Tuesday, 25 June 2019 15:40 (20 minutes)*

In this work, we implement an effective kinetic theory based scattering kernel in the anisotropic hydrodynamics (aHydro) formalism. We compare the realistic kernel results to those obtained from aHydro with the Anderson-Witting scattering kernel (RTA). For the purpose of this study, we consider a conformal system undergoing transversally-homogenous and boost-invariant Bjorken expansion. The collisional kernel is given by the leading order  $2 \leftrightarrow 2$  scattering kernel in the massless scalar  $\lambda\phi^4$ . We explicitly enforce number conservation through the incorporation of a dynamical chemical potential (fugacity) in the underlying aHydro distribution function and focus on the case of a system obeying classical statistics. We first compare the time evolution of the aHydro microscopic parameters and components of the energy-momentum tensor. Then, we determine the anisotropic non-equilibrium attractor for a system subject to this realistic collisional kernel. Our results indicate that when the near-equilibrium relaxation-times in the Anderson-Witting and scalar collisional kernels are matched, the aHydro dynamics receive quantitatively important corrections using the LO scalar kernel, however, the aHydro attractor itself is not substantially modified.

**Primary authors:** ALMAALOL, Dekrayat (Kent State University); STRICKLAND, Michael (Kent State University)

**Presenter:** ALMAALOL, Dekrayat (Kent State University)

**Session Classification:** Parallel: Approach to Equilibrium

**Track Classification:** Initial conditions for hydrodynamics & transport coefficients

Contribution ID: 13

Type: **Oral**

## Non-linear evolution in QCD at high-energy beyond leading order

*Tuesday, 25 June 2019 17:00 (20 minutes)*

The next-to-leading order (NLO) Balitsky-Kovchegov (BK) equation describing the high-energy evolution of the scattering between a dilute projectile and a dense target suffers from instabilities unless it is supplemented by a proper resummation of the radiative corrections enhanced by large transverse logarithms. Earlier studies have shown that if one expresses the evolution in terms of the rapidity of the dilute projectile, the dominant, anti-collinear, contributions can be resummed to all orders. However, in applications to physics, the results must be re-expressed in terms of the rapidity of the dense target (which corresponds to Bjorken  $x$ ). We show that although they lead to stable evolution equations, resummations expressed in the rapidity of the projectile show a strong, unwanted, scheme dependence when translated in the rapidity of the target. We circumvent this problem by working directly with the rapidity of the dense target [1]. This avoids the large anti-collinear contributions but introduces new, collinear, instabilities, which are however milder since disfavoured by the typical BK evolution. We propose several prescriptions for resumming these new double logarithms and find only little scheme dependence. The resummed equations are non-local in rapidity and can be extended to full NLO accuracy. We present the first applications of these resummed equations to deep inelastic scattering at HERA.

[1] B. Ducloué et al, e-Print: arXiv:1902.06637 [hep-ph]

**Primary authors:** DUCLOUE, Bertrand (IPhT Saclay); IANCU, Edmond (IPhT Saclay); MUELLER, Al (Columbia University); SOYEZ, Gregory (IPhT Saclay); TRIANTAFYLLOPOULOS, Dionysios (ECT\* Trento)

**Presenter:** DUCLOUE, Bertrand (IPhT Saclay)

**Session Classification:** Parallel: Forward/saturation/spin

**Track Classification:** Forward and saturation physics

Contribution ID: 14

Type: Oral

## Non-Gaussian fluctuations of $v_1, v_2, v_3$ and $v_4$ and their correlations in Pb+Pb collisions with the ATLAS detector

*Tuesday, 25 June 2019 17:40 (20 minutes)*

The measurements of the flow phenomena in Xe+Xe and Pb+Pb collisions provide an excellent opportunity to study the interplay of viscous effects – which diminish the azimuthal anisotropies more in Xe+Xe compared to Pb+Pb – and initial geometry fluctuations which have an opposite effect. With the recently developed techniques, used for suppression of non-flow correlations in small systems, applied to  $0.49 \text{ nb}^{-1}$  of Pb+Pb and  $3 \mu\text{b}^{-1}$  of Xe+Xe data significant reduction of non-flow biases is achieved with respect to the previous measurements both at high  $p_T$  and in peripheral collisions. An interesting scaling relationships is observed in the  $v_n$  across different centralities, where up to an overall scaling the  $v_n$  as a function of  $p_T$  have identical shapes. The origin of this scaling and its implications are discussed in the framework of hydrodynamic models. Multi-particle azimuthal cumulants measured in the Pb+Pb collisions provide information on the event-by-event fluctuations of harmonic flow coefficients  $v_n$  and correlated fluctuations between two harmonics  $v_n$  and  $v_m$ . For the first time, a non-zero four-particle cumulant is observed for dipolar flow,  $v_1$ . The four-particle cumulants for elliptic flow,  $v_2$ , and triangular flow,  $v_3$ , exhibit a strong centrality dependence and change sign in ultra-central collisions. Correlations between two harmonics are studied with three- and four-particle mixed-harmonic cumulants, which also decrease in strength towards central collisions and either approach zero or change sign in ultra-central collisions. To investigate the possible flow fluctuations arising from intrinsic centrality or volume fluctuations, the results are compared between two different event classes used for centrality definitions. In peripheral and mid-central collisions where the cumulant signals are large, only small differences are observed. In ultra-central collisions, the differences are much larger and transverse momentum dependent. These results provide new information to disentangle flow fluctuations from the initial and final states, as well as new insights on the influence of centrality fluctuations.

**Primary author:** ATLAS COLLABORATION**Presenter:** BEHERA, Arabinda (STAR)**Session Classification:** Parallel: Initial conditions for hydrodynamics & transport coefficients**Track Classification:** Initial conditions for hydrodynamics & transport coefficients

Contribution ID: 15

Type: **Poster**

## $v_n$ - $p_T$ correlations in 5.02 TeV Pb+Pb and p+Pb collisions with the ATLAS detector

*Tuesday, 25 June 2019 16:16 (1 minute)*

Recently collected data by the ATLAS experiment at the LHC offer opportunities to explore dynamic properties of quark-gluon plasma. A new tool to study these properties is the modified Pearson's correlation coefficient,  $\rho$ , that quantifies the correlation between the mean transverse momentum in an event,  $[p_T]$ , and the square of the flow harmonic magnitude,  $v_n^2$ . To suppress non-flow effects,  $v_n^2$  is calculated by correlating charged particles from two sub-events covering opposite pseudorapidity ranges of  $0.75 < |\eta| < 2.5$  while  $[p_T]$  is evaluated for particles with  $|\eta| < 0.5$ . The measurement of  $\rho$  is performed using minimum-bias p+Pb and Pb+Pb collisions at the same energy  $\sqrt{s_{NN}} = 5.02$  TeV allowing for a comparison of the medium dynamics in small and large systems. In Pb+Pb collisions, values of  $\rho$  coefficients are found to significantly deviate from zero for studied harmonics ( $v_2$ ,  $v_3$ , and  $v_4$ ). The coefficients as a function of centrality are observed to be weakly dependent on the transverse momentum range of the selected particles, despite large differences in the mean transverse momentum in an event and the magnitude of fluctuations of flow harmonics. The  $\rho$  coefficient in Pb+Pb collisions for the second order harmonics has a positive value for mid-central collisions and decreases in the most central events. In p+Pb collisions the  $\rho$  coefficient is measured only for the second order flow harmonics. In both p+Pb and peripheral Pb+Pb collisions, it is found to be negative. All measured coefficients are compared to theoretical models.

**Presenter:** BURKA, Klaudia (AGH-UST)**Session Classification:** Posters**Track Classification:** Collectivity in small systems



Contribution ID: 16

Type: Oral

## Recent ATLAS results on correlations in small collisions systems and photon-induced processes in ultra-peripheral Pb+Pb collisions at 5.02 TeV

*Wednesday, 26 June 2019 15:00 (20 minutes)*

This talk presents ATLAS measurements of azimuthal anisotropies in  $pp$  and  $p+Pb$  collisions, performed via two-, four- and six-particle correlations, with modifications to ensure suppression of correlations arising from jets and dijets. In  $pp$  collisions, the strength of the correlations quantified by the anisotropy parameter  $v_2$  does not show any dependence on the charged-particle multiplicity. Recent theoretical models suggest that this can be due to lack of correlation between the charged-particle multiplicity and the impact parameter of the  $pp$  collision. To test this hypothesis, correlation measurements are performed in  $pp$  collisions tagged by the presence of a Z boson – which acts as an independent handle on the impact parameter – and compared to inclusive  $pp$  collisions. Additionally, results of correlations between flow harmonics of different order in  $pp$  and  $p+Pb$  collisions measured via symmetric and asymmetric cumulants are also presented, and are shown to follow similar trends as those observed in Pb+Pb collisions. Measurements of HBT radii with respect to the second-order event-plane in  $p+Pb$  collisions are also presented. Azimuthal modulations in the HBT radii consistent with the hydrodynamic evolution of a short-lived medium are observed. The ultra-peripheral collisions (UPCs) of relativistic heavy ion beams lead to both photon-nucleus and photon-photon processes. The measurements of particle production in photo-nuclear reactions can shed light on the QCD dynamics of novel, extremely asymmetric colliding systems, with energies between those available at RHIC and the LHC. Understanding the hadronic fluctuation spectrum of the photon in this fashion is also critical for maximizing the precision of measurements at a future Electron Ion Collider facility. Finally, new measurements of light-by-light scattering with substantially reduced uncertainties will be presented in this talk. This process provides a precise and unique opportunity to investigate extensions to the Standard Model such as higher-dimension operators and axion-like particles.

**Primary author:** ATLAS COLLABORATION**Presenter:** SEIDLITZ, Blair (University of Colorado Boulder)**Session Classification:** Parallel: Collectivity in small systems 1**Track Classification:** Collectivity in small systems

Contribution ID: 17

Type: Oral

## ATLAS measurements of azimuthal anisotropy of heavy flavor hadrons in Pb+Pb, p+Pb and pp collisions

*Wednesday, 26 June 2019 14:20 (20 minutes)*

ATLAS measurements of azimuthal anisotropy and suppression of muons from heavy flavor decays in Pb+Pb collisions are presented. The measurements are extended to smaller systems of p+Pb and pp collisions, where no significant modification of the heavy flavor production are observed. In the smaller systems, a template fit method is used to subtract non-flow contributions using simultaneous fit to low and high charged-particle multiplicity samples. The heavy flavor flow in p+Pb is studied using multiple probes, including prompt  $D^0$  mesons,  $J/\psi$ , and muons from semi-leptonic decays of heavy flavor hadrons. In pp collisions, new measurements of flow coefficient of muons from heavy flavor decays are also presented. The observed heavy flavor azimuthal anisotropies in p+Pb and pp collisions are found to be qualitatively similar to those of light hadrons indicating a similar origin for both types of particles.

**Primary author:** ATLAS COLLABORATION**Presenter:** HILL, Kurt (University of Colorado)**Session Classification:** Parallel: Collectivity in small systems 1**Track Classification:** Collectivity in small systems

Contribution ID: 18

Type: Oral

## Jet and photon probes of small and large systems in ATLAS

*Wednesday, 26 June 2019 14:00 (20 minutes)*

Jets and photons have been studied to constrain the initial and final stages of collisions between two large nuclei at the Large Hadron Collider.

Measurements of photon and jet production  $p$ +Pb collisions are potentially sensitive to novel effects such as gluon saturation, the onset of non-linear QCD, and the energy loss of partons in the nuclear matter. In A+A collisions jets are modified as they pass through the hot nuclear matter. This talk presents recent ATLAS measurements of jets and photons in  $p$ +Pb, Pb+Pb and Xe+Xe collisions. Results on forward-forward and forward-central di-jet production in 5.02 TeV  $p$ +Pb and  $pp$  collisions in regions where the momentum fraction of a parton compared to a nucleon in the lead nucleus is small are presented. Also, measurements of photon production in 8.16 TeV  $p$ +Pb data over a large kinematic range is presented and compared to measurements in  $pp$  collisions and theoretical models. The comparison of di-jet balance in  $pp$ , Xe+Xe, and Pb+Pb collisions presented in this talk will provide information about the path-length dependence and role of fluctuations in the energy loss. Finally, measurements of the distributions of charged particles in and around jets as well as measurements of the energy and fragmentation functions of jets opposite photons are presented.

**Primary author:** ATLAS COLLABORATION

**Presenter:** PEREPELITSA, Dennis (University of Colorado Boulder)

**Session Classification:** Parallel: High pT probes of the initial state

**Track Classification:** High pT probes of the initial state

Contribution ID: 19

Type: Oral

## Heavy electroweak boson production in Pb+Pb collisions with ATLAS

*Tuesday, 25 June 2019 14:20 (20 minutes)*

Electroweak bosons provide a unique opportunity to extract the information about the beginning of the temporal evolution of the heavy-ion collision system and understand how the cold nuclear matter effects influence the observables that are measured in heavy-ion collisions.  $Z$  and  $W$  bosons decaying in leptonic channels are unaffected by the presence of the quark-gluon plasma and carry the information from the time when bosons were created, i.e. from the moment of the collision itself. Measurement of  $Z$  and  $W$  bosons allows to quantify the modification of the nuclear parton distribution functions and verify our understanding of the geometry of the colliding nuclei.

In the 2015 heavy-ion data-taking period at the LHC, the ATLAS experiment obtained 0.49/nb of the Pb+Pb data and 25/pb of the proton-proton data at the centre of mass energy of 5.02 TeV. The fully analysed data presented in this talk addresses the nuclear modification of the parton distribution functions PDF at a new level of precision. Comparison between the lead-lead and proton-proton systems gives an opportunity to subject the Glauber model used by all heavy ion experiments to a stringent test performed over a wide range of collision centralities.

**Primary author:** ATLAS COLLABORATION**Presenter:** DUMANCIC, Mirta (Weizmann Institute of Science)**Session Classification:** Parallel: nPDF/CNM**Track Classification:** nPDF, cold matter effects

Contribution ID: 26

Type: **Oral**

## Longitudinal fluctuations and decorrelations of anisotropic flows in relativistic heavy-ion collisions

*Tuesday, 25 June 2019 18:00 (20 minutes)*

We study the longitudinal decorrelations of elliptic, triangular and quadrangular flows in heavy-ion collisions at the LHC and RHIC energies. The event-by-event CLVisc (3+1)-dimensional hydrodynamics model, combined with the fully fluctuating AMPT initial conditions, is utilized to simulate the space-time evolution of the strongly-coupled quark-gluon plasma. Detailed analysis is performed for the longitudinal decorrelations of flow vectors, flow magnitudes and flow orientations. We find strong correlations between final-state longitudinal decorrelations of anisotropic flows and initial-state longitudinal structures and collision geometry: the decorrelation of elliptic flow shows a non-monotonic centrality dependence due to initial elliptic geometry, while the longitudinal flow decorrelations are typically larger in lower energy and less central collisions where the mean lengths of the string structure are shorter in the initial states.

**Primary authors:** QIN, Guang-You (Central China Normal University); WU, Xiang-Yu; PANG, Long-Gang; WANG, Xin-Nian (Lawrence Berkeley National Laboratory)

**Presenter:** QIN, Guang-You (Central China Normal University)

**Session Classification:** Parallel: Initial conditions for hydrodynamics & transport coefficients

**Track Classification:** Initial conditions for hydrodynamics & transport coefficients

Contribution ID: 27

Type: **Poster**

## Precision QCD with the LHeC and the FCC-eh

*Tuesday, 25 June 2019 16:11 (1 minute)*

The Large Hadron-electron Collider (LHeC) is a proposed upgrade of the LHC at CERN. An ERL will provide electrons to collide with the HL-LHC, HE-LHC and the FCC-hh proton beams to achieve centre-of-mass energies 1.3-3.5 TeV and luminosities  $10^{34} \text{ cm}^{-2}\text{s}^{-1}$ . They will enlarge the kinematic plane by more than one order of magnitude towards smaller  $x$  and larger  $Q^2$  than HERA. DIS measurements in such configurations offer unprecedented possibilities to enlarge our knowledge on parton densities through a complete unfolding of all flavours, both in a single experimental setup and combined with data from hadron colliders where precise factorisation tests can be performed. In this talk we review the most recent developments on the determination of proton PDFs and the measurement of  $\alpha_s$  at both the LHeC and the FCC-eh.

**Primary authors:** LHEC WORKING GROUP;; OLNESS, Fredrick (SMU)**Presenter:** OLNESS, Fredrick (SMU)**Session Classification:** Posters**Track Classification:** Future facilities

Contribution ID: 29

Type: **Oral**

## LHCb fixed target results and prospects

*Wednesday, 26 June 2019 17:30 (20 minutes)*

Among the main LHC experiments, LHCb is the only detector that can run both in collider and fixed-target mode. Internal gas targets of helium, neon and argon have been used so far to collect samples corresponding to integrated luminosities up to  $0.1 \text{ pb}^{-1}$ . An upgraded target, allowing for a wider choice of target gas species and increasing the gas density by up to two orders of magnitude, is going to be installed for the LHC Run 3. This offers a unique opportunity for measurements of great interest going from QCD to astroparticle in unexplored kinematic regions. Results and prospects on open and hidden charm productions will be presented, which can provide crucial constraints on cold nuclear matter effects and nPDF at large  $x$ .

**Primary author:** MUELLER, Katharina (Universitaet Zuerich (CH))

**Presenter:** DI NEZZA, Pasquale (INFN Frascati)

**Session Classification:** Parallel: Future facilities

**Track Classification:** Future facilities

Contribution ID: 33

Type: Oral

## A complete set of splitting functions in nuclear matter to any order in opacity and applications to jet physics

*Wednesday, 26 June 2019 15:40 (20 minutes)*

I will present the first calculation of all  $\mathcal{O}(\alpha_s)$  medium-induced branching processes to any order in opacity. The splitting functions results are given as iterative solutions to matrix equations with initial conditions set by the leading order branchings in the vacuum. The flavor and quark mass dependence of the in-medium  $q \rightarrow qg$ ,  $g \rightarrow gg$ ,  $q \rightarrow gq$ ,  $g \rightarrow q\bar{q}$  processes is fully captured by the light-front wavefunction formalism and the color representation of the parent and daughter partons. I will further present numerical results in a realistic QCD medium. The numerical simulations show that the second order in opacity corrections can change the energy dependence of the in-medium shower intensity. Corrections to the longitudinal and angular distributions of the in-medium splitting kernels that may have important implications for jet substructure phenomenology. Last but not least, I will show how these splitting function can be used to evaluate the modification of hadron jet production in SIDIS, such as at the future EIC.

**Primary author:** VITEV, Ivan (LANL)**Presenter:** VITEV, Ivan (LANL)**Session Classification:** Parallel: High pT probes of the initial state**Track Classification:** High pT probes of the initial state



Contribution ID: 34

Type: **Oral**

## Magnetic field in expanding quark-gluon plasma

*Tuesday, 25 June 2019 18:20 (20 minutes)*

Intense electromagnetic fields are created in the quark-gluon plasma by the external ultra-relativistic valence charges. The time-evolution and the strength of this field are strongly affected by the electrical conductivity of the plasma. Yet, it has recently been observed that the effect of the magnetic field on the plasma flow is small. We compute the effect of plasma flow on magnetic field and demonstrate that it is less than 10%. These observations indicate that the plasma hydrodynamics and the dynamics of electromagnetic field decouple. Thus, it is a very good approximation, on the one hand, to study QGP in the background electromagnetic field generated by external sources and, on the other hand, to investigate the dynamics of magnetic field in the background plasma. We also argue that the wake induced by the magnetic field in plasma is negligible.

**Primary author:** TUCHIN, Kirill**Presenter:** TUCHIN, Kirill**Session Classification:** Parallel: Forward/saturation/spin**Track Classification:** Vorticity and polarization

Contribution ID: 35

Type: **Oral**

## Stochastic hydrodynamics and long time tails of a non-equilibrium fluid

*Tuesday, 25 June 2019 14:20 (20 minutes)*

We investigate the impact of hydrodynamic fluctuations on correlation functions of a relativistic fluid with a conserved U(1) charge. The kinetic equations for the two-point functions of pressure, momentum and heat energy densities are derived within the framework of stochastic hydrodynamics. The leading non-analytic contributions to the energy-momentum tensor as well as the U(1) current are determined from the solutions to these kinetic equations. In the case of a static homogeneous background we show that the long time tails obtained from hydro-kinetic equations reproduce the one-loop results derived from statistical field theory. We use these results to establish bounds on transport coefficients. We generalize the stochastic equation to a scale invariant background flow undergoing Bjorken expansion. We compute the leading fractional power  $\mathcal{O}((\tau T)^{-3/2})$  correction to the U(1) current and compare with the first order gradient term. Finally, we discuss the extension of stochastic hydrodynamics to study the effects of critical behavior of the heat conductivity, shear and bulk viscosities in heavy ion collisions for a system close to the QCD critical point.

**Primary author:** Dr MARTINEZ , Mauricio (North Carolina State University)

**Co-authors:** Prof. SCHAEFER, Thomas (North Carolina State University); Dr SKOKOV, Vladimir (North Carolina State University)

**Presenter:** Dr MARTINEZ , Mauricio (North Carolina State University)

**Session Classification:** Parallel: Approach to Equilibrium

**Track Classification:** Approach to equilibrium (weak & strong coupling)

Contribution ID: 36

Type: **Poster**

## Non-perturbative aspects of hydrodynamization for the far-from-equilibrium Bjorken flow

*Tuesday, 25 June 2019 16:14 (1 minute)*

In relativistic kinetic theory, the one-particle distribution function is approximated by an asymptotic perturbative power series in Knudsen number which is divergent. For the Bjorken flow, we expand the distribution function in terms of its moments and study their nonlinear evolution equations. The resulting coupled dynamical system can be solved for each moment consistently using a multi-parameter transseries which makes the constitutive relations inherit the same structure. A new non-perturbative dynamical renormalization scheme is born out of this formalism that goes beyond the linear response theory. As a result, the transport coefficients get dynamically renormalized at every order in the time-dependent perturbative expansion by receiving non-perturbative corrections present in the transseries. The renormalized transport coefficients feature a transition to their equilibrium fixed point, which is a neat diagnostics of transient non-Newtonian behavior. Furthermore, we show that the first dissipative correction to the distribution function is not only determined by the known effective shear viscous term but also a new high energy non-hydrodynamic mode. Finally, we briefly discuss some possible phenomenological applications of the proposed non-hydrodynamic transport theory.

**Primary author:** Mr SHI, Haosheng (North Carolina State University)

**Co-authors:** Dr MARTINEZ , Mauricio (North Carolina State University); Mr BEHTASH, Alireza (North Carolina State University); Dr KAMATA, Syo (North Carolina State University)

**Presenter:** Mr SHI, Haosheng (North Carolina State University)

**Session Classification:** Posters

**Track Classification:** Approach to equilibrium (weak & strong coupling)

Contribution ID: 38

Type: **Poster**

## Linear and non-linear response of two-particle correlations to initial-geometry fluctuations

*Tuesday, 25 June 2019 16:08 (1 minute)*

We investigate the importance of different features of the initial geometry to anisotropic flow fluctuations in heavy-ion collisions. To that end, we explore the hydrodynamic response of differential flow harmonics  $v_n(p_T)$  to generalized eccentricities  $\epsilon_{n,m}$  of the initial density profile within a realistic hydrodynamic model. Special attention is paid to two-particle angular correlations, characterized in detail by the principal-component analysis (PCA). We address the relevance of non-linear response, as well as the stability of the results against the inclusion of extra eccentricities. Additionally, we study new effects from multiplicity fluctuations, which could lead to redundancies in the experimental PCA data.

**Primary authors:** Dr HIPPERT, Mauricio (Universidade de Sao Paulo); Dr DOBRIGKEIT CHINEL-LATO, David (Universidade Estadual de Campinas); Dr LUZUM, Matthew (Universidade de Sao Paulo); Dr NORONHA, Jorge (Universidade de Sao Paulo); Dr NUNES DA SILVA, Tiago (Universidade Federal de Santa Catarina); Dr TAKAHASHI, Jun (Universidade Estadual de Campinas)

**Presenter:** Dr HIPPERT, Mauricio (Universidade de Sao Paulo)

**Session Classification:** Posters

**Track Classification:** Initial conditions for hydrodynamics & transport coefficients

Contribution ID: 45

Type: **Oral**

## Initial state fluctuations in Pythia 8

*Wednesday, 26 June 2019 17:50 (20 minutes)*

Initial state geometry has proved to be decisively important for quantitative descriptions of collective behavior in large collision systems, such as PbPb and AuAu. One of the most remarkable lessons from LHC has been the discovery of collective behavior in small collision systems, but here the notion of spatial structure is not as obvious as in nuclear collisions. In Monte Carlo event generators, ad hoc phenomenological schemes are usually employed with clear room for improvement. In this talk we present progress on including a transverse space structure of pp collisions based on the Mueller dipole formulation of QCD, into the Pythia8 event generator. This formalism has the advantage that parameters can be estimated from inclusive quantities in ep and pp collisions, such that the spatial structure becomes a true prediction of the model. Besides the importance for collective behavior, in particular in pp, but also in fluctuation dominated pA and peripheral AA collisions, the dipole picture also serves as an important starting point for including electron-ion initial states in the model for heavy-ion collisions in Pythia8, the Angantyr framework, a perspective which also will be discussed.

**Primary authors:** RASMUSSEN, Christine (Lund University); BIERLICH, Christian (Lund University & University of Copenhagen)

**Presenter:** RASMUSSEN, Christine (Lund University)

**Session Classification:** Parallel: Collectivity in small systems 2

**Track Classification:** Collectivity in small systems

Contribution ID: 46

Type: **Oral**

## Computing the gluon Sivers function at low-x

*Wednesday, 26 June 2019 16:50 (20 minutes)*

The TMD parton distributions can have azimuthal asymmetry in the transverse plane for the transversely polarized nucleon. This is called the Sivers effect and is phenomenologically important for the description of single spin asymmetries. I will present our recent calculation of the gluon Sivers function at small-x obtained by using the known connection between the Sivers function and the odderon.

**Primary author:** Dr HATTA, Yoshitaka**Co-author:** Dr HATTA, Yoshitaka (BNL)**Presenter:** Dr HATTA, Yoshitaka (BNL)**Session Classification:** Parallel: Future facilities**Track Classification:** 3D nucleon structure

Contribution ID: 48

Type: **Oral**

## New paradigm for fluctuations in heavy-ion collisions

*Tuesday, 25 June 2019 18:40 (20 minutes)*

We present a first-principles description of the primordial state of relativistic nucleus-nucleus collisions, whose density fluctuations and anisotropy we evaluate in the color glass condensate (CGC) framework of high-energy QCD. Relating the primordial anisotropy of the system in our approach to the measured final-state anisotropic flow through a simple linear mapping, we achieve an excellent description of both RHIC and LHC data.

Our description does not make any explicit reference to the usual, ad hoc fluctuations due to the random positions of the incoming nucleons: Primordial initial-state fluctuations are generated solely by McLerran-Venugopalan correlators of color charges. The good agreement found with the data implies, then, that QCD interactions alone can provide the system with enough density fluctuations to explain the measured triangular flow, and elliptic flow fluctuations.

This suggests a fundamental paradigm shift in our understanding of fluctuations in heavy-ion collisions: At ultrarelativistic energies, the standard Monte Carlo Glauber picture of nuclear collisions, which until now has been understood as the dominant source of fluctuations, can be abandoned.

Based on:

<https://arxiv.org/abs/1902.07168>

**Primary author:** GIACALONE, Giuliano (IPhT - Saclay)

**Co-authors:** OLLITRAULT, Jean-Yves (Université Paris-Saclay); Dr LUZUM, Matthew (Universidade de Sao Paulo); MARQUET, Cyrille (Ecole Polytechnique); GUERRERO-RODRÍGUEZ, Pablo (Ecole Polytechnique)

**Presenter:** GIACALONE, Giuliano (IPhT - Saclay)

**Session Classification:** Parallel: Initial conditions for hydrodynamics & transport coefficients

**Track Classification:** Initial conditions for hydrodynamics & transport coefficients

Contribution ID: 49

Type: **Oral**

## Self-similarity and spectral functions of non-Abelian plasmas in 2+1D

*Tuesday, 25 June 2019 14:40 (20 minutes)*

To better understand the dynamics of initial stages in heavy-ion collisions, we perform classical-statistical simulations of SU(2) gauge theory in 2+1 dimensions. We find that highly occupied non-Abelian plasmas approach the same 2+1D self-similar state at late times of their far-from-equilibrium evolution, irrespective of details of their initial conditions, and we determine the scaling exponents. We extract the spectral function non-perturbatively and show that for larger momenta than the mass scale  $p \gg m$ , one sees a pronounced peak in the frequency domain, while at low momenta  $p < m$ , quasi-particle assumptions become invalid. The hard-thermal loop (HTL) formalism is not applicable to 2+1D gauge theories at low momenta  $p \leq m$ , and indeed, our results are inconsistent with its predictions. This challenges our detailed understanding of plasma instabilities at initial stages that is mostly based on HTL calculations of highly anisotropic gluonic matter.

**Primary authors:** Dr BOGUSLAVSKI, Kirill (Vienna University of Technology); Prof. KURKELA, Aleksi (CERN); Prof. LAPPI, Tuomas (University of Jyväskylä); Dr PEURON, Jarkko (ECT\*)

**Presenter:** Dr PEURON, Jarkko (ECT\*)

**Session Classification:** Parallel: Approach to Equilibrium

**Track Classification:** Approach to equilibrium (weak & strong coupling)



Contribution ID: 50

Type: **Oral**

## EPR paradox and quantum entanglement at sub-nucleonic scales

*Tuesday, 25 June 2019 18:00 (20 minutes)*

In quantum mechanics, Einstein, Podolsky, and Rosen (EPR) formulated an apparent paradox of quantum theory in 1935. They considered two quantum mechanical systems were first brought to interaction, then later separated to large distance. A measurement of a physical observable in one system would have an immediate effect on the conjugate observable in the other system, even when they are causally disconnected. Therefore, EPR concluded that there is an inconsistency in the quantum theory. In the parton model formulated by Bjorken, Feynman, and Gribov, the partons inside of a nucleon are viewed as “quasi-free” particles when they are boosted into the infinite momentum frame, where the parton probed by the virtual photon is causally disconnected from the rest of the nucleon. Since the parton and the rest of the nucleon have to form a color-singlet state due to confinement, we encounter the EPR paradox at sub-nucleonic states for the first time. In this work we propose a resolution of this apparent paradox via quantum entanglement. We test this idea by measuring the entanglement entropy of the system using data from proton-proton collisions at the Large Hadron Collider, and our results provide a strong direct indication of quantum entanglement at sub-nucleonic scales.

**Primary author:** TU, Zhoudunming (BNL)**Co-authors:** KHARZEEV, Dmitri (Stony Brook University and BNL); ULLRICH, Thomas (BNL)**Presenter:** TU, Zhoudunming (BNL)**Session Classification:** Parallel: Forward/saturation/spin**Track Classification:** Forward and saturation physics

Contribution ID: 53

Type: **Oral**

## Measurement of electroweak-boson production in p–Pb and Pb–Pb collisions at the LHC with ALICE

*Tuesday, 25 June 2019 14:00 (20 minutes)*

Electroweak bosons are created in the hard scattering processes at the initial stage of heavy-ion collisions and they are insensitive to the presence of the strongly-interacting medium. This makes them clean probes of the initial-state effects in heavy-ion collisions, such as the nuclear modification of the Parton Distribution Functions (nPDFs). Furthermore, their measurement in heavy-ion collisions is a powerful test of the binary scaling of hard processes as well as a reference for hot-matter effects on other probes.

The measurement of electroweak-boson production in p–Pb and Pb–Pb collisions at the LHC provides constraints on the nPDFs of (anti)quarks in phase-space regions which are poorly constrained from previous experiments. At forward rapidity ( $2.5 < y < 4$ ), ALICE can measure W and Z bosons via their muon decay in all collision systems provided by the LHC. These measurements are complementary to those by ATLAS and CMS at central rapidity.

In this contribution, focus will be given to the most recent ALICE electroweak-boson measurements. Exploiting the data collected by ALICE in 2015 and 2018, centrality and rapidity-differential measurements of the Z-boson production yield in Pb–Pb collisions at  $\sqrt{s_{NN}} = 5.02$  will be discussed. The first measurement of the Z-boson production cross-section in p–Pb collisions at  $\sqrt{s_{NN}} = 8.16$  TeV will also be shown as a function of rapidity. The status of ongoing W-boson analyses in various collision systems will also be reported. All the presented results will be compared to theoretical calculations including nPDFs.

**Primary author:** ANONYMOUS, submission (ALICE Collaboration)

**Presenter:** VALLE, Nicolo (Universita and INFN, Pavia (IT))

**Session Classification:** Parallel: nPDF/CNM

**Track Classification:** nPDF, cold matter effects

Contribution ID: 55

Type: **Poster**

## Particle production as a function of system size and underlying-event activity measured with ALICE at the LHC

ALICE has performed several measurements aimed at understanding the heavy-ion-like patterns observed in small collision systems. New approaches can be helpful to clarify particle production mechanisms in pp collisions, as well as the similarities observed among the systems created in pp, p-A and A-A collisions.

In this talk we report on charged-particle transverse momentum distributions as a function of event multiplicity. The distributions are obtained using a 2D-unfolding procedure. We compare unidentified charged-particle production at different collision energies, as well as that for pp, p-Pb and Pb-Pb collisions at the same energy. In order to understand the role of autocorrelations in small systems, it has been proposed to exploit the usage of the underlying event as a multiplicity estimator to factorize the hardest and the softer components of the events. This approach can also be used to study collective effects in events with exceptionally large activity in the underlying-event region with respect to the event-averaged mean. For this purpose, in this talk we also present the charged particle transverse momentum distributions as a function of underlying-event activity in pp collisions. All results will be compared with QCD-inspired event generators, as well as with existing measurements adopting the mid- and forward-pseudorapidity multiplicity estimators.

**Primary author:** ANONYMOUS, submission (ALICE Collaboration)

**Presenter:** ANONYMOUS, submission (ALICE Collaboration)

**Session Classification:** Posters

**Track Classification:** Collectivity in small systems

Contribution ID: 57

Type: **Oral**

## A Forward Rapidity Upgrade for the STAR Detector

*Wednesday, 26 June 2019 17:10 (20 minutes)*

The STAR Collaboration plans to design, construct, and install a suite of new detectors in the forward rapidity region ( $2.5 < \eta < 4$ ) over the next two years, enabling a program of novel measurements in pp, pA, and AA collisions. This extension of STAR's kinematic reach will allow detailed studies of cold QCD physics at both very high and very low partonic momentum fraction, i.e. when the colliding quarks and gluons carry very large or very small amounts of the nucleon energy. Previous STAR efforts using the Forward Pion Detector (FPD) and Forward Meson Spectrometer (FMS) detectors have demonstrated that there are outstanding QCD physics opportunities in the forward rapidity region. To fully explore these physics opportunities, a forward upgrade [1] with detection capability for neutral pions, photons, electrons, jets, and leading hadrons is proposed which adds charged-particle tracking, electromagnetic, and hadronic calorimetry to STAR's capabilities at high pseudorapidity. The upgrade will greatly expand the kinematic reach for ongoing measurements of the spin and flavor structure of the nucleon, and will enable studies of the longitudinal structure of the nuclear initial state that leads to breaking of boost invariance in heavy-ion collisions. Transport properties of the hot and dense matter formed in heavy-ion collisions will also become accessible with the proposed measurement capabilities at forward rapidity. Details on the proposed upgrade and the scientific opportunities it will enable will be presented.

[1] "The STAR Forward Calorimeter System and Forward Tracking System," <https://drupal.star.bnl.gov/STAR/starnotes/publications>

**Primary author:** BRANDENBURG, Daniel (Brookhaven National Laboratory)

**Presenter:** BRANDENBURG, Daniel (Brookhaven National Laboratory)

**Session Classification:** Parallel: Future facilities

**Track Classification:** Future facilities

Contribution ID: 60

Type: Oral

## Measurement of open heavy-flavour hadron production in pp, p-Pb and Pb-Pb collisions with ALICE

*Tuesday, 25 June 2019 15:20 (20 minutes)*

Heavy quarks (charm and beauty) are effective probes to test perturbative QCD-based calculations in pp collisions and to study cold nuclear matter (CNM) effects such as gluon saturation, shadowing,  $k_T$  broadening and energy loss in CNM in p-Pb collisions.

With the ALICE detectors, open heavy flavours are measured via the full reconstruction of hadronic decays of non-strange D mesons,  $D_s^+$  mesons and  $\Lambda_c^+$  baryons, and the semi-electronic decay channels of electrons from open charm and open beauty hadron decays and  $\Xi_c^0$  at mid-rapidity and, via the semi-muonic decays at forward rapidity. The measurements of  $\Lambda_c^+$  and  $\Xi_c^0$  production can shed light on charm hadronisation mechanisms in the absence of the hot and dense QCD medium. The comparisons of charmed baryon-to-meson ratios with models tuned to  $e^+e^-$  collisions allow us to examine the possible contributions of charm quark recombination or coalescence, which may be more prevalent at large charged-particle multiplicities, and to potentially disentangle these effects from pure vacuum fragmentation. Recent observations in pp and p-Pb collisions showed remarkable similarities with Pb-Pb collisions, which might suggest the presence of collectivity. To further explore the origin the collective-like effects observed in pp and p-Pb collisions, the study of open heavy-flavour production as a function of the charged-particle multiplicity naturally links soft and hard processes that occur in the collision and allows one to study their interplay.

In this contribution, the production cross sections of D mesons and open heavy-flavour decay electrons measured at mid-rapidity, and open heavy-flavour decay muons measured at forward rapidity in pp collisions at  $\sqrt{s} = 5.02$  TeV with ALICE detector will be presented. The latest ALICE results on  $\Lambda_c^+$  and  $\Xi_c^0$  production and  $\Lambda_c^+/D^0$  ratio in pp collisions at  $\sqrt{s} = 7$  TeV and in p-Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV, and  $\Xi_c^0/D^0$  ratio in pp collisions at  $\sqrt{s} = 7$  TeV and their comparison to model calculations and to equivalent results from the light-flavour sector will be discussed. The results of beauty production using beauty-decay electrons and non-prompt D mesons in pp collisions at  $\sqrt{s} = 5.02$  TeV and the self-normalized yield of open heavy-flavour decay electrons and muons as a function of multiplicity in pp and p-Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV will be presented. Finally, the nuclear modification factor ( $Q_{pPb}$ ) of D mesons in p-Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV and the comparison of the experimental results with theoretical models will be discussed as well.

**Primary author:** ALICE COLLABORATION

**Presenter:** FAGGIN, Mattia

**Session Classification:** Parallel: nPDF/CNM

**Track Classification:** nPDF, cold matter effects

Contribution ID: 61

Type: **Poster**

## **Non-equilibrium Green's functions for energy-momentum perturbations around Bjorken flow from the Boltzmann equation in relaxation time approximation**

*Tuesday, 25 June 2019 16:01 (1 minute)*

Non-equilibrium Green's functions provide an efficient tool to describe the evolution of the energy-momentum tensor during the early time pre-equilibrium stage, and provide a meaningful to address the question when and to what extent a hydrodynamic description of the system becomes applicable. We present a calculation of the Green's functions describing the evolution of energy density perturbations in the transverse plane, based on the Boltzmann equation in relaxation time approximation. We discuss the approach towards viscous hydrodynamics along with the emergence of various scaling phenomena for conformal systems. By comparing our results obtained in the relaxation time approximation to previous calculations in QCD kinetic theory, we further address the question which macroscopic features of the energy momentum tensor are sensitive to the underlying microscopic dynamics.

**Primary authors:** SCHLICHTING, Soeren (University of Washington); Dr MARTINEZ , Mauricio (North Carolina State University); Dr KAMATA, Syo (North Carolina State University)

**Presenters:** SCHLICHTING, Soeren (University of Washington); Dr MARTINEZ , Mauricio (North Carolina State University)

**Session Classification:** Posters

**Track Classification:** Approach to equilibrium (weak & strong coupling)

Contribution ID: 63

Type: **Oral**

## Measurement of heavy-flavour jets and correlations and elliptic flow in small systems with ALICE

*Wednesday, 26 June 2019 15:20 (20 minutes)*

The ALICE Collaboration studied extensively heavy-flavour production at mid- and forward rapidities in small systems at the LHC. The data provide precise tests for pQCD calculations based on the factorization approach and set constraints to Cold Nuclear-Matter effects that can modify heavy-flavour production in p-Pb collisions with respect to pp collisions. At mid-rapidity the study of the angular correlation of heavy-flavour particles with charged particles produced in the event allows us to further characterize charm and beauty production and fragmentation processes in pp collisions and investigate their possible modifications due to nuclear effects in p-Pb and Pb-Pb collisions. In p-Pb collisions these studies can also set constraints on the dependence of Cold Nuclear-Matter effects on the collision geometry and on the density of final-state particles. In addition, the study of the angular correlation pattern in p-Pb collisions as a function of the event multiplicity allows studying the features of long-range correlations similar to the one found in heavy-ion collisions whose origin is still debated. Finally, the measurement of heavy-flavour jets, besides constituting the necessary baseline for similar studies in the Pb-Pb collision system, gives more direct access to the initial parton kinematics and can provide further constraints on pQCD based models.

In this contribution, the latest ALICE results from pp and p-Pb collisions collected during the LHC Run 2 will be presented. In particular, measurements of the angular correlation of D mesons and heavy-flavour decay electrons with charged particles in pp and p-Pb collisions will be shown. The measurement of heavy-flavour decay leptons elliptic flow in p-Pb collisions at high multiplicity at mid- and forward rapidities will also be presented together with the D-meson central to peripheral ratio ( $Q_{CP}$ ). Finally, ALICE measurements on D-tagged jets will be discussed both in pp and p-Pb collisions together with the status of b-jet measurements. All the results shown will be compared with theoretical models.

**Primary author:** ALICE COLLABORATION**Presenter:** COLAMARIA, Fabio (INFN, Sezione di Bari (IT))**Session Classification:** Parallel: High pT probes of the initial state**Track Classification:** Collectivity in small systems

Contribution ID: 64

Type: **Poster**

## Toward a full mapping of the hydrodynamic response to initial conditions

*Tuesday, 25 June 2019 16:07 (1 minute)*

Anisotropic flow is well understood as a hydrodynamic response to spatial anisotropies in the system density at early times. This response function can be written explicitly as a systematic expansion in terms of length scales, such that the leading contribution is the familiar eccentricity  $\varepsilon_n$  that represents global structure. These relations have allowed for direct connections to be made between the initial state and experimental data, and constraints to be put on the initial stages.

However, the initial conditions for hydrodynamics consist of an entire tensor  $T^{\mu\nu}$  (as well as conserved currents). Although they are thought to be less important than energy density, other components such as momentum density and shear tensor can also contribute, and their affects should have increasing importance for smaller collision systems. It is therefore interesting to extend the response framework in order to probe these aspects of the initial stages and their affect on flow observables.

I will present a framework for including the effects of the full hydrodynamic initial conditions, along with numerical tests from full hydrodynamic simulations to demonstrate its efficacy. In addition, I will present an extension to include rapidity dependence.

**Primary authors:** LUZUM, Matthew (Universidade de São Paulo); SOUSA, Jefferson (Universidade de São Paulo); FRANCO, Rodrigo (Universidade de São Paulo); Dr NORONHA, Jorge (Universidade de Sao Paulo)

**Presenter:** LUZUM, Matthew (Universidade de São Paulo)

**Session Classification:** Posters

**Track Classification:** Initial conditions for hydrodynamics & transport coefficients



Contribution ID: 65

Type: Oral

## Quarkonium production in pp and p–Pb collisions with ALICE at the LHC

*Tuesday, 25 June 2019 15:00 (20 minutes)*

Quarkonia, bound states of either a charm and anti-charm quark pair (e.g.  $J/\psi$ ,  $\psi(2S)$ ), or a beauty and anti-beauty quark pair ( $\Upsilon(1S)$ ,  $(2S)$  and  $(3S)$ ), are considered a prominent tool to study the properties of the Quark-Gluon Plasma (QGP) formed in high-energy heavy-ion collisions such as those delivered by the LHC. However, their production is also sensitive to so called initial state effects, such as the modifications of the parton distribution functions in the nucleus or the occurrence of gluon saturation at low Bjorken  $x$ , and thus provide some insight on the initial conditions of such collisions.

Recently, proton-proton (pp) and proton-lead (p–Pb) collisions with high charged-particle multiplicities have been found to exhibit phenomena similar to those attributed to the QGP formation. Measuring quarkonia in such collisions could contribute to a better understanding of the underlying physics processes leading to these observations and in any case provide a more detailed understanding of their production mechanism.

ALICE has measured quarkonium production in both pp collisions for collision energies  $\sqrt{s}$  ranging from 2.76 to 13 TeV and p–Pb collisions at center of mass energies per nucleon-nucleon collisions  $\sqrt{s_{NN}} = 5.02$  and 8 TeV. These measurements have been carried out down to zero transverse momentum and at both mid and forward rapidity. Beyond quarkonium cross sections in pp collisions and nuclear modification factors in p–Pb collisions as a function of rapidity, transverse momentum and centrality, this presentation will also focus on correlations between the quarkonium and the underlying event, including recent results on the charged-particle multiplicity dependence of quarkonium relative yields and mean transverse momentum at both mid and forward rapidity,  $J/\psi$ -hadron correlations at mid rapidity and  $J/\psi$  azimuthal anisotropy in high-multiplicity p–Pb collisions.

**Primary author:** ANONYMOUS, submission (ALICE Collaboration)

**Presenter:** HAYASHI, Shinichi (CNS, the University of Tokyo)

**Session Classification:** Parallel: nPDF/CNM

**Track Classification:** nPDF, cold matter effects

Contribution ID: 67

Type: **Poster**

## Heavy-quark diffusion coefficient in out-of-equilibrium plasmas

*Tuesday, 25 June 2019 16:06 (1 minute)*

We study the heavy-quark momentum diffusion coefficient in gluon plasmas in a self-similar regime using real-time lattice techniques. We observe that the time-evolution of the momentum diffusion coefficient is consistent with a  $t^{-5/7}$  power law, as predicted by HTL perturbation theory and self-similarity. Using HTL with our recently acquired data on the spectral function of over-occupied gluodynamics, we find that the main contribution to the diffusion coefficient arises from the longitudinal Landau damping in the spectral function and study further features of the signal. Finding consistent results between lattice simulations and HTL results is an important step forward in understanding the evolution of the diffusion coefficient at initial stages.

**Primary authors:** Dr PEURON, Jarkko (European Centre for Theoretical Studies in Nuclear Physics and Related Areas (ECT\*)); Prof. LAPPI, Tuomas (University of Jyväskylä); Dr KURKELA, Aleks (CERN); BOGUSLAVSKI, Kirill (Vienna University of Technology)

**Presenter:** Dr PEURON, Jarkko (European Centre for Theoretical Studies in Nuclear Physics and Related Areas (ECT\*))

**Session Classification:** Posters

**Track Classification:** Initial conditions for hydrodynamics & transport coefficients

Contribution ID: 68

Type: **Poster**

## Measurement of long-range correlations in pp collisions at $\sqrt{s} = 13$ TeV with ALICE at the LHC

*Tuesday, 25 June 2019 16:23 (5 minutes)*

The observed azimuthal modulations of long-range correlations in pseudorapidity in small systems like pp or p-Pb collisions show strikingly similar features to those seen in heavy ion collisions. Many theoretical approaches to interpreting this effect have been developed. However, it is still unclear whether these long-range correlations are due to final or initial state effects.

To further investigate these effects, we studied long-range correlations as a function of transverse momentum in very high multiplicity pp collisions at  $\sqrt{s} = 13$  TeV, collected with the high multiplicity event trigger during 2016 and 2017 with ALICE. In this talk, we present the near side per-trigger yield at large pseudorapidity separation (ridge yield) as a function of transverse momentum in pp collisions at  $\sqrt{s}=13$  TeV. The results will be compared to previous measurements from CMS and ATLAS. In addition, we present the ridge yield in events where harder fragmentation processes are present, to explore possible physical origins of long range correlations.

**Primary author:** KIM, Junlee**Presenter:** KIM, Junlee**Session Classification:** Posters**Track Classification:** Collectivity in small systems

Contribution ID: 71

Type: **Oral**

## Holographic collisions with baryon number at intermediate coupling

*Tuesday, 25 June 2019 15:20 (20 minutes)*

In holographic heavy ion collisions it is possible to follow both the energy density and a globally conserved (baryon) number density throughout the evolution, dual to a metric and Maxwell field in the dual bulk theory, respectively. At infinite coupling, past work has shown that after the collision the baryon number ends up around mid-rapidity, which is different from high energy heavy ion collisions. In this talk I will present first results for the flow of baryon number in holographic collisions away from the infinite coupling limit, which also means we study the collisions at a larger shear viscosity over entropy density ratio than  $1/4\pi$ . I will give an introduction to the holographic set-up, which is interesting since it contains an extra coupling of the gravitational Ricci scalar with the Maxwell field. Remarkably, depending on the value of this extra coupling, we find that the flow of baryon number during the collision can be affected drastically. In particular, we find that at intermediate coupling it is possible for almost no baryon number to end up at mid-rapidity. We further show how the matter produced in the collision relaxes into a flow as described by hydrodynamics with a conserved baryon current.

**Primary authors:** VAN DER SCHEE, Wilke (Utrecht University); Prof. RAJAGOPAL, Krishna (MIT); Mr FOLKESTAD, Åsmund (MIT); Dr GROZDANOV, Sašo (MIT)

**Presenter:** VAN DER SCHEE, Wilke (Utrecht University)

**Session Classification:** Parallel: Approach to Equilibrium

**Track Classification:** Approach to equilibrium (weak & strong coupling)

Contribution ID: 76

Type: **Poster**

## Flow vs Nonflow in Ultracentral U-U and Be-Be Collisions

*Tuesday, 25 June 2019 16:17 (1 minute)*

An ongoing question in the field is if the collectivity originating in small systems arises from:

- Flow scenario: initial conditions coupled to relativistic hydrodynamics leading to independent particle emissions from the fluid hyper-surface.

or

- Non-flow scenario: genuine 2,4 etc particle correlations from, e.g. a saturation framework.

We note that, in general, the experimentally measured  $v_n\{m\}$ 's can include non-trivial contributions from both pictures (and mixing terms between the two) so we caution that the calculations of  $v_n\{m\}$  must consider both contributions, which look different in each framework.

Using the Color-Glass Condensate (CGC) formalism in the (semi)dilute-(semi)dilute regime for  $p_T \gg Q_s$ , we can compare the ultra-central scaling of azimuthal anisotropies with the multiplicities of deformed ion-ion collisions. We consider deformed ion-ion collisions as a testing bed for these comparisons because due to either tip-tip or side-side collisions, one expects the geometrical shape immediately after the collision (eccentricities) to scale inversely with the final multiplicity. Because hydrodynamics is predominately driven by linear response in ultracentral collisions, this inverse scaling of  $v_2\{2\}$  with  $dN/dy$  is preserved in the final hydrodynamic picture. In contrast to hydrodynamics, in the CGC framework  $v_2\{2\}$  and  $v_3\{2\}$  increase monotonically with the multiplicity. We repeat these calculations for Beryllium-Beryllium collisions and predict the same effect. Different parameterizations of Uranium are also studied and constrained by data. Thus, we argue that deformed ions can be a perfect testing ground for comparing the CGC to hydro pictures in small systems. Additionally, we calculate other flow observables such as  $(v_2\{4\}/v_2\{2\})^4$  and symmetric cumulants,  $NSC(3,2)$ . We find that due to the suppression of 4 particle correlations in our framework that the CGC picture produces an imaginary  $v_2\{4\}/v_2\{2\}$  while we find a positive  $NSC(3,2)$ .

**Primary author:** WERTEPNY, Douglas (Ben Gurion University of the Negev)

**Co-authors:** NORONHA-HOSTLER, Jacquelyn (Rutgers University); SIEVERT, Matthew (Rutgers University); RAO, Skanda (Rutgers undergraduate physics major, sophomore); Dr LUZUM, Matthew (Universidade de Sao Paulo); Mr CARZON, Patrick (Rutgers University); NUNES, Tiago (Universidade Estadual de Campinas)

**Presenter:** RAO, Skanda (Rutgers undergraduate physics major, sophomore)

**Session Classification:** Posters

**Track Classification:** Collectivity in small systems

Contribution ID: 77

Type: **Poster**

## A Principal Component Analysis of event-by-event fluctuations in hydrodynamic simulations at the LHC

Hydro simulations are compared with recent experimental data by CMS, on a Principle Component Analysis. The trends for the scaled leading and sub-leading components for  $n=2,3$  agree with data. In contrast, for  $n=0$  there is a qualitative disagreement: the leading component increases with  $p_T$  while it is constant in data. Using a toy model where the principal components can be computed analytically, it is shown that the  $n=0$  leading and sub-leading components are extremely dependent on fluctuations in  $N$  and  $\bar{p}_T$  as well as their covariance and provide a new and clean way to study fluctuations and rule out initial condition models.

**Primary authors:** Prof. GARDIM, Fernando (Universidade Federal de Alfenas-Brazil); GRASSI, Frederique (Universidade de São Paulo-Brazil); OLLITRAULT, Jean-Yves (Université Paris-Saclay); Dr LUZUM, Matthew (Universidade de Sao Paulo); Mr ISHIDA, Pedro (Universidade de São Paulo)

**Presenter:** GRASSI, Frederique (Universidade de São Paulo-Brazil)

**Session Classification:** Posters

**Track Classification:** Initial conditions for hydrodynamics & transport coefficients

Contribution ID: 79

Type: **Oral**

## Investigation of collectivity in small collision systems with ALICE

*Wednesday, 26 June 2019 15:20 (20 minutes)*

Measurements of anisotropic flow in heavy-ion collisions are an important tool to investigate the nature of the created collectively expanding medium called the Quark-Gluon Plasma (QGP). Recently, striking similarities have been observed in numerous measurements in high multiplicity proton-proton and proton-lead collisions, where no such medium was expected.

In this talk, we will present the latest ALICE measurements of flow coefficients, and their magnitude correlations using Symmetric Cumulants for charged particles in pp collisions at  $\sqrt{s} = 13$  TeV, p-Pb at  $\sqrt{s_{NN}} = 5.02$  TeV, Xe-Xe at  $\sqrt{s_{NN}} = 5.44$  TeV and Pb-Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV, collected during the Run 2 LHC data taking period. In addition, the flow coefficients of identified particles in p-Pb collisions will be presented. Non-flow effects, which are azimuthal correlations not originating from a common symmetry plane, are suppressed with a pseudorapidity separation and a subtraction method. Both methods are particularly important for studies of collectivity in small systems. The results from a broad spectrum of colliding systems and wide range of multiplicity are compared to various theoretical models, providing a deep insight into initial conditions and the nature of collective phenomena in different collision systems.

**Primary author:** ALICE COLLABORATION**Presenter:** PACIK, Vojtech (Niels Bohr Institute, University of Copenhagen, Denmark)**Session Classification:** Parallel: Collectivity in small systems 1**Track Classification:** Collectivity in small systems

Contribution ID: 80

Type: Oral

## Hydrodynamics far-from-equilibrium: a concrete example in kinetic theory

*Tuesday, 25 June 2019 17:00 (20 minutes)*

The applicability of hydrodynamical models in the extreme conditions produced in heavy ion collisions has not yet been properly understood theoretically. This happens mostly because the derivation of hydrodynamics from microscopic theory often relies on the assumption that the system is sufficiently close to equilibrium – something difficult to justify in the rapidly expanding systems created in heavy ion collisions. In this talk we propose a more general derivation of relativistic hydrodynamics from kinetic theory, in which the fluid is assumed to be close to an *isotropic nonequilibrium* state instead of an equilibrium one. We demonstrate that, for a wide variety of nonequilibrium states, a hydrodynamic theory that is identical to the traditional hydrodynamic equations applied to heavy ion collisions is obtained. The only difference appears in the form of the transport coefficients that enter the equations of motion. Simulations of the Boltzmann equation in 0+1D Bjorken flow in the ultrarelativistic limit are performed to demonstrate this effect, showing that the same evolution for the energy-momentum tensor is obtained even when the momentum distribution function of partons is very different from an equilibrium one.

**Primary authors:** SILVEIRA DENICOL, Gabriel (Universidade Federal Fluminense); Dr NORONHA, Jorge (Universidade de Sao Paulo)

**Presenter:** SILVEIRA DENICOL, Gabriel (Universidade Federal Fluminense)

**Session Classification:** Parallel: Initial conditions for hydrodynamics & transport coefficients

**Track Classification:** Initial conditions for hydrodynamics & transport coefficients



Contribution ID: 81

Type: **Oral**

## Connecting far-from-equilibrium hydrodynamics, resummed transport coefficients, and attractor solutions

*Tuesday, 25 June 2019 14:00 (20 minutes)*

After briefly reviewing recent developments in the field, we show how far-from-equilibrium hydrodynamics may be systematically defined, for arbitrary flow profiles, in terms of a generalized tensorial expansion with transport coefficients that contain an all order resummation in gradients. We discuss how this approach naturally relates to *hydrodynamic attractor* solutions found both at weak and strong coupling. In this formulation, the transport coefficients of far-from-equilibrium fluid dynamics depend not only on the microscopic properties of the system but also on the nonlinear properties of the underlying state of the fluid itself. In contrast to previous works, no additional assumptions about the symmetries of the flow are necessary. An example of this proposal is constructed using Israel-Stewart theory and, in this case, the novel resummed transport coefficients decrease with increasing Knudsen number according to formulas that can be readily investigated in current numerical simulations of the quark-gluon plasma, which can be especially relevant to small collision systems.

**Primary authors:** NORONHA, Jorge (Universidade de Sao Paulo); SILVEIRA DENICOL, Gabriel (Universidade Federal Fluminense)

**Presenter:** NORONHA, Jorge (Universidade de Sao Paulo)

**Session Classification:** Parallel: Approach to Equilibrium

**Track Classification:** Approach to equilibrium (weak & strong coupling)

Contribution ID: 82

Type: Oral

# Principal Component Analysis of collective flow in Heavy-Ion collisions

*Tuesday, 25 June 2019 18:20 (20 minutes)*

Principal Component Analysis (PCA) is a mathematical tool that can capture the most important information in data. As one of the unsupervised algorithms of machine learning, PCA is good at discovering modes or hidden patterns in huge amount of data. It has seen successful applications of PCA in computer vision, data science and physics. Compared with deep learning algorithms, the advantage of PCA lies in its simple and elegant mathematical formulation, which is understandable and traceable. In this talk, we implement PCA to analyze collective flow in Relativistic Heavy-Ion Collisions.

In the first part [1], we demonstrate the ability of PCA to automatically discover flow without any guidance from human beings. PCA is applied to particle yields distribution as a function of transverse plane angle  $\varphi$  in the reaction plane. The eigenmodes decomposed by PCA are similar to, but not identical with traditional Fourier bases. Furthermore, we define new flow harmonics with PCA modes and the new ones serve as better linear predictors for initial eccentricities than traditional ones. Specifically, correlations between same and different harmonics of initial and final states increase and decrease respectively, showing smaller mode-mixing effects.

In the second part [2], as another application of PCA, we study factorization breaking in two-particle correlation  $V_{n\Delta}(p_{T1}, p_{T2})$  with respect to transverse momentum  $p_T$  [3]. In particular, we focus on the sub-leading flow, which hopefully sheds light on different sources in initial geometries [4]. However, the stability and interpretation of PCA results have to be re-examined. We design different tests to explore the limitations of PCA, arguing that improper choice of  $p_T$  range and weight matrix might lead to confusing and inconsistent results. As a consequence, these mentioned technical issues should be addressed before we could come to any conclusions or truly understand physics from PCA results.

[1] Z. Liu, W. Zhao and H. Song, in preparation.

[2] Z. Liu, A. Behera, H. Song and J. Jia, in preparation.

[3] CMS Collaboration, Phys.Rev. C.96.064902

[4] A. Mazeliauskas and D. Teaney, Phys.Rev. C91 (2015) no.4, 044902

**Primary authors:** LIU, ZIMING (Peking University); SONG, Huichao (Peking); JIANGYONG, Jia (stony brook university & BNL)

**Presenter:** LIU, ZIMING (Peking University)

**Session Classification:** Parallel: Initial conditions for hydrodynamics & transport coefficients

**Track Classification:** Initial conditions for hydrodynamics & transport coefficients

Contribution ID: 83

Type: Oral

## Measurement of elliptic and triangular flow with multiparticle correlations in pPb collisions at 8.16 TeV

*Wednesday, 26 June 2019 16:30 (20 minutes)*

The second- and third-order azimuthal anisotropy Fourier harmonics are studied in pPb collisions at  $\sqrt{s_{\text{NN}}} = 8.16$  TeV over a wide range of event multiplicities. Multiparticle correlations are used to isolate global properties stemming from the collision overlap geometry. The second-order, elliptic harmonic moment is obtained with high precision through four-, six-, and eight-particle correlations and, for the first time, the third-order, triangular harmonic moment is studied using four-particle correlations. A sample of peripheral PbPb collisions at  $\sqrt{s_{\text{NN}}} = 5.02$  TeV that covers a similar range of event multiplicities to the pPb results is also analyzed. Model calculations of initial-state fluctuations in pPb and PbPb collisions can be directly compared to the high precision experimental results. This work provides new insight on the fluctuation-driven geometry at the earliest stages of heavy ion collisions.

**Primary author:** PETRUSHANKO, Sergey (Moscow State University)

**Presenter:** TUO, Shengquan (Vanderbilt University (US))

**Session Classification:** Parallel: Collectivity in small systems 2

**Track Classification:** Collectivity in small systems

Contribution ID: 84

Type: **Poster**

## Higher order flow correlations and their non-linear modes in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV with ALICE

*Tuesday, 25 June 2019 16:09 (1 minute)*

One of the primary goals of flow studies in heavy-ion collisions during recent years is a better understanding of the transport properties of the quark-gluon plasma (QGP), such as the temperature dependence of the shear viscosity to entropy ratio,  $\eta/s(T)$ . Flow observables, such as the higher order harmonics ( $n>3$ ) and their non-linear responses to the initial state anisotropy have a strong potential to constrain  $\eta/s(T)$  because of different sensitivities for various stages of heavy-ion collisions. These observables have been published in Pb-Pb collisions at  $\sqrt{s_{\text{NN}}} = 2.76$  TeV by the ALICE Collaboration. However, only harmonics up to the fifth order have been measured with good precision, leaving the orders highly sensitive to  $\eta/s$  as predicted un-investigated.

In this talk, we present the measurements of the symmetry-plane correlations and the non-linear coefficients up to the eighth harmonic order in Pb-Pb collisions at  $\sqrt{s_{\text{NN}}} = 5.02$  TeV for unidentified charged hadrons. In addition, we present the results of  $p_{\text{T}}$ -differential non-linear flow modes for charged pions, kaons, and (anti-)protons. The results are compared to the lower energy measurements at 2.76 TeV and calculations from state-of-the-art hydrodynamic models.

**Primary author:** ALICE COLLABORATION**Presenter:** PARKKILA, Jasper (University of Jyväskylä)**Session Classification:** Posters**Track Classification:** Initial conditions for hydrodynamics & transport coefficients

Contribution ID: 85

Type: **Poster**

## Early stage momentum anisotropies and electromagnetic probes of quark-gluon plasma

*Tuesday, 25 June 2019 16:05 (1 minute)*

Various microscopic models suggest that local rest frame momentum anisotropies can be large during the early stages of evolution of the quark-gluon plasma (QGP). In recent years, the framework of relativistic anisotropic hydrodynamics (aHydro) has been developed in order to incorporate momentum anisotropic distributions of the QGP constituents into the phenomenological studies of ultra-relativistic heavy-ion collision experiments. In this talk, the question of how much we can learn about the early-time momentum anisotropies by studying the yield and flow of electromagnetic probes will be addressed. In particular, we compare the sensitivity of hadronic and electromagnetic probes to the initial momentum anisotropy used in hydrodynamic calculations.

**Primary authors:** SALEHI KASMAEI, Babak (Kent State University); STRICKLAND, Michael (Kent State University)

**Presenter:** SALEHI KASMAEI, Babak (Kent State University)

**Session Classification:** Posters

**Track Classification:** Initial conditions for hydrodynamics & transport coefficients

Contribution ID: 86

Type: **Oral**

## Measurement of bottomonia in pp, pPb and PbPb collisions at 5.02 TeV with the CMS detector

Recent measurements of the  $\Upsilon(1S)$ ,  $\Upsilon(2S)$ , and  $\Upsilon(3S)$  mesons in pp, pPb and PbPb collisions at 5.02 TeV are presented. The analysis was performed as a function of rapidity and transverse momentum. In addition, the dependence on the event activity and collision centrality is studied in pPb and PbPb collisions, respectively. New results of the upsilon production in pPb collisions will be reported, compared with the results from PbPb collisions. In this presentation, the results are discussed in terms of the ‘cold nuclear matter’ effects in pPb collisions and sequential melting scenario in dense partonic matter, as well as the effect from recombination of uncorrelated quarks. The results are also compared with theory models, which can help to improve and constrain the theoretical calculations.

**Primary author:** PETRUSHANKO, Sergey (Moscow State University)

**Presenter:** PETRUSHANKO, Sergey (Moscow State University)

**Track Classification:** High pT probes of the initial state

Contribution ID: 88

Type: **Oral**

## Constraining nPDFs with Z boson and Drell-Yan measurements in pPb collisions with CMS

*Tuesday, 25 June 2019 14:40 (20 minutes)*

Nuclear parton distribution functions (nPDFs) of quarks and antiquarks affect the production electroweak bosons in proton-lead (pPb) collision. In this presentation the measurement of Z bosons in pPb collision at center of mass energies of 5.02 TeV and 8.16 TeV with CMS is presented. The rapidity dependence is particularly sensitive to nPDFs, but further information can be gained by studying the mass dependence of DY production, measured for the first time in pPb collisions at 8.16 TeV, down to 15 GeV. In addition, differential measurements in the dimuon  $p_T$  or  $\phi^*$  (an angular variable correlated with  $p_T$ , measured for the first time in pPb) provide insights on soft gluon emission at low  $p_T$ . Finally, prospects on Z boson production with future HL-LHC data are presented.

**Primary author:** PETRUSHANKO, Sergey (Moscow State University)**Presenter:** CHAPON, Émilien (CERN)**Session Classification:** Parallel: nPDF/CNM**Track Classification:** nPDF, cold matter effects

Contribution ID: 89

Type: **Poster**

## Studies of the top quark production in nuclear collisions and impact on nuclear PDFs in CMS

*Tuesday, 25 June 2019 16:10 (1 minute)*

In proton-nucleus collisions, the top quark is a novel and theoretically precise probe of the nuclear gluon density at high virtualities  $Q^2 \approx m_{\text{top}}$  and in the less explored high Bjorken- $x$  region. The first observation of the inclusive  $t\bar{t}$  production has been performed using  $174 \pm 6 \text{ nb}^{-1}$  of data in pPb collisions at  $\sqrt{s_{\text{NN}}} = 8.16 \text{ TeV}$ . However, the total uncertainty of about 17% is not sufficient for imposing constraints on current nPDF parameterizations, the dominant source of uncertainty in the theoretical prediction of  $\sigma(\text{pPb} \rightarrow t\bar{t} + X)$ . The prospects of measuring  $\sigma$  differentially have recently been examined and a feasibility study of the measurement with the CMS detector at the High-Luminosity LHC (HL-LHC) era is therefore carried out as a function of the reconstructed lepton  $p_T$  and rapidity. The relative statistical uncertainty in both variables is found to be at the level of 4–5% in each bin, and it is expected to be the dominant uncertainty at  $\sqrt{s_{\text{NN}}} = 8.16 \text{ TeV}$  for an integrated luminosity scenario of  $2 \text{ pb}^{-1}$ . The motivations for measurements of top quarks in nucleus-nucleus collisions are multifold and are discussed: the top quark decay products are sensitive to the energy loss of heavy quarks, and a probe of the space-time structure of the QGP at HL-LHC.

**Primary author:** PETRUSHANKO, Sergey (Moscow State University)**Presenter:** LINDSEY, Cole (University of Kansas)**Session Classification:** Posters**Track Classification:** High  $p_T$  probes of the initial state



Contribution ID: 91

Type: **Poster**

## Influence of neutron skin in heavy ion collisions at RHIC energies

*Tuesday, 25 June 2019 16:00 (1 minute)*

A key ingredient to model heavy ion collisions dynamically is the initial spatial distribution of protons and neutrons inside the nucleus. Traditionally in most theoretical calculations there is no difference between them and their positions are sampled with the Woods-Saxon distribution. However, this assumption has been invalidated by experimental measurements [1]: The diffusiveness of the neutron distribution is larger than the proton one, resulting in a larger amount of neutrons distributed on the outer layers of the nucleus.

We implement the neutron skin together with nucleon-nucleon correlations, in the initial nuclear distribution of the SMASH transport model [2]. With this new state-of-the-art nuclear parametrization [3] we look at different collision systems (Pb+Pb, Zr+Zr, Ru+Ru) at RHIC energies i.e.  $7.7 < \sqrt{s} < 200 \text{ GeV}$ . We compute eccentricity distributions, charge and isospin densities, and the corresponding magnetic fields. As expected, we observe an increased number of neutron-neutron interactions for peripheral collisions leading to a modification of electric charge dependent observables. In addition, we found a  $\sim 10\%$  enhancement of the strength of the magnetic field. The implication of this result for the Chiral Magnetic Effect searches with the isobar run at RHIC is discussed.

[1] Phys. Rev. Lett. 112, 242502

[2] Phys. Rev. C 94, 054905

[3] arXiv:1811.10078v1

**Primary author:** Mr HAMMELMANN, Jan (FIAS Frankfurt Institute for Advanced Studies, Institute for theoretical physics Frankfurt)

**Co-authors:** Dr SOTO ONTOSO, Alba (Brookhaven National Laboratory); Dr ALVIOLI, Massimiliano (Istituto di Ricerca per la Protezione Idrogeologica, Consiglio Nazionale delle Ricerche); Prof. ELFNER, Hannah (FIAS Frankfurt Institute for Advanced Studies, Institute for theoretical physics Frankfurt, 3GSI Helmholtzzentrum für Schwerionenvorschung)

**Presenter:** Mr HAMMELMANN, Jan (FIAS Frankfurt Institute for Advanced Studies, Institute for theoretical physics Frankfurt)

**Session Classification:** Posters

**Track Classification:** 3D nucleon structure

Contribution ID: 92

Type: **Oral**

## IP-Jazma critical assessment of physics attributions

*Wednesday, 26 June 2019 14:00 (20 minutes)*

The IP-Jazma model was constructed to elucidate which features of calculations in the color glass condensate framework are attributable to simple geometry and scaling, and which are manifestations of more complex physics. In this talk we detail comparisons with CGC calculations in the dilute-dense limit, in the dense-dense limit (via IP-Glasma), and more phenomenological models such as Trento. Our result indicate that many results attributed in the literature to complex QCD phenomena are in fact dominated by simple geometric effects.

**Primary author:** Dr NAGLE, Jamie (University of Colorado Boulder)

**Co-authors:** Dr LIM, Sanghoon (University of Colorado Boulder); Dr ZAJC, Bill (Columbia University)

**Presenters:** Dr ZAJC, Bill (Columbia University); ZAJC, William (Columbia University)

**Session Classification:** Parallel: Collectivity in small systems 1

**Track Classification:** Collectivity in small systems

Contribution ID: 94

Type: **Oral**

## Measurements of nuclear parton distribution functions using dijets, forward jets, and photo-nuclear jets at the CMS detector & prospects for measurements in Run III

*Tuesday, 25 June 2019 17:20 (20 minutes)*

Measurements of dijet production and photo-nuclear interactions in heavy-ion collisions probe several nuclear mechanisms. In particular, dijet measurements in pPb collisions have been shown to be one of the most important tools for constraining the gluon nuclear parton distribution functions (PDFs) at large Bjorken- $x$ . Dijet production in pp and pPb collisions at a nucleon-nucleon centre-of-mass energy of 5.02 TeV is reported with the data samples collected with the Compact Muon Solenoid detector at the Large Hadron Collider. The dijet pseudorapidity distributions are measured as a function of dijet average transverse momentum in order to study the nuclear modifications of PDFs at various factorization scales. The final results from pp and pPb data samples are compared with next-to-leading-order perturbative QCD predictions obtained from both nucleon and nuclear PDFs. A significant modification of dijet pseudorapidity distributions in pPb collisions with respect to the measured pp reference is observed which indicates that the gluon PDF in lead ions is modified and the results are incompatible with predictions with DSSZ PDF without gluon EMC effects. Photo-nuclear jets are also measured in pp and pPb collision systems. The yield and angular correlation of low- $p_T$  jets at forward rapidity,  $5.0 < |\eta| < 6.5$ , are studied using the CASTOR calorimeter, which is sensitive to PDFs at low values of  $x$  and  $Q^2$ . The prospects of future measurements of forward and ultra-peripheral jets in various collision systems as well as dijet production in pPb at 8.16 TeV and in Run III will also be discussed.

**Primary author:** PETRUSHANKO, Sergey (Moscow State University)**Presenter:** BYLINKIN, Alexander (The University of Kansas (US))**Session Classification:** Parallel: Forward/saturation/spin**Track Classification:** Future facilities

Contribution ID: 95

Type: **Oral**

## First measurement of Diffraction in pPb collisions and recent results on ultra-peripheral heavy-ion processes

*Tuesday, 25 June 2019 17:40 (20 minutes)*

In this talk, first results on diffraction measurements in pPb collisions will be presented. This measurement utilizes the larger rapidity acceptance of the CMS detector using forward calorimeters such as the forward hadronic calorimeters HF, CASTOR and ZDC. This measurement provides important information for models related to the initial state in pPb collisions, and can also be used to tune Monte Carlo event generators of cosmic ray physics. In addition, recent results on exclusive vector meson photoproduction of  $\rho^0$ ,  $J/\psi$  and  $\Upsilon$  in both pPb and PbPb collisions will be presented. The measured integrated and differential cross sections as a function of rapidity and transverse momentum will be compared to theoretical models.

**Primary author:** PETRUSHANKO, Sergey (Moscow State University)

**Presenter:** WALCZAK, Marek (University of Warsaw)

**Session Classification:** Parallel: Forward/saturation/spin

**Track Classification:** Forward and saturation physics

Contribution ID: 96

Type: Oral

## Partonic spatial imaging at an electron-ion collider

*Wednesday, 26 June 2019 18:10 (20 minutes)*

The 2015 U.S. Nuclear Physics Long-Range Plan recommended the realization of an electron-ion collider (EIC) as the next large construction project in the United States. A U.S.-based EIC has also recently been endorsed by the U.S. National Academy of Sciences.

With the design of an EIC, advancements in theory and further development of phenomenological tools, we are now preparing for the next step in subnuclear tomographic imaging. The collider's large range of center-of-mass energy, in combination with very high luminosity and polarization of both the lepton and the hadron beams, will open a unique opportunity for very high precision measurements of both cross sections and spin-asymmetries. This will allow us for a detailed investigation of the partonic substructure of hadrons in multi-dimensions, as well as addressing the role of orbital angular momentum with respect to the nucleon spin.

Generalized parton distributions (GPDs) describe the multi-dimensional partonic structure of a nucleon in coordinate space, providing new information about the internal dynamics of quarks and gluons. Extraction of GPDs from hard exclusive processes and all related probes, is a pillar of the EIC science program.

This talk will highlight key measurements, experimental challenges, and finally discuss the EIC's expected impact over the current knowledge of the partonic multidimensional structure of hadrons in space coordinates.

**Primary author:** FAZIO, Salvatore (Brookhaven National Laboratory)

**Presenter:** LEE, J.H. (Brookhaven National Laboratory)

**Session Classification:** Parallel: Future facilities

**Track Classification:** 3D nucleon structure

Contribution ID: 97

Type: **Oral**

## Semi-inclusive Deep-Inelastic Scattering, Parton Distributions and \\ Fragmentation Functions at a Future Electron-Ion Collider

*Wednesday, 26 June 2019 17:50 (20 minutes)*

We present a quantitative assessment of the impact a future Electron-Ion Collider would have in the determination of parton distribution functions in the proton and parton-to-hadron fragmentation functions through semi-inclusive deep-inelastic electron-proton scattering data. Specifically, we estimate the kinematic regions for which the forthcoming data are expected to have the most significant impact in the precision of these distributions, computing the respective correlation and sensitivity coefficients. Using a reweighting technique for the sets of simulated data with their realistic uncertainties for two different center-of-mass energies, we analyse the resulting new sets of parton distribution functions and fragmentation functions, which have significantly reduced uncertainties.

**Primary authors:** Dr ASCHENAUER, E. C. (BNL); BORSA SANJUAN, Ignacio (Universidad de Buenos Aires); SASSOT, Rodolfo (Universidad de Buenos Aires); VAN HULSE, Charlotte (University of the Basque Country –UPV/EHU, Spain)

**Presenter:** BORSA SANJUAN, Ignacio (Universidad de Buenos Aires)

**Session Classification:** Parallel: Future facilities

**Track Classification:** Future facilities

Contribution ID: 98

Type: Oral

## Measurement of rapidity-odd directed flow for $D^0$ and $\overline{D}^0$ mesons using the STAR detector at RHIC

Heavy quarks, owing to their large masses, are predominantly created in the initial hard scatterings in heavy-ion collisions. Therefore, they can play a crucial role in probing the initial-time dynamics in these collisions. Hydrodynamic model calculations suggest that the rapidity-odd directed flow ( $v_1(y)$ ) of particles produced at mid-rapidity can originate from a tilt in the reaction plane of the thermalized matter caused by the asymmetry between the number of participants from projectile and target nuclei as a function of rapidity. Recently, it has been predicted that the slope of the directed flow at mid-rapidity of  $D^0$  mesons, arising from the transport of charm quarks in the tilted medium, can be several times larger than that of light flavor hadrons. A notable feature from the model calculation is the enhanced sensitivity of the  $D$ -meson  $v_1$  slope ( $dv_1/dy$ ), compared to that of light flavor hadrons, to the initial longitudinal profile of the QGP source. It has also been predicted that the transient electromagnetic field at early times can induce a much larger charge dependent directed flow for heavy quarks than for light quarks.

The Heavy Flavor Tracker (HFT) detector at STAR has demonstrated an excellent performance in reconstructing  $D^0$  and  $\overline{D}^0$  via hadronic decay channels. In this talk, we will report on the first evidence for a non-zero rapidity-odd directed flow for  $D^0$  and  $\overline{D}^0$  mesons in 10-80% central Au+Au collisions at  $\sqrt{s_{NN}} = 200$ -GeV using high statistics data collected with the HFT during the 2014 and 2016 RHIC runs. The average  $dv_1/dy$  for  $D^0$  and  $\overline{D}^0$  mesons is  $-0.081 \pm 0.021 \pm 0.017$ , while that of charged kaons is  $-0.0030 \pm 0.0001 \pm 0.0002$ , suggesting a significantly larger slope of the  $D^0$  mesons. The results will be compared to model calculations and physics implications will also be discussed.

**Primary author:** SINGHA, Subhash (Kent State University)

**Presenter:** SINGHA, Subhash (Kent State University)

**Session Classification:** Parallel: Initial conditions for hydrodynamics & transport coefficients

**Track Classification:** Initial conditions for hydrodynamics & transport coefficients

Contribution ID: 101

Type: Oral

## Probing collision dynamics of small system collisions via high $p_T$ hadrons and direct photons by the PHENIX experiment at RHIC

*Wednesday, 26 June 2019 14:40 (20 minutes)*

The recent finding of collective behavior of particles emitted in small system collisions calls for reexamination of such systems to determine whether or effects beyond the initial state nuclear effects play a significant role.

High  $p_T$  hadrons from jets and direct photons have been powerful probes to explore the dynamics of the heavy ion collisions, since they are produced at the early stages of the collisions. While photons do not interact strongly and therefore escape the medium largely unmodified, the high  $p_T$  hadrons originate from color charged objects that lose significant energy while traversing the medium. Taken together, these provide significant evidence in support of QGP formation in heavy ion collisions. Measurements of these two probes in small system collisions may provide insight to the dynamics of small collision systems and help assess their apparent similarity to heavy ion collisions.

PHENIX has measured mid to high  $p_T$  hadrons over wide range of rapidity in  $p/d/{}^3\text{He}+A$  collisions. PHENIX also has measured direct photons in  $p/d+A$  collisions. In this talk, the latest results on the high  $p_T$  hadrons and direct photons are presented and the possible additional effects explored by these probes will be discussed.

**Primary author:** SAKAGUCHI, Takao (BNL)

**Presenter:** SAKAGUCHI, Takao (BNL)

**Session Classification:** Parallel: High  $p_T$  probes of the initial state

**Track Classification:** High  $p_T$  probes of the initial state



Contribution ID: 103

Type: Oral

## How to infer the shape of the QGP droplet from the data

*Wednesday, 26 June 2019 15:00 (20 minutes)*

We propose an approach to extract the spatial anisotropy of QGP formed in ultrarelativistic heavy-ion collisions from measured high-pt observables  $R_{AA}$  and  $v_2$ . We show, through analytical arguments, numerical calculations, and comparison with experimental data, that  $v_2/(1 - R_{AA})$  reaches a well-defined saturation value at high  $p_\perp$ , which is in turn proportional to the initial anisotropy. We provide first anisotropy estimates from our approach, and compare them with predictions of various (fundamentally unrelated) initial state models. With expected future significant reduction of experimental errors, the anisotropy extracted from experimental data will strongly constrain the calculations of initial particle production in heavy-ion collisions and thus test our understanding of QGP physics.

**Primary authors:** DJORDJEVIC, Magdalena (Institute of Physics Belgrade); Mr STEFAN, Stojku (Institute of Physics Belgrade); Prof. DJORDJEVIC, Marko (University of Belgrade); Dr HUOVINEN, Pasi (Institute of Physics Belgrade)

**Presenter:** DJORDJEVIC, Magdalena (Institute of Physics Belgrade)

**Session Classification:** Parallel: High pT probes of the initial state

**Track Classification:** High pT probes of the initial state

Contribution ID: 104

Type: Oral

## PHENIX measurements of muon pairs from $c\bar{c}$ , $b\bar{b}$ , and Drell-Yan in p+p and p+Au at 200 GeV

*Tuesday, 25 June 2019 15:40 (20 minutes)*

In this talk we present the measurement of the muon pair continuum in p+p and p+Au collisions at a center of mass energy of 200 GeV. Our novel analysis technique enables the isolation of correlated pairs from semi-leptonic decays of charm and bottom hadrons and from the Drell-Yan process. The measured azimuthal correlations of muon pairs from heavy flavor decays are used to constrain the relative contributions of different production mechanisms of  $c\bar{c}$  and  $b\bar{b}$  pairs in p+p collisions. For bottom production, data from p+Au places limits on possible cold nuclear modifications. Measuring the in Drell-Yan cross-section in p+p and p+A collisions constrains nuclear parton distribution functions and furthers our understanding of initial state effects.

**Primary author:** DREES, Axel (Stony Brook University)**Presenter:** DREES, Axel (Stony Brook University)**Session Classification:** Parallel: nPDF/CNM**Track Classification:** nPDF, cold matter effects

Contribution ID: 106

Type: Oral

## Centrality dependence of collectivity in kinetic theory

*Wednesday, 26 June 2019 15:40 (20 minutes)*

To what extent are fluid-dynamic or particle-like excitations at the origin of the flow phenomena observed in pp, pA and AA collisions? And how does the interplay between these two sources of collectivity change as a function of system size and energy density? Here, we address this question in a simple transport theory that interpolates between free-streaming and viscous fluid dynamics. We discuss how this transport theory accounts for the centrality dependence of  $v_2$  and  $v_3$  between 1% and 90% centrality. Investigating the properties of the energy-momentum tensor evolved in this kinetic theory, we conclude that kinetic theory is consistent with a fluid dynamic picture of central PbPb collisions at the LHC, but that it strongly deviates from such a picture in peripheral PbPb collisions.

based on A. Kurkela, U.A. Wiedemann and B. Wu, arXiv:1803.02072 arXiv:1805.04081 and work in preparation.

**Primary authors:** Dr KURKELA, Aleksi (CERN); WIEDEMANN, Urs; WU, Bin (The Ohio State University)

**Presenters:** WIEDEMANN, Urs; Dr WIEDEMANN, Urs (CERN)

**Session Classification:** Parallel: Collectivity in small systems 1

**Track Classification:** Collectivity in small systems

Contribution ID: 107

Type: Oral

## Relaxation dynamics of chiral transports and spin polarization in Quark-Gluon Plasma

*Tuesday, 25 June 2019 18:40 (20 minutes)*

We study relaxation dynamics of chiral transport phenomena and spin polarization in Quark-Gluon Plasma in both weakly and strongly coupled regimes. These relaxation dynamics determine the important dynamical time scale for achieving equilibrium spin-polarization of quasi-particles in the presence of magnetic field and fluid vorticity, which are time-dependent in heavy-ion collisions. This is also important in the time-dependence of the Chiral Vortical Effect. Our results should be crucial in a reliable quantitative study of  $\Lambda$  baryon polarization in off-central heavy-ion collisions.

**Primary authors:** LI, Shiyong (UIC); YEE, Ho-Ung (University of Illinois at Chicago)

**Presenter:** LI, Shiyong (UIC)

**Session Classification:** Parallel: Forward/saturation/spin

**Track Classification:** Vorticity and polarization

Contribution ID: **109**Type: **Poster**

## Forward-backward centrality fluctuation in heavy ion collisions

*Tuesday, 25 June 2019 16:19 (1 minute)*

Centrality fluctuations is one of the main uncertainties for interpreting the centrality dependence of many experimental observables. The centrality fluctuation is constrained by selection based on particle multiplicity in a reference subevent, and contributes to observables measured in another subevent. Due to the asymmetry between forward- and backward-going participating nucleons, the number of sources for particle production is a function of  $\eta$  even in a single event, which leads to centrality decorrelation between different rapidities. Using a Glauber-based independent source model, we study the influence of centrality decorrelations on multi-particle cumulants. In mid-central collisions, a general relation is established between the multiplicity fluctuation and resulting centrality resolution in given subevent. In ultra-central collisions, where distribution of particle production sources is strongly distorted, we find these cumulants exhibit different behavior, due to observable-dependent non-Gaussianity in the underlying distributions. Furthermore, we investigate the influence centrality fluctuations in HIJING and UrQMD models. This study can be considered as a first step towards detailed understanding of the longitudinal dynamics for particle production in heavy-ion collisions.

**Primary author:** JIANGYONG, Jia (stony brook university & BNL)

**Co-author:** Mr ZHANG, Chunjian (Stony Brook University, Shanghai Institute of Applied Physics)

**Presenter:** JIANGYONG, Jia (stony brook university & BNL)

**Session Classification:** Posters

**Track Classification:** Initial conditions for hydrodynamics & transport coefficients

Contribution ID: 111

Type: **Poster**

## Examination of Flow and Non-Flow Factorization Methods in Small Collision Systems

*Tuesday, 25 June 2019 16:18 (1 minute)*

Two particle correlations have been used extensively to study hydrodynamic flow patterns in heavy-ion collisions. In small collision systems, such as p+p and p+A, where particle multiplicities are much smaller than in A+A collisions, non-flow effects from jet correlations, momentum conservation, particle decays, etc. can be significant, even when imposing a large pseudorapidity gap between the particles. A number of techniques to subtract the non-flow contribution have been developed by experiments at the LHC for use in p+p and p+Pb collisions. Recently, experiments at RHIC have explored the possibility of adopting these techniques for small collision systems at lower energies. In this talk, we systematically test these techniques using the Monte Carlo generators PYTHIA and HIJING, which do not include any collective flow, and AMPT, which does. We find that it is crucial to examine the results of such tests at the LHC and RHIC both as a function of multiplicity and particle  $p_T$ . Our results indicate reasonable non-flow subtraction for p+p collisions at the highest LHC energies, while failing if applied to p+p collisions at RHIC. In the case of p+Au collisions at RHIC, both HIJING and AMPT results indicate a substantial over-subtraction of non-flow for  $p_T \geq 1$  GeV/c and hence an underestimate of elliptic flow.

**Primary authors:** Dr LIM, Sanghoon (University of Colorado Boulder); Dr HU, Qipeng (University of Colorado Boulder); BELMONT, Ron (University of North Carolina Greensboro); HILL, Kurt (University of Colorado); NAGLE, James (University of Colorado); PEREPELITSA, Dennis (University of Colorado Boulder)

**Presenter:** NAGLE, James (University of Colorado)

**Session Classification:** Posters

**Track Classification:** Collectivity in small systems

Contribution ID: 115

Type: **Oral**

## Observation of collectivity in p+Au, d+Au and $^3\text{He}$ +Au collisions with PHENIX

*Wednesday, 26 June 2019 16:50 (20 minutes)*

In order to investigate the origin of collectivity in small systems, the PHENIX experiment has collected data of p+Au, d+Au and  $^3\text{He}$ +Au collisions at 200 GeV, giving a unique set of initial geometries. In this talk we present a complete set of elliptic and triangular flow measurements, which taken together provide unprecedented model discrimination between initial-state momentum correlation effects and final-state effects. This talk also covers the elliptic anisotropies measured in the d+Au beam energy scan, with collision energies ranging from 200 GeV to 19.6 GeV. The d+Au beam energy scan data leaves the intrinsic geometry unchanged but varies the system lifetime and the duration of the possible QGP phase. Measurements of  $v_2(p_T)$  and  $v_2(\eta)$  at different energies provide important insights about the system dynamics and allow further input for model comparison.

**Primary author:** Dr XU, Qiao (Vanderbilt University)**Presenter:** Dr XU, Qiao (Vanderbilt University)**Session Classification:** Parallel: Collectivity in small systems 2**Track Classification:** Collectivity in small systems

Contribution ID: 116

Type: **Poster**

## Investigation of the linear and mode-coupled flow harmonics in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV

*Tuesday, 25 June 2019 16:02 (1 minute)*

The elliptic and triangular flow coefficients,  $v_2$  and  $v_3$ , are expected to be driven by the initial anisotropy coefficients of the same order,  $\epsilon_2$  and  $\epsilon_3$ , respectively. However, the higher order flow coefficients,  $v_n$  ( $n > 3$ ), are comprised of linear contributions driven by  $\epsilon_n$ , as well as mode-coupled contributions derived from the lower order coefficients. The study of these disparate contributions to  $v_n$  can give important insight to discern initial-state models and to constrain the temperature-dependent specific shear viscosity,  $\frac{\eta}{s}(T)$ . In recent work, we have made detailed measurements of both the linear and the mode-coupled coefficients,  $v_n$  ( $n=4,5$ ), in Au+Au collisions ( $\sqrt{s_{NN}}=200$  GeV) using 2- and multi-particle correlations based on the standard and subevent cumulant methods. These measurements will be presented as a function of centrality,  $p_T$  and particle species. The comparisons to the LHC measurements and different theoretical calculations will be presented. The implications of these comparisons for initial-state models and  $\frac{\eta}{s}(T)$  will be discussed.

**Primary author:** Dr ABDELRAHMAN, Niseem (Department of Physics, University of Illinois at Chicago, Chicago)

**Presenter:** Dr ABDELRAHMAN, Niseem (Department of Physics, University of Illinois at Chicago, Chicago)

**Session Classification:** Posters

**Track Classification:** Initial conditions for hydrodynamics & transport coefficients



Contribution ID: 117

Type: **Poster**

## Viscous Hydrodynamics with Finite Baryon Number, Strangeness, and Electric Charge

*Tuesday, 25 June 2019 16:03 (1 minute)*

Relativistic viscous hydrodynamics has been an essential tool in studying the evolution of the Quark Gluon Plasma (QGP) produced in heavy-ion collisions as well as in searching for the critical point expected to be present for more baryon dense systems. With the coming runs of the Beam Energy Scan II at the Relativistic Heavy Ion Collider (RHIC), it will be necessary to implement conserved Baryon number,  $B$ , electric charge,  $Q$ , and strangeness,  $S$ , into the hydrodynamic description in order to have relevant theoretical predictions. This requires knowledge of an equation of state that contains thermodynamic information on the associated chemical potentials for  $B$ ,  $Q$ , and  $S$ . In this work, we use the most up-to-date equation of state which includes all of the necessary thermodynamic information to make relevant predictions about the effect of transport coefficients on system dynamics. In particular, we will explore how a large bulk viscosity near the critical point could lead to effects such as the possibility of a cavitating system, as well as the slowing down of dynamics leading to a longer lifetime of the fluid.

**Primary authors:** DORE, Travis (Rutgers University); NORONHA-HOSTLER, Jacquelyn (University of Houston); SIEVERT, Matthew (Rutgers University)

**Presenter:** DORE, Travis (Rutgers University)

**Session Classification:** Posters

**Track Classification:** Initial conditions for hydrodynamics & transport coefficients

Contribution ID: 120

Type: Oral

## Correlations between mid-rapidity charged tracks and large-rapidity event activity in p+Au collisions at $\sqrt{s_{NN}}=200$ GeV

*Wednesday, 26 June 2019 17:10 (20 minutes)*

Measurements of “small,” p+A or d+A, systems at the LHC and RHIC have suggested possible collective flow and, for high event activity collisions, jet modification that may scale with Bjorken  $x$ . They also provide input to related questions of the initial state of the proton prior to and throughout its collision with the opposing heavy ion nucleus. This talk presents preliminary measurements of correlations between mid-rapidity charged tracks and high-rapidity event activity measured by scintillator tiles in  $\sqrt{s_{NN}} = 200$  GeV p+Au collisions at STAR. These correlations are important because they inform the current discussion regarding the use of the Glauber model in small systems and have implications for calculating nuclear modification and quenching observables in these systems. The results support concerns about centrality binning in p+Au collisions, and as such motivate using ratios of semi-inclusive, as opposed to fully inclusive, jet spectra to look for jet enhancement or suppression.

**Primary author:** STEWART, David (Yale University)**Presenter:** STEWART, David (Yale University)**Session Classification:** Parallel: Collectivity in small systems 2**Track Classification:** Collectivity in small systems

Contribution ID: 121

Type: **Poster**

## NLO+NLLx computation of inclusive photon+dijet production in e+A DIS as a probe of gluon saturation

*Tuesday, 25 June 2019 16:20 (1 minute)*

We present the first computation of the NLO photon+dijet impact factor in e+A DIS at small  $x$ . When combined with the recent derivation of JIMWLK small  $x$  evolution to NLL $x$  accuracy, this result provides us with a prediction of the photon+dijet cross-section in e+A DIS to  $O(\alpha_s^3 \ln(1/x))$  accuracy. The comparison of this result with photon+dijet measurements at a future EIC therefore provides a precision test of the systematics of gluon saturation. In the soft photon limit, one obtains a compact representation of the state-of-the art results for fully inclusive DIS. The novel techniques developed in this computation can also be applied to promote existing LO computations of photon+dijet production in p+A collisions to NLO+NLL $x$  accuracy.

**Primary author:** ROY, Kaushik (Stony Brook University and Brookhaven National Laboratory)

**Co-author:** VENUGOPALAN, Raju (BNL)

**Presenter:** ROY, Kaushik (Stony Brook University and Brookhaven National Laboratory)

**Session Classification:** Posters

**Track Classification:** Forward and saturation physics

Contribution ID: 126

Type: Oral

## Off-equilibrium infrared structure of self-interacting scalar fields: Universal scaling, Vortex-antivortex superfluid dynamics and Bose-Einstein condensation

*Tuesday, 25 June 2019 15:00 (20 minutes)*

We map the infrared dynamics of a relativistic single component ( $N=1$ ) interacting scalar field theory to that of nonrelativistic complex scalar fields. The Gross-Pitaevskii (GP) equation, describing the real time dynamics of single component ultracold Bose gases, is obtained at first nontrivial order in an expansion proportional to the powers of  $\lambda\phi^2/m^2$  where  $\lambda$ ,  $\phi$  and  $m$  are the coupling constant, the scalar field and the particle mass respectively. Our analytical studies are corroborated by numerical simulations of the spatial and momentum structure of overoccupied scalar fields in (2+1)-dimensions. Universal scaling of infrared modes, vortex-antivortex superfluid dynamics and the off-equilibrium formation of a Bose-Einstein condensate are observed. Our results for the universal scaling exponents are in agreement with those extracted in the numerical simulations of the GP equation. As in these simulations, we observe coarsening phase kinetics in the Bose superfluid with strongly anomalous scaling exponents relative to that of vertex resummed kinetic theory. Our relativistic field theory framework further allows one to study more closely the coupling between superfluid and normal fluid modes, specifically the turbulent momentum and spatial structure of the coupling between a quasi-particle cascade to the infrared and an energy cascade to the ultraviolet. We outline possible applications of the formalism to the dynamics of vortex-antivortex formation and to the off-equilibrium dynamics of the strongly interacting matter formed in heavy-ion collisions.

**Primary authors:** VENUGOPALAN, Raju (BNL); SCHLICHTING, Soeren (University of Washington); WANG, Qun (University of science and technology of China)

**Presenter:** WANG, Qun (University of science and technology of China)

**Session Classification:** Parallel: Approach to Equilibrium

**Track Classification:** Approach to equilibrium (weak & strong coupling)

Contribution ID: 127

Type: **Oral**

## Factorization breaking - flow angle and magnitude decorrelation

*Tuesday, 25 June 2019 17:20 (20 minutes)*

The factorization breaking of collective flow in transverse momentum or in pseudorapidity is equivalent to a small decorrelation of flow in different phasespace regions. ATLAS has measured, besides the usual factorization breaking (including both flow magnitude and angle decorrelation), the flow angle decorrelation in pseudorapidity. The latter accounts for about half of the total decorrelation. These observations are confirmed in hydrodynamic calculations for the flow decorrelation, both in pseudorapidity [1] and in transverse momentum [2]. The general relation between flow angle and flow magnitude decorrelation can be understood within a simple model with a random component of the flow vector [3].

Our work makes also predictions for additional observables, that could be tested experimentally.

[1] P. Bozek, W. Broniowski, Phys. Rev. C 97 (2018) 034913.

[2] P. Bozek, Phys. Rev. C98 (2018) 064906.

[3] P. Bozek, H. Mehrabpour, in preparation

**Primary author:** BOZEK, Piotr (AGH University of Science and Technology)

**Presenter:** BOZEK, Piotr (AGH University of Science and Technology)

**Session Classification:** Parallel: Initial conditions for hydrodynamics & transport coefficients

**Track Classification:** Initial conditions for hydrodynamics & transport coefficients

Contribution ID: 129

Type: **Poster**

## Emergence of collectivity near a critical point

*Tuesday, 25 June 2019 16:15 (1 minute)*

Ever since the discovery of the quark-gluon plasma (QGP) the location of the critical point in the QCD phase diagram - the end point of the supposed first-order transition between hadronic matter and the QGP - has been a principal research goal for heavy-ion collision experiments at RHIC. We use the gauge/gravity duality to study a four-dimensional, strongly-coupled gauge theory with a first-order, second-order and crossover thermal phase transition. In the dual gauge theory we calculate the formation, evolution and saturation of the spinodal instability. We uncover a new surprising example of the applicability of hydrodynamics to systems with large gradients (JHEP 1706 (2017) 129 + upcoming work).

We discover with out-of-equilibrium shockwave collisions that in theories near a critical point a long-lived, quasi-static state may be formed. moreover, we show the Mueller-Israel-Stewart-type formulation of hydrodynamics to fail to describe pressures (Phys.Rev.Lett. 121 (2018), no.26, 261601) near a critical point. We provide the necessary correction terms and demonstrate that large second-order spatial derivatives need to be accounted for.

**Primary authors:** ATTEMS, Maximilian (Instituto Galego de Física de Altas Enerxías); Dr BEA, Yago (University of Barcelona); Prof. CASALDERREY SOLANA, Jorge (University of Barcelona); Prof. MATEOS, David (University of Barcelona); Dr ZILHAO, Miguel (University of Lisbon)

**Presenter:** ATTEMS, Maximilian (Instituto Galego de Física de Altas Enerxías)

**Session Classification:** Posters

**Track Classification:** Approach to equilibrium (weak & strong coupling)

Contribution ID: 131

Type: **Oral**

## Probing initial stages with scale dependent observables of the QGP in sPHENIX

*Wednesday, 26 June 2019 16:30 (20 minutes)*

Our understanding of QCD under extreme conditions has advanced tremendously following the discovery of the Quark Gluon Plasma and its detailed characterisation in heavy ion collisions at RHIC and the LHC. The sPHENIX experiment at RHIC will provide precision measurements of jet, upilon and open heavy flavor probes, complementing analogous measurements at the LHC. The physics program enabled by these measurements will advance understanding of QCD dynamics through all phases of the collision, connecting the initial stage in which one expects large temperature and parton density differences between collisions at RHIC and LHC energies, to subsequent stages in which the properties of scale sensitive probes can be further modified. We will describe the current status of the sPHENIX detector and its physics program, with an emphasis on the physics program enabled by the experiment's large coverage, high rate capability and precision vertexing.

**Primary authors:** MORRISON, David (BNL); ROLAND, Gunther (MIT)

**Presenter:** REED, Rosi (Lehigh University)

**Session Classification:** Parallel: Future facilities

**Track Classification:** Future facilities

Contribution ID: 132

Type: **Poster**

## Cold QCD Physics with sPHENIX and Potential Forward Upgrades

*Tuesday, 25 June 2019 16:12 (1 minute)*

The sPHENIX detector at RHIC, together with the accelerator's unique capabilities, will enable a spectrum of new or improved measurements enhancing our understanding of the initial state for nuclear collisions. Specifically, sPHENIX measurements in spin polarized proton-proton and proton-nucleus collisions will reveal more about how partons behave in a nuclear environment, explore spin-spin and spin-momentum correlations in the nucleon, and provide data to investigate effects of non-universality. A potential upgrade to sPHENIX with forward instrumentation could significantly enhance these physics capabilities. The cold QCD nuclear physics program for the proposed sPHENIX midrapidity detector as well as the enhanced program enabled with forward upgrades will be presented.

**Primary authors:** MORRISON, David (BNL); ROLAND, Gunther (MIT)

**Presenter:** JI, Zhongling (Stony Brook University)

**Session Classification:** Posters

**Track Classification:** nPDF, cold matter effects



Contribution ID: **133**

Type: **not specified**

## IAC Dinner

*Wednesday, 26 June 2019 19:00 (2 hours)*

Hosted by the Center for Frontiers in Nuclear Science at Miss Mamie's Spoonbread Too, 366 W 110th St, New York, NY 10025 (10 minute walk from Columbia)

Contribution ID: **134**

Type: **not specified**

## Poster session 2

Contribution ID: 135

Type: **not specified**

## Theory overview

*Monday, 24 June 2019 08:45 (30 minutes)*

**Presenter:** SCHLICHTING, Soeren (University of Washington)

**Session Classification:** Initial stages: current status

Contribution ID: **136**

Type: **not specified**

## **RHIC/LHC overview**

*Monday, 24 June 2019 09:15 (30 minutes)*

**Presenter:** MOHAPATRA, Soumya (Columbia University)

**Session Classification:** Initial stages: current status

Contribution ID: **137**

Type: **not specified**

## **Low-x frontier**

*Monday, 24 June 2019 09:45 (30 minutes)*

**Presenter:** WING, Matthew (UCL)

**Session Classification:** Initial stages: current status

Contribution ID: **138**

Type: **not specified**

## **CMS overview**

**Presenter:** WANG, Jing

**Session Classification:** Initial stages: current status

Contribution ID: **139**

Type: **not specified**

## STAR overview

**Presenter:** HUANG, shengli (Stony Brook University)

**Session Classification:** Initial stages: current status

Contribution ID: **140**

Type: **not specified**

## **LHCb overview**

**Presenter:** Dr SCHMIDT, Burkhard (CERN)

**Session Classification:** Initial stages: current status



Contribution ID: **141**

Type: **not specified**

## **ALICE overview**

**Presenter:** CAFFARRI, Davide (Nikhef)

**Session Classification:** Initial stages: current status

Contribution ID: **142**

Type: **not specified**

## **ATLAS Overview**

*Monday, 24 June 2019 14:00 (30 minutes)*

**Presenter:** JIANGYONG, Jia (stony brook university & BNL)

**Session Classification:** Experimental overviews 2

Contribution ID: **143**

Type: **not specified**

## **PHENIX overview**

*Monday, 24 June 2019 14:30 (30 minutes)*

**Presenters:** Dr LIM, Sanghoon (University of Colorado Boulder); LIM, Sanghoon (University of Colorado Boulder)

**Session Classification:** Experimental overviews 2

Contribution ID: 144

Type: **not specified**

## **Panel discussion: reactions to new data**

*Monday, 24 June 2019 15:00 (30 minutes)*

**Presenter:** EVDOKIMOV, Olga (UIC)

**Session Classification:** Experimental overviews 2

Contribution ID: 145

Type: **not specified**

## PDF+small x resummation

*Monday, 24 June 2019 16:00 (30 minutes)*

**Presenter:** BONVINI, Marco (INFN Sezione di Roma 1)

**Session Classification:** PDFs and saturation

Contribution ID: **146**

Type: **not specified**

## **nPDF overview**

*Monday, 24 June 2019 16:30 (30 minutes)*

**Presenter:** ZURITA, Maria (University of Regensburg)

**Session Classification:** PDFs and saturation

Contribution ID: **147**

Type: **not specified**

## PDF from lattice

*Monday, 24 June 2019 17:00 (30 minutes)*

**Presenter:** SHANAHAN, Phiala (MIT)

**Session Classification:** PDFs and saturation

Contribution ID: **148**

Type: **not specified**

## Saturation overview

*Monday, 24 June 2019 17:30 (30 minutes)*

**Presenter:** Dr ALTINOLUK, Tolga (NCBJ Warsaw)

**Session Classification:** PDFs and saturation



Contribution ID: 149

Type: **not specified**

## Deep Inelastic Scattering in the Valence Regime

*Tuesday, 25 June 2019 08:45 (30 minutes)*

**Presenter:** KEPPEL, Cynthia (Thomas Jefferson National Accelerator Facility)

**Session Classification:** Imaging the nucleon

Contribution ID: **150**

Type: **not specified**

## Nucleon imaging

*Tuesday, 25 June 2019 09:15 (30 minutes)*

**Presenter:** BOUSSARIE, Renaud (Brookhaven National Lab)

**Session Classification:** Imaging the nucleon

Contribution ID: **151**

Type: **not specified**

## **GPDs/Wigner functions**

*Tuesday, 25 June 2019 09:45 (30 minutes)*

**Presenter:** DUMITRU, Adrian (Dept. of Natural Sciences, Baruch College (CUNY))

**Session Classification:** Imaging the nucleon

Contribution ID: 152

Type: **not specified**

## Thermalization across energy scales

*Tuesday, 25 June 2019 10:45 (30 minutes)*

**Presenter:** Prof. BERGES, Jürgen (Heidelberg University)

**Session Classification:** Approach to hydro

Contribution ID: **153**

Type: **not specified**

## Hydrodynamic matching

*Tuesday, 25 June 2019 11:15 (30 minutes)*

**Presenter:** MAZELIAUSKAS, Aleksas (University of Heidelberg)

**Session Classification:** Approach to hydro

Contribution ID: 154

Type: **not specified**

## Hydro attractors

*Tuesday, 25 June 2019 11:45 (30 minutes)*

**Presenter:** HELLER, Michal P. (Max Planck Institute for Gravitational Physics (Albert Einstein Institute) / National Centre for Nuclear Research)

**Session Classification:** Approach to hydro

Contribution ID: 155

Type: **not specified**

## **Panel discussion: approach to hydro**

*Tuesday, 25 June 2019 12:15 (30 minutes)*

**Presenter:** Dr WIEDEMANN, Urs (CERN)

**Session Classification:** Approach to hydro

Contribution ID: **156**

Type: **not specified**

## **Flow: small & large systems**

*Wednesday, 26 June 2019 08:45 (30 minutes)*

**Presenter:** KRIZKOVA GAJDOSOVA, Katarina (Czech Technical University in Prague)

**Session Classification:** Initial vs. final state correlations



Contribution ID: **157**

Type: **not specified**

## **Initial state correlations**

*Wednesday, 26 June 2019 09:15 (30 minutes)*

**Presenters:** MACE, Mark (Stony Brook University); MACE, Mark (University of Jyväskylä)

**Session Classification:** Initial vs. final state correlations

Contribution ID: **158**

Type: **not specified**

## **Hydro in small systems: How small can a QGP be?**

*Wednesday, 26 June 2019 09:45 (30 minutes)*

**Presenter:** BRONIOWSKI, Wojciech (IFJ PAN & UJK)

**Session Classification:** Initial vs. final state correlations

Contribution ID: 159

Type: **not specified**

## EMC from short range correlations

*Wednesday, 26 June 2019 10:45 (30 minutes)*

**Presenter:** FOMIN, Nadia (University of Tennessee)

**Session Classification:** Aspects of the initial state

Contribution ID: **160**

Type: **not specified**

## Proton shape fluctuations

*Wednesday, 26 June 2019 11:15 (30 minutes)*

**Presenter:** Dr MÄNTYSAARI, Heikki (University of Jyväskylä)

**Session Classification:** Aspects of the initial state

Contribution ID: **161**

Type: **not specified**

## **Hydro perspectives on BES / Longitudinal dynamics**

*Wednesday, 26 June 2019 11:45 (30 minutes)*

**Presenter:** SHEN, Chun (Wayne State University)

**Session Classification:** Aspects of the initial state

Contribution ID: **162**

Type: **not specified**

## **Panel: Input from Rice/INT workshops & discussion**

*Wednesday, 26 June 2019 12:15 (30 minutes)*

**Presenter:** LI, Wei (Rice University)

**Session Classification:** Aspects of the initial state

Contribution ID: **163**

Type: **not specified**

## Chiral magnetohydrodynamics

*Thursday, 27 June 2019 08:45 (30 minutes)*

**Presenter:** LIAO, JINFENG (INDIANA UNIVERSITY)

**Session Classification:** Electromagnetic fields, chirality and vorticity

Contribution ID: **164**

Type: **not specified**

## **Hydrodynamics with spin / Connection to vorticity**

*Thursday, 27 June 2019 09:15 (30 minutes)*

**Presenter:** RYBLEWSKI, Radoslaw (Institute of Nuclear Physics Polish Academy of Sciences)

**Session Classification:** Electromagnetic fields, chirality and vorticity



Contribution ID: **165**

Type: **not specified**

## **Evidence for EM fields**

*Thursday, 27 June 2019 09:45 (30 minutes)*

**Presenter:** DUBLA, Andrea

**Session Classification:** Electromagnetic fields, chirality and vorticity

Contribution ID: **166**

Type: **not specified**

## Entanglement at collider energies

*Thursday, 27 June 2019 10:45 (30 minutes)*

**Presenter:** FLOERCHINGER, Stefan (Heidelberg University, Institute for Theoretical Physics)

**Session Classification:** Strong vs. weak coupling

Contribution ID: **167**

Type: **not specified**

## **Strong & weak coupling approaches**

*Thursday, 27 June 2019 11:15 (30 minutes)*

**Presenter:** Dr GROZDANOV, Sašo (MIT)

**Session Classification:** Strong vs. weak coupling

Contribution ID: **168**

Type: **not specified**

## Transport coefficients

*Thursday, 27 June 2019 11:45 (30 minutes)*

**Presenter:** WU, Bin (The Ohio State University)

**Session Classification:** Strong vs. weak coupling

Contribution ID: **169**

Type: **not specified**

## **Panel: strong vs. weak coupling approaches**

*Thursday, 27 June 2019 12:15 (30 minutes)*

**Presenter:** Prof. ARNOLD, Peter (University of Virginia)

**Session Classification:** Strong vs. weak coupling

Contribution ID: 170

Type: **not specified**

## Initial conditions from jets: pA & AA

*Thursday, 27 June 2019 14:00 (30 minutes)*

**Presenter:** ANDRES, Carlota (Jefferson Lab)

**Session Classification:** High pT probes of the initial state

Contribution ID: 171

Type: **not specified**

## **LHC forward physics from hard probes**

*Thursday, 27 June 2019 14:30 (30 minutes)*

**Presenter:** RYBAR, Martin (Columbia University (US))

**Session Classification:** High  $p_T$  probes of the initial state

Contribution ID: 172

Type: **not specified**

## Initial stages using EM probes

*Thursday, 27 June 2019 15:00 (30 minutes)*

**Presenter:** YANG, Shuai (BNL)

**Session Classification:** High  $p_T$  probes of the initial state



Contribution ID: 173

Type: **not specified**

## Heavy flavor in small systems

*Thursday, 27 June 2019 16:00 (30 minutes)*

**Presenter:** CHEN, Zhenyu (Stony Brook University)

**Session Classification:** Heavy flavor, e+e- & UPC

Contribution ID: 174

Type: **not specified**

## Ridge in $e+e^-/\text{DIS}$

*Thursday, 27 June 2019 16:30 (30 minutes)*

**Presenter:** BATY, Austin

**Session Classification:** Heavy flavor,  $e+e^-$  & UPC

Contribution ID: 175

Type: **not specified**

## UPC: theory & experiment

*Thursday, 27 June 2019 17:00 (30 minutes)*

**Presenter:** SEGER, Janet (Creighton University)

**Session Classification:** Heavy flavor, e+e- & UPC

Contribution ID: 176

Type: Oral

## $v_n$ - $p_T$ correlations in 5.02 TeV Pb+Pb and p+Pb collisions with the ATLAS detector

*Friday, 28 June 2019 09:00 (15 minutes)*

Recently collected data by the ATLAS experiment at the LHC offer opportunities to explore dynamic properties of quark-gluon plasma. A new tool to study these properties is the modified Pearson's correlation coefficient,  $\rho$ , that quantifies the correlation between the mean transverse momentum in an event,  $[p_T]$ , and the square of the flow harmonic magnitude,  $v_n^2$ . To suppress non-flow effects,  $v_n^2$  is calculated by correlating charged particles from two sub-events covering opposite pseudorapidity ranges of  $0.75 < |\eta| < 2.5$  while  $[p_T]$  is evaluated for particles with  $|\eta| < 0.5$ . The measurement of  $\rho$  is performed using minimum-bias p+Pb and Pb+Pb collisions at the same energy  $\sqrt{s_{NN}} = 5.02$ -TeV allowing for a comparison of the medium dynamics in small and large systems. In Pb+Pb collisions, values of  $\rho$  coefficients are found to significantly deviate from zero for studied harmonics ( $v_2$ ,  $v_3$ , and  $v_4$ ). The coefficients as a function of centrality are observed to be weakly dependent on the transverse momentum range of the selected particles, despite large differences in the mean transverse momentum in an event and the magnitude of fluctuations of flow harmonics. The  $\rho$  coefficient in Pb+Pb collisions for the second order harmonics has a positive value for mid-central collisions and decreases in the most central events. In p+Pb collisions the  $\rho$  coefficient is measured only for the second order flow harmonics. In both p+Pb and peripheral Pb+Pb collisions, it is found to be negative. All measured coefficients are compared to theoretical models.

**Presenter:** BURKA, Klaudia (AGH-UST)**Session Classification:** Flash talks & toward the EIC

Contribution ID: 177

Type: **Oral**

## Linear and non-linear response of two-particle correlations to initial-geometry fluctuations

*Friday, 28 June 2019 09:15 (15 minutes)*

We investigate the importance of different features of the initial geometry to anisotropic flow fluctuations in heavy-ion collisions. To that end, we explore the hydrodynamic response of differential flow harmonics  $v_n(pT)$  to generalized eccentricities  $e_{n,m}$  of the initial density profile within a realistic hydrodynamic model. Special attention is paid to two-particle angular correlations, characterized in detail by the principal-component analysis (PCA). We address the relevance of non-linear response, as well as the stability of the results against the inclusion of extra eccentricities. Additionally, we study new effects from multiplicity fluctuations, which could lead to redundancies in the experimental PCA data.

**Presenter:** HIPPERT, Mauricio (Universidade de Sao Paulo)

**Session Classification:** Flash talks & toward the EIC

Contribution ID: **178**

Type: **not specified**

## Spin at low-x

*Friday, 28 June 2019 09:30 (30 minutes)*

**Presenter:** SIEVERT, Matthew (Rutgers University)

**Session Classification:** Flash talks & toward the EIC

Contribution ID: **179**

Type: **not specified**

## **pA vs. eA universality**

*Friday, 28 June 2019 10:00 (30 minutes)*

**Presenter:** XIAO, Bowen (Central China Normal University)

**Session Classification:** Flash talks & toward the EIC

Contribution ID: **180**

Type: **not specified**

## **RHIC/LHC forward physics & upgrades**

*Friday, 28 June 2019 11:00 (30 minutes)*

**Presenter:** LAJOIE, John (Iowa State University)

**Session Classification:** The future of initial stages (chair TBA)



Contribution ID: **181**

Type: **not specified**

## **EIC perspective**

*Friday, 28 June 2019 11:30 (30 minutes)*

**Presenter:** SICHTERMANN, Ernst (Lawrence Berkeley National Laboratory)

**Session Classification:** The future of initial stages (chair TBA)

Contribution ID: **182**

Type: **not specified**

## **CERN perspective**

*Friday, 28 June 2019 12:00 (30 minutes)*

**Presenter:** ARMESTO, Nestor (Universidade de Santiago de Compostela)

**Session Classification:** The future of initial stages (chair TBA)

Contribution ID: **183**

Type: **Oral**

## Conference summary

*Friday, 28 June 2019 12:30 (30 minutes)*

**Presenter:** NORONHA-HOSTLER, Jacquelyn (University of Houston)

**Session Classification:** The future of initial stages (chair TBA)

Contribution ID: **184**

Type: **not specified**

## **Measurement of electroweak-boson production in p–Pb and Pb–Pb collisions at the LHC with ALICE**

**Presenter:** Dr VALLE, Nicolo (Universita and INFN, Pavia (IT))

**Session Classification:** Parallel: nPDF/CNM

Contribution ID: **185**

Type: **not specified**

## **Heavy electroweak boson production in Pb+Pb collisions with ATLAS**

**Presenter:** DUMANCIC, Mirta (Weizmann Institute of Science)

**Session Classification:** Parallel: nPDF/CNM

Contribution ID: **186**

Type: **not specified**

## **Constraining nPDFs with Z boson and Drell-Yan measurements in pPb collisions with CMS**

**Presenter:** CHAPON, Émilien (CERN)

**Session Classification:** Parallel: nPDF/CNM

Contribution ID: **187**

Type: **not specified**

## **Quarkonium production in pp and p–Pb collisions with ALICE at the LHC**

**Presenter:** HAYASHI, Shinichi (CNS, the University of Tokyo)

**Session Classification:** Parallel: nPDF/CNM

Contribution ID: **188**

Type: **not specified**

## **Measurement of open heavy-flavour hadron production in pp, p-Pb and Pb-Pb collisions with ALICE**

**Presenter:** FAGGIN, Mattia

**Session Classification:** Parallel: nPDF/CNM



Contribution ID: **189**

Type: **not specified**

## **PHENIX measurements of muon pairs from $c\bar{c}$ , $b\bar{b}$ , and Drell-Yan in p+p and p+Au at 200 GeV**

**Presenter:** DREES, Axel (Stony Brook University)

**Session Classification:** Parallel: nPDF/CNM

Contribution ID: **190**

Type: **not specified**

## **Non-linear evolution in QCD at high-energy beyond leading order**

**Presenter:** DUCLOUE, Bertrand (IPhT Saclay)

**Session Classification:** Parallel: Forward/saturation/spin

Contribution ID: 191

Type: **not specified**

## **Measurements of nuclear parton distribution functions using dijets, forward jets, and photo-nuclear jets at the CMS detector & prospects for measurements in Run III**

**Presenter:** BYLINKIN, Alexander (The University of Kansas (US))

**Session Classification:** Parallel: Forward/saturation/spin

Contribution ID: 192

Type: **not specified**

## **First measurement of Diffraction in pPb collisions and recent results on ultra-peripheral heavy-ion processes**

**Presenter:** WALCZAK, Marek (University of Warsaw)

**Session Classification:** Parallel: Forward/saturation/spin

Contribution ID: **193**

Type: **not specified**

## **EPR paradox and quantum entanglement at sub-nucleonic scales**

**Presenter:** TU, Zhoudunming (BNL)

**Session Classification:** Parallel: Forward/saturation/spin

Contribution ID: **194**

Type: **not specified**

## **Magnetic field in expanding quark-gluon plasma**

**Presenter:** TUCHIN, Kirill

**Session Classification:** Parallel: Forward/saturation/spin

Contribution ID: 195

Type: **not specified**

## **Relaxation dynamics of chiral transports and spin polarization in Quark-Gluon Plasma**

**Presenter:** LI, Shiyong (UIC)

**Session Classification:** Parallel: Forward/saturation/spin

Contribution ID: 196

Type: **not specified**

## **Connecting far-from-equilibrium hydrodynamics, resummed transport coefficients, and attractor solutions**

**Presenter:** NORONHA, Jorge (Universidade de Sao Paulo)

**Session Classification:** Parallel: Approach to Equilibrium



Contribution ID: **197**

Type: **not specified**

## **Stochastic hydrodynamics and long time tails of a non-equilibrium fluid**

**Presenter:** Dr MARTINEZ , Mauricio (North Carolina State University)

**Session Classification:** Parallel: Approach to Equilibrium

Contribution ID: **198**

Type: **not specified**

## **Self-similarity and spectral functions of non-Abelian plasmas in 2+1D**

**Presenter:** Dr PEURON, Jarkko (European Centre for Theoretical Studies in Nuclear Physics and Related Areas (ECT\*))

**Session Classification:** Parallel: Approach to Equilibrium

Contribution ID: 199

Type: **not specified**

## **Off-equilibrium infrared structure of self-interacting scalar fields: Universal scaling, Vortex-antivortex superfluid dynamics and Bose-Einstein condensation**

**Presenter:** WANG, Qun (University of science and technology of China)

**Session Classification:** Parallel: Approach to Equilibrium

Contribution ID: **200**

Type: **not specified**

## **Holographic collisions with baryon number at intermediate coupling**

**Presenter:** VAN DER SCHEE, Wilke (Utrecht University)

**Session Classification:** Parallel: Approach to Equilibrium

Contribution ID: **201**

Type: **not specified**

## **Anisotropic hydrodynamics with a realistic collisional kernel**

**Presenter:** ALMAALOL, Dekrayat (Kent State Uiniversity)

**Session Classification:** Parallel: Approach to Equilibrium

Contribution ID: 202

Type: **not specified**

## **Hydrodynamics far-from-equilibrium: a concrete example in kinetic theory**

**Presenter:** SILVEIRA DENICOL, Gabriel (Universidade Federal Fluminense)

**Session Classification:** Parallel: Initial conditions for hydrodynamics & transport coefficients

Contribution ID: 203

Type: **not specified**

## **Factorization breaking - flow angle and magnitude decorrelation**

**Presenter:** BOZEK, Piotr (AGH University of Science and Technology)

**Session Classification:** Parallel: Initial conditions for hydrodynamics & transport coefficients

Contribution ID: 204

Type: **not specified**

## **Non-Gaussian fluctuations of $v_1, v_2, v_3$ and $v_4$ and their correlations in Pb+Pb collisions with the ATLAS detector**

**Presenter:** BEHERA, Arabinda (STAR)

**Session Classification:** Parallel: Initial conditions for hydrodynamics & transport coefficients



Contribution ID: 205

Type: **not specified**

## **Longitudinal fluctuations and decorrelations of anisotropic flows in relativistic heavy-ion collisions**

**Presenter:** QIN, Guang-You (Central China Normal University)

**Session Classification:** Parallel: Initial conditions for hydrodynamics & transport coefficients

Contribution ID: 206

Type: **not specified**

## **Principal Component Analysis of collective flow in Heavy-Ion collisions**

**Presenter:** LIU, ZIMING (Peking University)

**Session Classification:** Parallel: Initial conditions for hydrodynamics & transport coefficients

Contribution ID: 207

Type: **not specified**

## **New paradigm for fluctuations in heavy-ion collisions**

**Presenter:** GIACALONE, Giuliano (IPhT - Saclay)

**Session Classification:** Parallel: Initial conditions for hydrodynamics & transport coefficients

Contribution ID: 208

Type: **not specified**

## **Investigation of the linear and mode-coupled flow harmonics in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV**

*Wednesday, 26 June 2019 14:40 (20 minutes)*

**Presenter:** ABDELRAHMAN, Niseem (Stony Brook University)

**Session Classification:** Parallel: Collectivity in small systems 1

Contribution ID: 209

Type: **not specified**

## **ATLAS measurements of azimuthal anisotropy of heavy flavor hadrons in Pb+Pb, p+Pb and pp collisions**

**Presenter:** HILL, Kurt (University of Colorado)

**Session Classification:** Parallel: Collectivity in small systems 1

Contribution ID: **210**

Type: **not specified**

## **Centrality dependence of collectivity in kinetic theory**

**Presenter:** WIEDEMANN, Urs

**Session Classification:** Parallel: Collectivity in small systems 1

Contribution ID: 211

Type: **not specified**

## **Recent ATLAS results on correlations in small collisions systems and photon-induced processes in ultra-peripheral Pb+Pb collisions at 5.02 TeV**

**Presenter:** SEIDLITZ, Blair (University of Colorado Boulder)

**Session Classification:** Parallel: Collectivity in small systems 1

Contribution ID: 212

Type: **not specified**

## **Investigation of collectivity in small collision systems with ALICE**

**Presenter:** PACIK, Vojtech (Niels Bohr Institute, University of Copenhagen, Denmark)

**Session Classification:** Parallel: Collectivity in small systems 1



Contribution ID: 213

Type: **not specified**

## **IP-Jazma critical assessment of physics attributions**

**Presenter:** Dr ZAJC, Bill (Columbia University)

**Session Classification:** Parallel: Collectivity in small systems 1

Contribution ID: 214

Type: **not specified**

## **Measurement of elliptic and triangular flow with multiparticle correlations in pPb collisions at 8.16 TeV**

**Presenter:** Dr TUO, Shengquan (Vanderbilt University (US))

**Session Classification:** Parallel: Collectivity in small systems 2

Contribution ID: 215

Type: **not specified**

## **Observation of collectivity in p+Au, d+Au and $^3\text{He}$ +Au collisions with PHENIX**

**Presenter:** XU, Qiao (Vanderbilt University)

**Session Classification:** Parallel: Collectivity in small systems 2

Contribution ID: 216

Type: **not specified**

## **Correlations between mid-rapidity charged tracks and large-rapidity event activity in p+Au collisions at $\sqrt{s_{NN}}=200$ GeV**

**Presenter:** STEWART, David (Yale University)

**Session Classification:** Parallel: Collectivity in small systems 2

Contribution ID: 217

Type: **not specified**

# **Hydrodynamic simulations of relativistic nuclear collisions with nucleon substructure: combined analysis of p+Pb and Pb+Pb collision systems at 5.02 TeV**

**Presenter:** BASS, Steffen (Duke)

**Session Classification:** Parallel: Collectivity in small systems 2

Contribution ID: **218**

Type: **not specified**

## **Initial state fluctuations in Pythia 8**

**Presenter:** RASMUSSEN, Christine (Lund University)

**Session Classification:** Parallel: Collectivity in small systems 2

Contribution ID: 219

Type: **not specified**

## Measurement of rapidity-odd directed flow for $D^0$ and $\overline{D}^0$ mesons using the STAR detector at RHIC

*Wednesday, 26 June 2019 18:10 (20 minutes)*

**Presenter:** SINGHA, Subhash (Kent State University)

**Session Classification:** Parallel: Collectivity in small systems 2

Contribution ID: 220

Type: **not specified**

## **Jet and photon probes of small and large systems in ATLAS**

**Presenter:** PEREPELITSA, Dennis (University of Colorado Boulder)

**Session Classification:** Parallel: High pT probes of the initial state



Contribution ID: 221

Type: **not specified**

## **Measurement of bottomonia in pp, pPb and PbPb collisions at 5.02 TeV with the CMS detector**

*Wednesday, 26 June 2019 14:20 (20 minutes)*

**Presenter:** KIM, Yongsun (UIUC)

**Session Classification:** Parallel: High pT probes of the initial state

Contribution ID: 222

Type: **not specified**

## **Probing collision dynamics of small system collisions via high $p_T$ hadrons and direct photons by the PHENIX experiment at RHIC**

**Presenter:** SAKAGUCHI, Takao (BNL)

**Session Classification:** Parallel: High  $p_T$  probes of the initial state

Contribution ID: 223

Type: **not specified**

## **How to infer the shape of the QGP droplet from the data**

**Presenter:** DJORDJEVIC, Magdalena (Institute of Physics Belgrade)

**Session Classification:** Parallel: High  $p_T$  probes of the initial state

Contribution ID: 224

Type: **not specified**

## **A complete set of splitting functions in nuclear matter to any order in opacity and applications to jet physics**

**Presenter:** VITEV, Ivan (LANL)

**Session Classification:** Parallel: High pT probes of the initial state

Contribution ID: 225

Type: **not specified**

## **Measurement of heavy-flavour jets and correlations and elliptic flow in small systems with ALICE**

**Presenter:** COLAMARIA, Fabio (INFN, Sezione di Bari (IT))

**Session Classification:** Parallel: High  $p_T$  probes of the initial state

Contribution ID: 226

Type: **not specified**

## **Partonic spatial imaging at an electron-ion collider**

**Presenter:** LEE, J.H. (Brookhaven National Laboratory)

**Session Classification:** Parallel: Future facilities

Contribution ID: 227

Type: **not specified**

## Computing the gluon Sivers function at low- $x$

**Presenter:** HATTA, Yoshitaka (BNL)

**Session Classification:** Parallel: Future facilities

Contribution ID: 228

Type: **not specified**

## A Forward Rapidity Upgrade for the STAR Detector

**Presenter:** BRANDENBURG, Daniel (Brookhaven National Laboratory)

**Session Classification:** Parallel: Future facilities



Contribution ID: **229**

Type: **not specified**

## **LHCb fixed target results and prospects**

**Presenter:** Dr DI NEZZA, Pasquale (INFN Frascati)

**Session Classification:** Parallel: Future facilities

Contribution ID: 230

Type: **not specified**

## **Semi-inclusive Deep-Inelastic Scattering, Parton Distributions and Fragmentation Functions at a Future Electron-Ion Collider**

**Presenter:** BORSA SANJUAN, Ignacio (Universidad de Buenos Aires)

**Session Classification:** Parallel: Future facilities

Contribution ID: 231

Type: **not specified**

## **Probing initial stages with scale dependent observables of the QGP in sPHENIX**

**Presenter:** REED, Rosi (Lehigh University)

**Session Classification:** Parallel: Future facilities

Contribution ID: 232

Type: **Poster**

## **Influence of neutron skin in heavy ion collisions at RHIC energies**

**Presenter:** HAMMELMANN, Jan

**Session Classification:** Posters

Contribution ID: 233

Type: **Oral**

## Welcome and conference logistics

**Presenters:** MUELLER, Berndt (BNL); STEINBERG, Peter (BNL); VENUGOPALAN, Raju (BNL)

**Session Classification:** Initial stages: current status

Contribution ID: 234

Type: **not specified**

## **Tribute to Roy Glauber (1925-2018)**

*Thursday, 27 June 2019 17:30 (30 minutes)*

**Presenters:** Dr ZAJC, Bill (Columbia University); ZAJC, William (Columbia University)

**Session Classification:** Heavy flavor, e+e- & UPC

Contribution ID: 235

Type: **not specified**

## Welcome from Berndt Mueller

*Monday, 24 June 2019 08:40 (5 minutes)*

**Presenter:** MUELLER, Berndt (BNL)

**Session Classification:** Introduction to IS2019

Contribution ID: **236**

Type: **not specified**

## Conference information

*Monday, 24 June 2019 08:30 (10 minutes)*

**Presenters:** STEINBERG, Peter (BNL); VENUGOPALAN, Raju (BNL)

**Session Classification:** Introduction to IS2019



Contribution ID: **237**

Type: **not specified**

## **CMS Overview**

*Monday, 24 June 2019 10:45 (30 minutes)*

**Presenter:** WANG, Jing

**Session Classification:** Experimental overviews 1

Contribution ID: 238

Type: **not specified**

## STAR overview

*Monday, 24 June 2019 11:15 (30 minutes)*

**Presenter:** HUANG, shengli (Stony Brook University)

**Session Classification:** Experimental overviews 1

Contribution ID: **239**

Type: **not specified**

## **LHCb overview**

*Monday, 24 June 2019 11:45 (30 minutes)*

**Presenter:** SCHMIDT, Burkhard (CERN)

**Session Classification:** Experimental overviews 1

Contribution ID: **240**

Type: **not specified**

## **ALICE overview**

*Monday, 24 June 2019 12:15 (30 minutes)*

**Presenter:** CAFFARRI, Davide (Nikhef)

**Session Classification:** Experimental overviews 1

Contribution ID: **241**

Type: **Poster**

## **Conformal invariance of TMD rapidity evolution**

I discuss conformal properties of TMD operators and present the result of the conformal rapidity evolution of TMD operators in the Sudakov region.

**Primary author:** VENUGOPALAN, Raju (BNL)

**Presenter:** VENUGOPALAN, Raju (BNL)

**Track Classification:** 3D nucleon structure

Contribution ID: 242

Type: **Poster**

## Conformal invariance of TMD rapidity evolution

*Tuesday, 25 June 2019 16:22 (1 minute)*

I discuss conformal properties of TMD operators and present the result of the conformal rapidity evolution of TMD operators in the Sudakov region

**Presenter:** VENUGOPALAN, Raju (BNL)

**Session Classification:** Posters

Contribution ID: **243**

Type: **not specified**

## Acknowledgements

*Friday, 28 June 2019 10:45 (15 minutes)*

**Presenters:** STEINBERG, Peter (BNL); VENUGOPALAN, Raju (BNL)