



UNIVERSITÄT
HEIDELBERG
ZUKUNFT
SEIT 1386

Long live the Higgs Portal!

Anastasia Filimonova


Based on 1812.04628 and current work

Brookhaven Forum 2019, 26/09

Image credit:
ATLAS collaboration

Higgs portal for fermionic dark matter

Higgs portal for fermionic dark matter




Pure electroweak singlet, doublet, triplet.

Dark matter candidate **only at Higgs threshold** or **too heavy for colliders**.

[Beniwal et al., 1512.06458]

[Cirelli et al., 1507.05519]

Higgs portal for fermionic dark matter




Pure electroweak singlet, doublet, triplet.


Dark matter candidate **only at Higgs threshold** or **too heavy for colliders**.

[Beniwal et al., 1512.06458]

[Cirelli et al., 1507.05519]



Minimal renormalizable extensions:
singlet-singlet, singlet-doublet, doublet-triplet.



Tiny portal coupling: **naturalness?**

[Freitas, Westhoff, Zupan, 1506.04149]

[Lee et al., 0803.2932] [Mahbubani, Senatore, hep-ph/0510064]

[Dedes, Karamitros, 1403.7744]

Singlet-triplet model for dark matter

2 majorana fields: $SU(2)$ singlet χ_S and triplet χ_T

$$\mathcal{L}_{\text{eff}} \supset -\frac{m_S}{2}\bar{\chi}_S\chi_S - \frac{m_T}{2}\text{Tr}[\bar{\chi}_T\chi_T] + \frac{\kappa_{ST}}{\Lambda} [(H^\dagger\bar{\chi}_TH)\chi_S + h.c.]$$

$$\text{with } \chi_S = \chi_S^0, \quad \chi_T = \begin{pmatrix} \chi_T^0/\sqrt{2} & \chi^+ \\ \chi^- & -\chi_T^0/\sqrt{2} \end{pmatrix}$$

Singlet-triplet model for dark matter

2 majorana fields: $SU(2)$ singlet χ_S and triplet χ_T **Naturally small**

$$\mathcal{L}_{\text{eff}} \supset -\frac{m_S}{2}\bar{\chi}_S\chi_S - \frac{m_T}{2}\text{Tr}[\bar{\chi}_T\chi_T] + \frac{\kappa_{ST}}{\Lambda} [(H^\dagger\bar{\chi}_TH)\chi_S + h.c.]$$

$$\text{with } \chi_S = \chi_S^0, \quad \chi_T = \begin{pmatrix} \chi_T^0/\sqrt{2} & \chi^+ \\ \chi^- & -\chi_T^0/\sqrt{2} \end{pmatrix}$$

Singlet-triplet model for dark matter

2 majorana fields: $SU(2)$ singlet χ_S and triplet χ_T **Naturally small**

$$\mathcal{L}_{\text{eff}} \supset -\frac{m_S}{2}\bar{\chi}_S\chi_S - \frac{m_T}{2}\text{Tr}[\bar{\chi}_T\chi_T] + \frac{\kappa_{ST}}{\Lambda} [(H^\dagger\bar{\chi}_TH)\chi_S + h.c.]$$

$$\text{with } \chi_S = \chi_S^0, \quad \chi_T = \begin{pmatrix} \chi_T^0/\sqrt{2} & \chi^+ \\ \chi^- & -\chi_T^0/\sqrt{2} \end{pmatrix}$$

- Three new parameters: m_S , m_T , $\mu = \frac{\kappa_{ST}v^2}{\sqrt{2}\Lambda}$
- Freedom in the field definitions: can choose $m_T, \kappa_{ST} > 0$

Singlet-triplet model for dark matter

2 majorana fields: $SU(2)$ singlet χ_S and triplet χ_T **Naturally small**

$$\mathcal{L}_{\text{eff}} \supset -\frac{m_S}{2} \bar{\chi}_S \chi_S - \frac{m_T}{2} \text{Tr}[\bar{\chi}_T \chi_T] + \frac{\kappa_{ST}}{\Lambda} [(H^\dagger \bar{\chi}_T H) \chi_S + h.c.]$$

$$\text{with } \chi_S = \chi_S^0, \quad \chi_T = \begin{pmatrix} \chi_T^0/\sqrt{2} & \chi^+ \\ \chi^- & -\chi_T^0/\sqrt{2} \end{pmatrix}$$

- Three new parameters: m_S , m_T , $\mu = \frac{\kappa_{ST} v^2}{\sqrt{2} \Lambda}$
- Freedom in the field definitions: can choose $m_T, \kappa_{ST} > 0$

Tree-level structure + electroweak corrections:

$$(\Delta m_{hc})^{\text{ew}} \left(\begin{array}{c} \frac{\chi_h}{\chi^+} \\ \frac{\chi_l}{\chi^+} \end{array} \right) \begin{pmatrix} (\Delta m_{hc})^{\text{mix}} \\ \Delta m_{cl} \end{pmatrix}$$

Singlet-triplet model for dark matter

2 majorana fields: $SU(2)$ singlet χ_S and triplet χ_T **Naturally small**

$$\mathcal{L}_{\text{eff}} \supset -\frac{m_S}{2} \bar{\chi}_S \chi_S - \frac{m_T}{2} \text{Tr}[\bar{\chi}_T \chi_T] + \frac{\kappa_{ST}}{\Lambda} [(H^\dagger \bar{\chi}_T H) \chi_S + h.c.]$$

$$\text{with } \chi_S = \chi_S^0, \quad \chi_T = \begin{pmatrix} \chi_T^0/\sqrt{2} & \chi^+ \\ \chi^- & -\chi_T^0/\sqrt{2} \end{pmatrix}$$

- Three new parameters: m_S , m_T , $\mu = \frac{\kappa_{ST} v^2}{\sqrt{2} \Lambda}$
- Freedom in the field definitions: can choose $m_T, \kappa_{ST} > 0$

Tree-level structure + electroweak corrections:

$$(\Delta m_{hc})^{\text{ew}} \left(\begin{array}{c} \underline{\chi_h} \\ \underline{\chi^+} \\ \dots\dots\dots \\ \underline{\chi_l} \end{array} \right) \begin{array}{l} (\Delta m_{hc})^{\text{mix}} \\ \\ \Delta m_{cl} \end{array}$$

Small μ :

$$\left(\begin{array}{c} \underline{\chi^+} \\ \underline{\chi_h} \\ \underline{\chi_l} \end{array} \right) \lesssim (\Delta m_{hc})^{\text{ew}}$$

Singlet-triplet model for dark matter

2 majorana fields: $SU(2)$ singlet χ_S and triplet χ_T **Naturally small**

$$\mathcal{L}_{\text{eff}} \supset -\frac{m_S}{2} \bar{\chi}_S \chi_S - \frac{m_T}{2} \text{Tr}[\bar{\chi}_T \chi_T] + \frac{\kappa_{ST}}{\Lambda} [(H^\dagger \bar{\chi}_T H) \chi_S + h.c.]$$

$$\text{with } \chi_S = \chi_S^0, \quad \chi_T = \begin{pmatrix} \chi_T^0/\sqrt{2} & \chi^+ \\ \chi^- & -\chi_T^0/\sqrt{2} \end{pmatrix}$$

- Three new parameters: m_S , m_T , $\mu = \frac{\kappa_{ST} v^2}{\sqrt{2} \Lambda}$
- Freedom in the field definitions: can choose $m_T, \kappa_{ST} > 0$

Tree-level structure + electroweak corrections:

$$(\Delta m_{hc})^{\text{ew}} \begin{pmatrix} \chi_h \\ \chi^+ \\ \chi_l \end{pmatrix} \begin{pmatrix} (\Delta m_{hc})^{\text{mix}} \\ \Delta m_{cl} \end{pmatrix}$$


Small μ :

$$\begin{pmatrix} \chi^+ \\ \chi_h \\ \chi_l \end{pmatrix} \lesssim (\Delta m_{hc})^{\text{ew}}$$

Dark matter candidate


Couplings of dark fermions

Gauge and Higgs couplings



Couplings of dark fermions

Gauge and Higgs couplings



Two physical scenarios, depending on parameters of the theory:

Scalar case:


Couplings $\propto 1, \gamma_\mu$

Pseudo-scalar case:

Couplings $\propto \gamma_5, \gamma_\mu \gamma_5$

Direct detection


Direct detection



A Feynman diagram illustrating the direct detection of dark matter. On the left, two incoming lines labeled χ_l (dark matter particles) meet at a vertex. A dashed line labeled h (Higgs boson) connects this vertex to another vertex on the right. From the right vertex, two outgoing lines labeled q (quarks) emerge. The diagram is followed by the proportionality relation $\propto \mu/v \sin(2\theta)$.

$$\propto \mu/v \sin(2\theta)$$

Direct detection




$$\propto \mu/v \sin(2\theta)$$



$$\frac{\chi_h}{\chi^+} \quad (\Delta m_{hc})^{\text{mix}} \simeq \mu \sin(2\theta)/2$$

Direct detection

[1805.12562]




$$\frac{\chi_h}{\chi^+} \quad (\Delta m_{hc})^{\text{mix}} \simeq \mu \sin(2\theta)/2$$






Direct detection

[1805.12562]




$$\frac{\chi_h}{\chi^+} \quad (\Delta m_{hc})^{\text{mix}} \simeq \mu \sin(2\theta)/2$$





Direct detection


[1805.12562]




$$\frac{\chi_h}{\chi^+} \quad (\Delta m_{hc})^{\text{mix}} \simeq \mu \sin(2\theta)/2$$



Small portal couplings needed




Surviving regions: scalar case




Annihilation

$$(g \sin \theta)^4$$



Co-annihilation

$$(g \sin \theta)^2$$




$$\begin{array}{c} \chi^+ \\ \chi_l \end{array} \overline{15 - 30 \text{ GeV}}$$

Warning: co-scattering regime

Small portal

	process	scaling		process	scaling
pair annihilation	$\chi_l \chi_l \rightarrow W^+ W^-$ $\chi_l \chi_l \rightarrow h^* \rightarrow f \bar{f}, VV$ $\chi_l \chi_l \rightarrow hh$	$(g \sin \theta)^4$ $(\mu \sin(2\theta)/v)^2$ $(\mu \cos(2\theta)/v)^4$	mediator annihilation	$\chi_h \chi_h \rightarrow W^+ W^-$ $\chi_h \chi^+ \rightarrow f \bar{f}', VV$ $\chi^+ \chi^- \rightarrow f \bar{f}, VV$	$(g \cos \theta)^2$ $(g \cos \theta)^2$ g^2
co-annihilation	$\chi_l \chi^+ \rightarrow f f', VV$ $\chi_l \chi_h \rightarrow W^+ W^-$ $\chi_l \chi_h \rightarrow h^* \rightarrow f \bar{f}, VV$	$(g \sin \theta)^2$ $(g \sin \theta)^2$ $(\mu/v)^2$	mediator decays	$\chi^+ \rightarrow \chi_l f f'$ $\chi_h \rightarrow \chi_l f \bar{f}$	$(g \sin \theta)^2$ $(\mu/v)^2$
co-scattering	$\chi_l f \rightarrow \chi^+ f'$ $\chi_l f \rightarrow \chi_h f$	$(g \sin \theta)^2$ $(\mu/v)^2$	scattering	$\chi_l f \rightarrow \chi_l f$	$(\mu \sin \theta/v)^2$




Warning: co-scattering regime

Small portal

	process	scaling		process	scaling
pair annihilation	$\chi_\ell \chi_\ell \rightarrow W^+ W^-$ $\chi_\ell \chi_\ell \rightarrow h^* \rightarrow f \bar{f}, VV$ $\chi_\ell \chi_\ell \rightarrow hh$	$(g \sin \theta)^4$ $(\mu \sin(2\theta)/v)^2$ $(\mu \cos(2\theta)/v)^4$	mediator annihilation	$\chi_h \chi_h \rightarrow W^+ W^-$ $\chi_h \chi^+ \rightarrow f \bar{f}', VV$ $\chi^+ \chi^- \rightarrow f \bar{f}, VV$	$(g \cos \theta)^2$ $(g \cos \theta)^2$ g^2
co-annihilation	$\chi_\ell \chi^+ \rightarrow f f', VV$ $\chi_\ell \chi_h \rightarrow W^+ W^-$ $\chi_\ell \chi_h \rightarrow h^* \rightarrow f \bar{f}, VV$	$(g \sin \theta)^2$ $(g \sin \theta)^2$ $(\mu/v)^2$	mediator decays	$\chi^+ \rightarrow \chi_\ell f f'$ $\chi_h \rightarrow \chi_\ell f \bar{f}$	$(g \sin \theta)^2$ $(\mu/v)^2$
co-scattering	$\chi_\ell f \rightarrow \chi^+ f'$ $\chi_\ell f \rightarrow \chi_h f$	$(g \sin \theta)^2$ $(\mu/v)^2$	scattering	$\chi_\ell f \rightarrow \chi_\ell f$	$(\mu \sin \theta/v)^2$

- Decays become very slow
- Mediators are still in chemical equilibrium
- Equilibrium is lost for dark matter
- Relic abundance is set by co-scattering + mediator annihilation




Warning: co-scattering regime

Small portal

	process	scaling		process	scaling
pair annihilation	$\chi_\ell \chi_\ell \rightarrow W^+ W^-$ $\chi_\ell \chi_\ell \rightarrow h^* \rightarrow f \bar{f}, VV$ $\chi_\ell \chi_\ell \rightarrow hh$	$(g \sin \theta)^4$ $(\mu \sin(2\theta)/v)^2$ $(\mu \cos(2\theta)/v)^4$	mediator annihilation	$\chi_h \chi_h \rightarrow W^+ W^-$ $\chi_h \chi^+ \rightarrow f \bar{f}', VV$ $\chi^+ \chi^- \rightarrow f \bar{f}, VV$	$(g \cos \theta)^4$ $(g \cos \theta)^2$ g^2
co-annihilation	$\chi_\ell \chi^+ \rightarrow f f', VV$ $\chi_\ell \chi_h \rightarrow W^+ W^-$ $\chi_\ell \chi_h \rightarrow h^* \rightarrow f \bar{f}, VV$	$(g \sin \theta)^2$ $(g \sin \theta)^2$ $(\mu/v)^2$	mediator decays	$\chi^+ \rightarrow \chi_\ell f f'$ $\chi_h \rightarrow \chi_\ell f \bar{f}$	$(g \sin \theta)^2$ $(\mu/v)^2$
co-scattering	$\chi_\ell f \rightarrow \chi^+ f'$ $\chi_\ell f \rightarrow \chi_h f$	$(g \sin \theta)^2$ $(\mu/v)^2$	scattering	$\chi_\ell f \rightarrow \chi_\ell f$	$(\mu \sin \theta/v)^2$

- Decays become very slow
- Mediators are still in chemical equilibrium
- Equilibrium is lost for dark matter
- Relic abundance is set by co-scattering + mediator annihilation




These processes are not included in public codes e.g. MicrOMEGAs!

Is a common feature of theories with nontrivial dark sector in a small-coupling regime


see [1904.07513], [1705.08450], [1705.09292] and [1906.09269] and **the talk by P. Wang tomorrow** for more examples

Small-coupling regime: phases




Co-scattering @ decoupling & colliders

Co-scattering @ decoupling & colliders



- Mediator decays are slow during decoupling.
- **But they are also long-lived at colliders!**

Co-scattering @ decoupling & colliders



- Mediator decays are slow during decoupling.
- **But they are also long-lived at colliders!**


- Both dark states present at decoupling—**compressed spectrum.**

$$\begin{array}{l} \text{M} \\ \text{DM} \end{array} \begin{array}{c} \text{—————} \\ \text{—————} \end{array} \quad \frac{\Delta m}{m} \simeq 10\%$$

Promising soft displaced signatures @ LHC

Promising soft displaced signatures @ LHC


Soft displaced lepton pair



$$\Delta M_{+,l} = 15 - 30 \text{ GeV}$$

Promising soft displaced signatures @ LHC

Soft displaced lepton pair




displaced

$$\Delta M_{+,l} = 15 - 30 \text{ GeV}$$


Promising soft displaced signatures @ LHC

Soft displaced lepton pair



$$\Delta M_{+,l} = 15 - 30 \text{ GeV}$$

Displaced soft leptons + prompt jets




$$\Delta M_{+,l} = 15 - 30 \text{ GeV}$$

$$\Delta M_{+,h} \text{ arbitrarily small}$$


Promising soft displaced signatures @ LHC

Soft displaced lepton pair



$$\Delta M_{+,l} = 15 - 30 \text{ GeV}$$


Displaced soft leptons + prompt jets



$$\Delta M_{+,l} = 15 - 30 \text{ GeV}$$


$$\Delta M_{+,h} \text{ arbitrarily small}$$

Collider searches




Collider searches


Prompt searches can not help



Lifetimes accessible @ LHC



How displaced/soft?




thermal relic

PRELIMINARY


How displaced/soft?

Current cut
[CMS-PAS-EXO-16-022]



PRELIMINARY

How displaced/soft?





PRELIMINARY

How displaced/soft?

Displacement for proposed cut →

Current cut
[CMS-PAS-EXO-16-022]




thermal relic

PRELIMINARY

Summary

- The singlet-triplet Higgs portal is a minimal fermionic extension to the SM weak sector with a naturally small coupling.
- Due to strong direct detection constraints, the thermal relic abundance of dark matter is set by mediator annihilation and co-scattering processes.
- This also leads to a natural appearance of the long-lived states. The most promising collider signatures involve displaced particles.
- New searches are needed to conclusively test this scenario: displaced soft leptons, appearing tracks, etc.




[S.Westhoff's talk at 5th LLP Workshop]

Thank you!


Backup slides

Surviving regions: pseudoscalar case

Mediator annihilation + further decays to χl




$$(g \cos \theta)^2$$



$$\begin{array}{c} \chi^+ \\ \chi l \end{array} \quad \underline{\quad 15 - 30 \text{ GeV} \quad}$$

LHC: prompt and displaced soft particles



Displaced jets


Nagata, Otono, Shirai [1506.08206]

Disappearing charged tracks


ATLAS [1712.02118]
Mahbubani, Schwaller, Zurita [1703.05327]

Soft leptons

Bharuchaa, Brümmer, Desai [1804.02357]
CMS [1206.3949]



Collider searches



Going to the mass basis: two cases

Scalar scenario

$$m_S > \mu^2/m_T$$

$$\begin{pmatrix} \chi_\ell \\ \chi_h \end{pmatrix} = \begin{pmatrix} \cos \theta \chi_S^0 - \sin \theta \chi_T^0 \\ \sin \theta \chi_S^0 + \cos \theta \chi_T^0 \end{pmatrix}$$

$$m_{h,\ell} = \frac{1}{2} \left(m_T + m_S \pm \Delta m_{h\ell} \right)$$

$$m_c = m_T$$

Pseudo-scalar scenario

$$m_S < \mu^2/m_T$$

$$\begin{pmatrix} \chi_\ell \\ \chi_h \end{pmatrix} = \begin{pmatrix} \cos \theta \chi_S^0 + \sin \theta i\gamma_5 \chi_T^0 \\ \sin \theta i\gamma_5 \chi_S^0 + \cos \theta \chi_T^0 \end{pmatrix}$$

$$m'_{h,\ell} = \frac{1}{2} \left(\Delta m_{h\ell} \pm (m_T + m_S) \right) = \pm m_{h,\ell}$$




$$m_c = m_T$$

$$\text{with } \Delta m_{h\ell} = \sqrt{(m_T - m_S)^2 + 4\mu^2} \quad \theta \simeq \frac{\mu}{m_T - m_S}$$


In both scenarios $m_\ell/m'_\ell > 0$

Disappearing charged tracks

Charged mediator leaves track in the inner detector




Relic density



Higgs resonance
region

$$\theta \simeq \frac{\mu}{m_T - m_S}$$



Blue/green: $\Omega_\psi h^2 = 0.1199 \pm 0.0022$

[Planck coll., 1502.01589]

Scalar vs. Pseudo-scalar decay rates


$$\Gamma_\psi = \theta^2 G_F^2 \Delta M^x \times \text{PS}$$

ΔM^5 (3-body) or $f_\pi^2 \Delta M^3$ (2-body)

Example:


$$\chi_+ \rightarrow \chi_h \pi^+ : \text{PS} = \left[1 - \left(\frac{m_\pi}{\Delta M} \right)^2 \right]^{1/2}$$

For small mass splittings:



$$\Gamma_+^{\text{sc}} \propto \frac{\Delta M_{Cl}^5}{120}$$

$$\Gamma_+^{\text{ps-sc}} \propto \frac{\Delta M_{Cl}^5}{40}$$



$$\Gamma_h^{\text{sc}} \propto \frac{\Delta M_{hl}^5}{60}$$

$$\Gamma_h^{\text{ps-sc}} \propto \frac{\Delta M_{hl}^7}{560}$$