


$(g - 2)_\mu$ RECAP


$(g - 2)_\mu$: **discrepancy** between exp vs theory ($\gtrsim 3\sigma$)
hadronic contributions dominate the error



$$a_\mu = \frac{\alpha}{\pi} \int \frac{ds}{s} K(s, m_\mu) \frac{\text{Im}\Pi(s)}{\pi} \quad [\text{Brodsky, de Rafael '68}]$$

analyticity $\hat{\Pi}(s) = \Pi(s) - \Pi(0) = \frac{s}{\pi} \int_{4m_\pi^2}^{\infty} dx \frac{\text{Im}\Pi(x)}{x(x - s - i\epsilon)}$

unitarity



$$\text{Im} \left[\text{vacuum polarization} \right] = \sum_X \left| \text{transition to } X \right|^2$$



$$\frac{4\pi^2\alpha}{s} \frac{\text{Im}\Pi(s)}{\pi} = \sigma_{e^+e^- \rightarrow \gamma^* \rightarrow \text{had}}$$

At present $O(30)$ channels: $\pi^0\gamma, \pi^+\pi^-, 3\pi, 4\pi, K^+K^-, \dots$

$\pi\pi$ channel is $\sim 70\%$ of signal and $\sim 70\%$ of error




MOTIVATIONS FOR τ



$V - A$ current

Final states $I = 1$ charged



EM current

Final states $I = 0, 1$ neutral

τ data can improve $a_\mu[\pi\pi]$
 $\rightarrow 72\%$ of total Hadronic LO

or $a_\mu^{ee} \neq a^\tau \rightarrow \text{NP}$ [Cirigliano et al '18]



ISOSPIN CORRECTIONS

Restriction to $e^+e^- \rightarrow \pi^+\pi^-$ and $\tau^- \rightarrow \pi^-\pi^0\nu_\tau$

$$v_0(s) = \frac{s}{4\pi\alpha^2} \sigma_{\pi^+\pi^-(\gamma)}(s)$$

$$v_-(s) = \frac{m_\tau^2}{6|V_{ud}|^2} \frac{\mathcal{B}_{\pi\pi^0}}{\mathcal{B}_e} \frac{1}{N_{\pi\pi^0}} \frac{dN_{\pi\pi^0}}{ds} \left(1 - \frac{s}{m_\tau^2}\right)^{-1} \left(1 + \frac{2s}{m_\tau^2}\right)^{-1} \frac{1}{S_{EW}}$$

Isospin correction $v_0 = R_{IB}v_-$ $R_{IB} = \frac{\text{FSR}}{G_{EM}} \frac{\beta_0^3 |F_\pi^0|^2}{\beta_-^3 |F_\pi^-|^2}$ [Alemani et al. '98]

0. S_{EW} electro-weak radiative correct. [Marciano, Sirlin '88][Braaten, Li '90]

1. Final State Radiation of $\pi^+\pi^-$ system [Schwinger '89][Drees, Hikasa '90]

2. G_{EM} (long distance) radiative corrections in τ decays

Chiral Resonance Theory [Cirigliano et al. '01, '02]

Meson Dominance [Flores-Talpa et al. '06, '07]

3. Phase Space ($\beta_{0,-}$) due to $(m_{\pi^\pm} - m_{\pi^0})$



LONG DISTANCE QED - I

At low energies relevant degrees of freedom are mesons

Chiral Perturbation Theory


[Cirigliano et al. '01, '02]

Meson dominance model

[Flores-Talpa et al. '06, '07]

Corrections casted in one function $v_-(s) \rightarrow v_-(s) G_{\text{EM}}(s)$


Real photon corrections



Real + virtual


→ IR divergences cancel

Virtual photon corrections



PION FORM FACTORS



$$F_{\pi}^0(s) \propto \frac{m_{\rho}^2}{D_{\rho}(s)}$$




[Gounaris, Sakurai '68]

[Kühn, Santamaria '90]


$$\times \left[1 + \delta_{\rho\omega} \frac{s}{D_{\omega}(s)} \right]$$



$$+ \frac{m_X^2}{D_X(s)} \quad X = \rho', \rho''$$



$$F_{\pi}^{-}(s) \propto \frac{m_{\rho^{-}}^2}{D_{\rho^{-}}(s)} + (\rho', \rho'')$$



Sources of IB breaking in phenomenological models

$$m_{\rho^0} \neq m_{\rho^{\pm}}, \Gamma_{\rho^0} \neq \Gamma_{\rho^{\pm}}, m_{\pi^0} \neq m_{\pi^{\pm}}$$

$$\rho - \omega \text{ mixing } \delta_{\rho\omega} \simeq O(m_u - m_d) + O(e^2)$$



STATUS

$$a_{\mu}^{\text{HVP,LO}}[\pi\pi, ee] = 503.51(3.5) \times 10^{-10} \text{ with } E \in [2m_{\pi}, 1.8 \text{ GeV}]$$

$$a_{\mu}^{\text{HVP,LO}}[\pi\pi, \tau] = 531.3(3.3) \times 10^{-10}$$

$$a_{\mu}[\pi\pi, ee] - a_{\mu}[\pi\pi, \tau] = -12.0(2.6) \quad [\text{Cirigliano et al.}]$$

$$a_{\mu}[\pi\pi, ee] - a_{\mu}[\pi\pi, \tau] = -16.1(1.8) \quad [\text{Davier et al. '09}]$$

(≈ -12 due to S_{EW} , rest R_{IB})

$$a_{\mu}[\tau] : \left\{ \begin{array}{l} \text{model dependence} \\ e^+e^- \text{ data more precise} \end{array} \right. = \text{abandoned}$$

Additional $\rho\gamma$ mixing correction

[Jegerlehner, Szafron '11]

partly accounted in $m_{\rho^0} - m_{\rho^-}$ in [Davier et al. '09]

$$a_{\mu}[\pi\pi, ee] = 385.2(1.5) \text{ with } E \in [0.582 - 0.975] \text{ GeV}$$

$$a_{\mu}[\pi\pi, \tau] = 386.0(2.4) \text{ after } R_{IB}$$



CONTRIBUTION TO a_μ

Time-momentum representation

[Bernecker, Meyer, '11]

$$G^\gamma(t) = \frac{1}{3} \sum_k \int d\vec{x} \langle j_k^\gamma(x) j_k^\gamma(0) \rangle \rightarrow a_\mu = 4\alpha^2 \sum_t w_t G^\gamma(t)$$

Isospin decomposition of u, d current

$$j_\mu^\gamma = \frac{i}{6} (\bar{u}\gamma_\mu u + \bar{d}\gamma_\mu d) + \frac{i}{2} (\bar{u}\gamma_\mu u - \bar{d}\gamma_\mu d) = j_\mu^{(0)} + j_\mu^{(1)}$$

$$G_{00}^\gamma \leftarrow \langle j_k^{(0)}(x) j_k^{(0)}(0) \rangle = \text{[diagram: circle with arrow and dot on left]} + \text{[diagram: circle with arrow and dot on right]} + \text{[diagram: circle with arrow and dot on top]} + \text{[diagram: circle with arrow and dot on bottom and wavy line]} + \text{[diagram: circle with arrow and dot on top and cross]} + \dots$$

$$G_{01}^\gamma \leftarrow \langle j_k^{(0)}(x) j_k^{(1)}(0) \rangle = \text{[diagram: circle with arrow and dot on bottom and wavy line]} + \text{[diagram: circle with arrow and dot on top and cross]} + \dots$$

$$G_{11}^\gamma \leftarrow \langle j_k^{(1)}(x) j_k^{(1)}(0) \rangle = \text{[diagram: circle with arrow and dot on left]} + \text{[diagram: circle with arrow and dot on bottom and wavy line]} + \text{[diagram: circle with arrow and dot on top and cross]} + \dots$$

Decompose $a_\mu = a_\mu^{(0,0)} + a_\mu^{(0,1)} + a_\mu^{(1,1)}$



NEUTRAL VS CHARGED

$$\frac{i}{2}(\bar{u}\gamma_\mu u - \bar{d}\gamma_\mu d), \left[\begin{array}{c} I=1 \\ I_3=0 \end{array} \right] \rightarrow j_\mu^{(1,-)} = \frac{i}{\sqrt{2}}(\bar{u}\gamma_\mu d), \left[\begin{array}{c} I=1 \\ I_3=-1 \end{array} \right]$$

$$\text{Isospin 1 charged correlator } G_{11}^W = \frac{1}{3} \sum_k \int d\vec{x} \langle j_k^{(1,+)}(x) j_k^{(1,-)}(0) \rangle$$

$$\delta G^{(1,1)} \equiv G_{11}^\gamma - G_{11}^W \quad [\text{MB et al.' Latt18}]$$

$$= Z_V^4 (4\pi\alpha) \frac{(Q_u - Q_d)^4}{4} \left[\text{Bubble} + \text{Bubble} \right]$$


$$G_{01}^\gamma = Z_V^4 \frac{(Q_u^2 - Q_d^2)^2}{2} (4\pi\alpha) \left[\text{Bubble} + 2 \times \text{Bubble} + \text{Bubble} + \dots \right]$$

$$+ Z_V^2 \frac{Q_u^2 - Q_d^2}{2} (m_u - m_d) \left[2 \times \text{Bubble} + \dots \right]$$

... = subleading diagrams currently not included




SYNERGY - I



from QCD we need a **4-point function** $f(x, y, z, t)$:
known kernel with details of photons and muon line
1 pair of point sources (x, y) , sum over z, t exact at sink
stochastic sampling over (x, y) (based on $|x - y|$)

Successful strategy: x10 error reduction

[RBC '16]



from QCD we need a **4-point function** $f(x, y, z, t)$:
 $(g - 2)_\mu$ kernel + photon propagator

Similar problem $\rightarrow$ re-use HLbL point sources!



The RBC & UKQCD collaborations

[BNL and BNL/RBRC](#)

Yasumichi Aoki (KEK)
Taku Izubuchi
Yong-Chull Jang
Chulwoo Jung
Meifeng Lin
Aaron Meyer
Hiroshi Ohki
Shigemi Ohta (KEK)
Amarjit Soni

[UC Boulder](#)

Oliver Witzel

[CERN](#)

Mattia Bruno

[Columbia University](#)

Ryan Abbot
Norman Christ
Duo Guo
Christopher Kelly
Bob Mawhinney
Masaaki Tomii
Jiqun Tu

Bigeng Wang
Tianle Wang
Yidi Zhao

[University of Connecticut](#)

Tom Blum
Dan Hoying (BNL)
Luchang Jin (RBRC)
Cheng Tu

[Edinburgh University](#)

Peter Boyle
Luigi Del Debbio
Felix Erben
Vera Gülpers
Tadeusz Janowski
Julia Kettle
Michael Marshall
Fionn Ó hGáin
Antonin Portelli
Tobias Tsang
Andrew Yong
Azusa Yamaguchi

[KEK](#)

Julien Frison

[University of Liverpool](#)

Nicolas Garron

[MIT](#)

David Murphy

[Peking University](#)

Xu Feng

[University of Regensburg](#)

Christoph Lehner (BNL)

[University of Southampton](#)

Nils Asmussen
Jonathan Flynn
Ryan Hill
Andreas Jüttner
James Richings
Chris Sachrajda


[Stony Brook University](#)

Jun-Sik Yoo
Sergey Syritsyn (RBRC)



SYNERGY - II

Stat. improvements from data of HLbL project [Phys.Rev.Lett. 118 (2017)]
contribution of diagrams V, S to a_μ



$O(2000)$ point-src per conf.
 ~ 3000 combinations
 $O(10)$ configurations

$\times 4$ reduction in error

expected QED conn. error $\leq 3 \times 10^{-10} \rightarrow$ matches target



LAST SLIDE, THEN PLOTS!

Restriction to $2\pi \rightarrow$ neglect pure $I = 0$ part $a_\mu^{(0,0)}[\pi^0\gamma, 3\pi, \dots]$

$$\text{Lattice: } \Delta a_\mu[\pi\pi, \tau] = 4\alpha^2 \sum_t w_t \times [G_{01}^\gamma(t) + G_{11}^\gamma(t) - G_{11}^W(t)]$$

$$\text{Pheno: } \Delta a_\mu[\pi\pi, \tau] = \int_{4m_\pi^2}^{m_\tau^2} ds K(s) \left[v_0(s) - v_-(s) \right]$$

Conversion to Euclidean time for direct comparison



$$\Delta a_\mu[\pi\pi, \tau] = 4\alpha^2 \sum_t w_t \times \left\{ \frac{1}{12\pi^2} \int d\omega \, \omega^2 e^{-\omega t} [R_{\text{IB}}(\omega^2) - 1] v_-(\omega^2) \right\}$$



LATTICE: PRELIMINARY RESULTS - I

$$\Delta a_\mu \rightarrow G_{01} + \delta G_{11}:$$

Pure $I = 1$ only $O(\alpha)$ terms:



$$V = \text{diagram of a wavy line with two vertices}$$

$$F = \text{diagram of two loops connected by a wavy line}$$

$$S = \text{diagram of a wavy line with two vertices and a loop}$$

$$M = \text{diagram of a loop with a cross on the top line}$$

$$O = \text{diagram of two loops connected by a wavy line}$$

relevant, negative, not-included




LATTICE: PRELIMINARY RESULTS - II

Study **integrand** in euclidean time $\rightarrow$ **as important as integral**


direct comparison
Lattice vs. EFT+Pheno

1. validate previous estimates of R_{IB}
2. study neutral/charged ρ and ω properties

Preliminary lattice (full) calculation: $G_{01}^\gamma + \delta G$



Not included:

1.  **relevant**
2. **sub-leading** $1/N_c$, $SU(N_f)$

Under consideration:


3. **finite-volume** errors
4. discretization errors



SYSTEMATIC ERRORS

$$a_{\mu}^{\text{QED,conn}} = V + 2S$$

FV study at coarse
 $a^{-1} \sim 1 \text{ GeV}$



Finite volume errors

empirical observation: diagrams may have largish FV errors

cancellation of FV effects in **physical combinations**




SYSTEMATIC ERRORS - II

Lattice fully **inclusive**

1. cut $E < m_\tau$

2. higher multiplicity channels, 4π

effects above ~ 1.8 GeV suppressed by (muon) kernel



Smeared step function

$$\Theta(s) = (1 + e^{2k(E-E_0)})^{-1}$$

→ control syst. effect of 4π

manipulate $G(t)$ (e.g. Backus-Gilbert)

preliminary: w/o cut $\Delta a_\mu[4\pi] \approx 2(1)$

$$E_0 = 2 \text{ GeV}, 1/(2k) = 0.25 \text{ GeV}$$




TOWARDS A COMPARISON

Lattice contains $\pi^0\pi^-\gamma$ states $\rightarrow$ 

Re-evaluation of $G_{\text{EM}} \rightarrow G_{\text{EM}}^{\pi}$ [in collab. with Cirigliano]


Real photon corrections



G_{EM}^{π} w/o $\pi^0\pi^-\gamma$ FSR

$\frac{v_-}{G_{\text{EM}}^{\pi}}$ w $\pi^0\pi^-\gamma$ FSR

Virtual photon corrections



OUTLOOK

Inclusive studies:

- kernels to suppress high channels

- suppression of short/long distances (cutoff effects/noise)

- expand τ -decay program [with M. Gonzalez-Alonso]

- e.g. $K\pi$ channel in vector-vector correlator

- e.g. $SU(3)$ breaking $\pi\pi - K\pi$

Exclusive study: long term goal, proper isospin-breaking in $\pi\pi$ form factor



CONCLUSIONS

These are exciting times for $(g - 2)_\mu$:

1% goal for lattice results to be expected soon

QED+SIB crucial to reach target uncertainty

As a bi-product we get $\Delta a_\mu[\tau]$:

1. first lattice calculation of $\Delta a_\mu[\tau]$ almost complete

2. tests/checks previous calculations

comparing v_- with experiment requires G_{EM}^π

study G_{01}^γ alone $\rightarrow \rho\omega$ mixing; $\delta G^{(1,1)}$ alone $\rightarrow \rho^0$ vs ρ^-


3. possibly bound new physics

Thanks for your attention



EXPERIMENTAL RESULTS

$$\Delta a_\mu(t) = 4\alpha^2 \sum_t w_t \left\{ \int ds h(s, t) \left[v_0(s) - \frac{v_1(s)}{G_{\text{EM}}(s)} \right] \right\}$$



v_0 BaBar, v_1 Aleph

preliminary



$\rho\gamma$ MIXING - I

Gounaris-Sakurai based on VMD model w/o EM gauge invariance

- generation of a photon mass

- + based on phase shift (proper pion rescattering behavior)

widely used: e.g. PDG estimates of m_ρ , Γ_ρ

VMD model with gauge-invariance
at 1-loop s -dependent mass matrix

[Kroll, Lee, Zumino '67]

[Jegerlehner, Szafron '11]





limits of validity pion-loop? high enough energy must break down



$\rho\gamma$ MIXING - II

[Jegerlehner, Szafron '11]



30% correction at 1 GeV

δ_1^1 in good agreement $E < 800$ MeV


perhaps restrict the $\rho\gamma$ below 800 MeV?



$\rho\gamma$ MIXING-III

[1] = [Jegelehner, Szafron '17]

modified $\rho\gamma$ coupling
large negative Δa_μ



Modified $\rho\gamma$ suggests different behavior from lattice data

direct comparison with lattice not possible $\rightarrow$ hard cut at 1 GeV

