



Contribution ID: 271

Type: **Contributed Talk**

## Measurement of transverse single-spin asymmetries for di-jet production in polarized p+p collisions at $\sqrt{s} = 200$ GeV at STAR

*Thursday, 26 March 2020 09:20 (20 minutes)*

Our understanding of the origin of the nucleon spin remains incomplete. Determining the partonic orbital angular momentum (OAM) contributions is particularly challenging experimentally. One possible signature for the partonic OAM would be detection of a non-zero Sivers effect, which characterizes the correlation between the transverse momentum of a parton ( $\vec{k}_T$ ) and the transverse spin ( $\vec{S}$ ) of its proton, moving in the longitudinal ( $\vec{p}$ ) direction. Experimentally, the Sivers function can be accessed by searching for a preference in the bisector ray of di-jet transverse opening angle, which reverses direction when the beam polarization direction is flipped. A previous effort by STAR using  $1\text{ pb}^{-1}$  of  $p+p$  data taken in 2006 at  $\sqrt{s} = 200$  GeV did not find a significant effect due to limited statistical precision. In 2012 and 2015, STAR accumulated much larger datasets ( $\sim 33$  times the di-jets after event selection) at  $\sqrt{s} = 200$  GeV. Moreover, a new technique has been implemented to tag the flavor of the fragmenting partons, to avoid cancellation between  $u$  and  $d$  quark effects, which are expected to have opposite signs. In this talk, a preliminary result of the measurement of Sivers asymmetries using STAR 2012 and 2015  $p+p$  data at  $\sqrt{s} = 200$  GeV will be reported.

**Primary author:** LIU, Huanzhao

**Presenter:** LIU, Huanzhao

**Session Classification:** Spin Physics

**Track Classification:** Spin Physics