

XXVIII International Workshop on Deep Inelastic Scattering and Related Subjects



Contribution ID: 146

Type: **Contributed Talk**

A model calculation of unpolarized and polarized transverse-momentum-dependent distribution functions

Thursday, 26 March 2020 09:00 (20 minutes)

While significant steps toward the formal definition of quark TMDs and their extraction from experimental data through global fits has been made in the last years, the gluon-TMD field represents a largely unexplored territory. Pursuing the goal of extending our knowledge of this sector, we present analytic expressions for all T -even gluon TMDs at twist-2, calculated in a spectator model for the parent nucleon. At variance with respect to previous works, our approach encodes a flexible parametrization for the spectator-mass spectral density, allowing us to improve the description in the small- x region.

We build a common framework where valence, sea quark and gluon densities are concurrently generated. Our results can be used to predict the behavior of observables sensitive to TMD dynamics.

Primary authors: BACCHETTA, Alessandro (University of Pavia and INFN); CELIBERTO, Francesco Giovanni (University of Pavia and INFN); Prof. RADICI, Marco (INFN, Sezione di Pavia); Dr TAELS, Pieter (Ecole Polytechnique, CPHT & Università di Cagliari & INFN)

Presenter: CELIBERTO, Francesco Giovanni (University of Pavia and INFN)

Session Classification: Spin Physics

Track Classification: Spin Physics