

# XXVIII International Workshop on Deep Inelastic Scattering and Related Subjects



Contribution ID: 33

Type: **Contributed Talk**

## Accessing gluon saturation at electron-ion colliders seen via dissociative production of vector mesons

*Wednesday, 25 March 2020 12:26 (17 minutes)*

We present a model with which we predict the cross sections for exclusive and dissociative photo and electro-production of light and heavy vector mesons off protons; the model describes correctly available experimental data. The model is based on the color-dipole approach and incorporates geometric fluctuations of the target-proton partonic structure in the impact-parameter plane. The number of fluctuations grows with decreasing Bjorken- $x$  and they are generated as event-by-event randomly placed areas of high gluonic density, so-called hot spots. A striking feature of the model is the prediction of a maximum of the dissociative cross section as a function of the center-of-mass energy  $W_{\gamma p}$ , followed by a steep decrease as the hot spots start to overlap with increasing energy. We use these maxima to define a geometric saturation scale which grows linearly with energy as a function of the scale of the process. This phenomenon can be measured at the proposed electron-ion colliders such as eRHIC or LHeC. We present a comparison of their envisioned kinematic reach with the geometric saturation scale.

**Primary author:** BENDOVA, Dagmar (Czech Technical University in Prague (FNSPE))

**Co-authors:** CEPILA, Jan (Czech Technical University in Prague); CONTRERAS NUNO, Jesus Guillermo (Czech Technical University in Prague)

**Presenter:** BENDOVA, Dagmar (Czech Technical University in Prague (FNSPE))

**Session Classification:** Small- $x$ , Diffraction and Vector Mesons

**Track Classification:** Small- $x$ , Diffraction and Vector Mesons