



Contribution ID: 308

Type: **Contributed Talk**

## Non-zero transverse single spin asymmetry of very forward $\pi^0$ in polarized p+p collisions at $\sqrt{s} = 510$ GeV

*Wednesday, 25 March 2020 15:30 (20 minutes)*

Through the 21st and 22nd International Workshop on Deep Inelastic Scattering, a possibility of the non-perturbative contribution for the non-zero transverse single spin asymmetry of forward  $\pi^0$  ( $2 < \eta < 4$ ) was brought up. Bigger asymmetry was observed with more isolated final state which was connected with the non-perturbative event topology. Since the non-perturbative contribution has been studied by only the forward  $\pi^0$  production where the perturbative process was expected to be the major interaction rather than non-perturbative one, the RHICf experiment measured the very forward ( $\eta > 6$ )  $\pi^0$  to study the non-perturbative contribution in more detail. A new electromagnetic calorimeter was installed for the operation at the zero-degree region of the STAR experiment at RHIC and measured the  $\pi^0$  over the kinematic range of  $x_F > 0.25$  and  $0 < p_T < 1$  GeV/c in June, 2017. A clear non-zero asymmetry was observed even in low  $p_T < 1$  GeV/c showing a similar  $x_F$  dependence with the forward  $\pi^0$  one as the  $p_T$  approached to 1 GeV/c. We present the first measurement of the very forward  $\pi^0$  asymmetry, its analysis procedure and preliminary result. A future aspect to more precisely study the non-perturbative contribution will be also introduced.

**Primary author:** Mr KIM, Minho (Korea University/ RIKEN)

**Presenter:** Mr KIM, Minho (Korea University/ RIKEN)

**Session Classification:** Spin Physics

**Track Classification:** Spin Physics