


Composite nature of Dark Matter

© Francesco Sannino

CP³-Origins, Danish IAS, Univ. Southern Denmark

LME - BNL Decemer 2013


CP³ Origins

The logo for CP³ Origins features the text "CP³ Origins" in a white serif font. Below the text, a red horizontal line spans the width of the text. Two red arrows point towards the center of this line, one from the left and one from the right, meeting at the center under the letter 'g'.


Cosmology & Particle Physics

Dark matter

- ◆ Interacts at least gravitationally
- ◆ Electrically neutral & decoupled from primordial plasma
- ◆ Leads to density profile for galaxy rotation curves
- ◆ “if” cold: clusters & leads to structure formation
- ◆ Either stable or very long lived




What makes dark matter ?



Atoms: 3 forces and many fund. particles

What makes dark matter ?



Atoms
5%

Atoms: 3 forces and many fund. particles


DM oversimplification

DM Particle



Dark Matter
27%

What makes dark matter ?




Atoms: 3 forces and many fund. particles

DM oversimplification


DM Particle



Elementary




Composite



Incomplete DM list


Incomplete DM list

Elementary



Incomplete DM list


Elementary



- ◆ Unnatural, classical conformality, delayed naturality
- ◆ Perturbative natural conformality Antipin, Mojaza, Sannino 13
- ◆ Natural extensions [Susy,..]
- ◆ Axions
- ◆ ...


Incomplete DM list

Elementary




- ◆ Unnatural, classical conformality, delayed naturality
- ◆ Perturbative natural conformality Antipin, Mojaza, Sannino 13
- ◆ Natural extensions [Susy,..]
- ◆ Axions
- ◆ ...

Composite




Incomplete DM list

Elementary



- ◆ Unnatural, classical conformality, delayed naturality
- ◆ Perturbative natural conformality Antipin, Mojaza, Sannino 13
- ◆ Natural extensions [Susy,..]
- ◆ Axions
- ◆ ...

Composite




- ◆ New QCD-like sectors added by hand
- ◆ Composite DM from composite EW [Technicolor,..]
- ◆ Composite axions
- ◆ ...


In practice

In practice

Elementary




Composite




In practice

Elementary




Composite




- ◆ Perturbative: We can compute

In practice

Elementary




Composite




- ◆ Perturbative: We can compute
- ◆ Nonperturbative:

In practice

Elementary




Composite




- ◆ Perturbative: We can compute
- ◆ Nonperturbative:
 - ◆ Effective lagrangians
 - ◆ First principle lattice simulations

Building composite DM




Building composite DM




- ◆ Gauge theories with fermionic matter

Building composite DM




- ◆ Gauge theories with fermionic matter
- ◆ Gauge group: SU, SO, Sp, Exceptional

Building composite DM




- ◆ Gauge theories with fermionic matter
- ◆ Gauge group: SU, SO, Sp, Exceptional
- ◆ Choose matter repr.

Building composite DM



- ◆ Gauge theories with fermionic matter
- ◆ Gauge group: SU, SO, Sp, Exceptional
- ◆ Choose matter repr.
- ◆ Embed SM [e.g. Dynamical EWSB]


Building composite DM



- ◆ Gauge theories with fermionic matter
- ◆ Gauge group: SU, SO, Sp, Exceptional
- ◆ Choose matter repr.
- ◆ Embed SM [e.g. Dynamical EWSB]

Resulting composite DM can be

Building composite DM




- ◆ Gauge theories with fermionic matter
- ◆ Gauge group: SU, SO, Sp, Exceptional
- ◆ Choose matter repr.
- ◆ Embed SM [e.g. Dynamical EWSB]

Resulting composite DM can be

- ◆ A fermion

Building composite DM




- ◆ Gauge theories with fermionic matter
- ◆ Gauge group: SU, SO, Sp, Exceptional
- ◆ Choose matter repr.
- ◆ Embed SM [e.g. Dynamical EWSB]

Resulting composite DM can be

- ◆ A fermion
- ◆ A (Goldstone) boson

Building composite DM




- ◆ Gauge theories with fermionic matter
- ◆ Gauge group: SU, SO, Sp, Exceptional
- ◆ Choose matter repr.
- ◆ Embed SM [e.g. Dynamical EWSB]

Resulting composite DM can be

- ◆ A fermion
- ◆ A (Goldstone) boson

Solve new strong dynamics!




(2-flavors) QCD - like

- ◆ Number of odd colors > 3
- ◆ Complex representations
- ◆ If below conformal window

$$SU(2) \times SU(2) \rightarrow SU(2)$$

- ◆ In Technicolor-like embedding the TC-neutron = DM (1 - 3 TeV)
- ◆ Different flavors/SM embedding more possibilities



$SU(2) = Sp(2)$ - Template

Appelquist, Sannino, 98, 99

Ryttov, Sannino, 2008


Järvinen, Ryttov, Sannino, 2009

Lewis, Pica, Sannino 2012


Hietanen, Lewis, Pica, Sannino 2013

Sp(2)

$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}^a F^{a\mu\nu} + \overline{Q}(i\gamma^\mu D_\mu)Q$$



Sp(2)




$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}^a F^{a\mu\nu} + \overline{Q}(i\gamma^\mu D_\mu)Q$$

$$Q = \begin{pmatrix} U_L \\ D_L \\ -i\sigma^2 C \overline{U}_R^T \\ -i\sigma^2 C \overline{D}_R^T \end{pmatrix}$$

◆ Act. invariant under SU(4) transf.

$$Q \rightarrow \left(1 + i \sum_{k=1}^{15} \alpha^k T^k \right) Q$$

Sp(2)



$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}^a F^{a\mu\nu} + \overline{Q}(i\gamma^\mu D_\mu)Q$$

$$Q = \begin{pmatrix} U_L \\ D_L \\ -i\sigma^2 C \overline{U}_R^T \\ -i\sigma^2 C \overline{D}_R^T \end{pmatrix}$$


◆ Act. invariant under SU(4) transf.

$$Q \rightarrow \left(1 + i \sum_{k=1}^{15} \alpha^k T^k \right) Q$$


◆ Mass term respects Sp(4) with 10 generators

$$\delta\mathcal{L} = \frac{m}{2} Q^T (-i\sigma^2) C E Q + \text{h.c.} \quad E = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix}$$

Theoretical expectations




Theoretical expectations



- ◆ @ m=0

$$\langle Q^T (-i\sigma^2) C EQ \rangle = \langle \bar{U}U + \bar{D}D \rangle = \Lambda^3 \neq 0$$

Theoretical expectations




- ◆ @ $m=0$

$$\langle Q^T (-i\sigma^2) C EQ \rangle = \langle \bar{U}U + \bar{D}D \rangle = \Lambda^3 \neq 0$$

- ◆ SU(4) can break spontaneously to Sp(4)
- ◆ Composite spectrum are representations of Sp(4)
- ◆ 5 Goldstones

Hadronic operators



- ◆ Mesons


$$\mathcal{O}_{\overline{U}D}^{(\Gamma)} \equiv \overline{U}(x)\Gamma D(x) \ ,$$

$$\mathcal{O}_{\overline{D}U}^{(\Gamma)} \equiv \overline{D}(x)\Gamma U(x) \ ,$$

$$\mathcal{O}_{\overline{U}U \pm \overline{D}D}^{(\Gamma)} \equiv \frac{1}{\sqrt{2}} \left(\overline{U}(x)\Gamma U(x) \pm \overline{D}(x)\Gamma D(x) \right)$$

$$\Gamma = 1, \gamma^5, \gamma^\mu, \dots$$

Hadronic operators



- ◆ Mesons

$$\mathcal{O}_{\overline{U}D}^{(\Gamma)} \equiv \overline{U}(x)\Gamma D(x) ,$$

$$\mathcal{O}_{\overline{D}U}^{(\Gamma)} \equiv \overline{D}(x)\Gamma U(x) ,$$


$$\mathcal{O}_{\overline{U}U\pm\overline{D}D}^{(\Gamma)} \equiv \frac{1}{\sqrt{2}} \left(\overline{U}(x)\Gamma U(x) \pm \overline{D}(x)\Gamma D(x) \right)$$

$$\mathcal{O}_{UD}^{(\Gamma)} \equiv U^T(x)(-i\sigma^2)C\Gamma D(x) , \quad \text{◆ Baryons (diquarks)}$$


$$\mathcal{O}_{DU}^{(\Gamma)} \equiv D^T(x)(-i\sigma^2)C\Gamma U(x) ,$$

$$\mathcal{O}_{UU\pm DD}^{(\Gamma)} \equiv \frac{1}{\sqrt{2}} \left(U^T(x)(-i\sigma^2 C)\Gamma U(x) \pm D^T(x)(-i\sigma^2 C)\Gamma D(x) \right)$$

$$\Gamma = 1, \gamma^5, \gamma^\mu, \dots$$




Facts



Facts


- ◆ Mesons and baryons are mass-degenerate
- ◆ Equal angular momentum & opposite parity



Facts

- ◆ Mesons and baryons are mass-degenerate
- ◆ Equal angular momentum & opposite parity

$$\begin{aligned} J \left(\mathcal{O}_{UD}^{(\Gamma)} \right) &= J \left(\mathcal{O}_{\bar{U}D}^{(\Gamma)} \right), \\ P \left(\mathcal{O}_{UD}^{(\Gamma)} \right) &= -P \left(\mathcal{O}_{\bar{U}D}^{(\Gamma)} \right) \end{aligned}$$




Facts

- ◆ Mesons and baryons are mass-degenerate
- ◆ Equal angular momentum & opposite parity

$$\begin{aligned} J \left(\mathcal{O}_{UD}^{(\Gamma)} \right) &= J \left(\mathcal{O}_{\bar{U}D}^{(\Gamma)} \right), \\ P \left(\mathcal{O}_{UD}^{(\Gamma)} \right) &= -P \left(\mathcal{O}_{\bar{U}D}^{(\Gamma)} \right) \end{aligned}$$

- ◆ 5 Goldstones



Facts

- ◆ Mesons and baryons are mass-degenerate
- ◆ Equal angular momentum & opposite parity


$$\begin{aligned} J \left(\mathcal{O}_{UD}^{(\Gamma)} \right) &= J \left(\mathcal{O}_{\overline{U}D}^{(\Gamma)} \right), \\ P \left(\mathcal{O}_{UD}^{(\Gamma)} \right) &= -P \left(\mathcal{O}_{\overline{U}D}^{(\Gamma)} \right) \end{aligned}$$

- ◆ 5 Goldstones

pseudoscalar

Π^+

Π^-



Facts

- ◆ Mesons and baryons are mass-degenerate
- ◆ Equal angular momentum & opposite parity

$$\begin{aligned} J \left(\mathcal{O}_{UD}^{(\Gamma)} \right) &= J \left(\mathcal{O}_{\overline{U}D}^{(\Gamma)} \right), \\ P \left(\mathcal{O}_{UD}^{(\Gamma)} \right) &= -P \left(\mathcal{O}_{\overline{U}D}^{(\Gamma)} \right) \end{aligned}$$


- ◆ 5 Goldstones

pseudoscalar

Π^+ Π^-

scalar baryon

Π_{UD} $\Pi_{\overline{UD}}$



Facts

- ◆ Mesons and baryons are mass-degenerate
- ◆ Equal angular momentum & opposite parity

$$\begin{aligned} J \left(\mathcal{O}_{UD}^{(\Gamma)} \right) &= J \left(\mathcal{O}_{\overline{UD}}^{(\Gamma)} \right), \\ P \left(\mathcal{O}_{UD}^{(\Gamma)} \right) &= -P \left(\mathcal{O}_{\overline{UD}}^{(\Gamma)} \right) \end{aligned}$$

- ◆ 5 Goldstones

pseudoscalar

Π^+

Π^-


Π^0

scalar baryon

Π_{UD}

$\Pi_{\overline{UD}}$

Minimal TC Model & DM




Appelquist, Sannino, 98, 99

Ryttov, Sannino, 2008

Järvinen, Ryttov, Sannino, 2009

Minimal TC Model & DM



- ◆ Gauge $SU_L(2) \times U_Y(1)$ in $SU(4)$


$$f_\Pi \simeq 246 \text{ GeV}$$

Appelquist, Sannino, 98, 99

Ryttov, Sannino, 2008

Järvinen, Ryttov, Sannino, 2009

Minimal TC Model & DM



- ◆ Gauge $SU_L(2) \times U_Y(1)$ in $SU(4)$

$$f_\Pi \simeq 246 \text{ GeV}$$


- ◆ Such that

Appelquist, Sannino, 98, 99

Ryttov, Sannino, 2008

Järvinen, Ryttov, Sannino, 2009

Minimal TC Model & DM



- ◆ Gauge $SU_L(2) \times U_Y(1)$ in $SU(4)$

$$f_\Pi \simeq 246 \text{ GeV}$$

Appelquist, Sannino, 98, 99


Ryttov, Sannino, 2008

Järvinen, Ryttov, Sannino, 2009

- ◆ Such that

Π^+ Π^- Π^0 Longitudinal W and Z

Minimal TC Model & DM



- ◆ Gauge $SU_L(2) \times U_Y(1)$ in $SU(4)$

$$f_\Pi \simeq 246 \text{ GeV}$$

Appelquist, Sannino, 98, 99

Ryttov, Sannino, 2008


Järvinen, Ryttov, Sannino, 2009

- ◆ Such that

Π^+ Π^- Π^0 Longitudinal W and Z

Π_{UD} $\Pi_{\overline{UD}}$ Goldstone DM & anti-DM, SM singlet

Minimal TC Model & DM



- ◆ Gauge $SU_L(2) \times U_Y(1)$ in $SU(4)$

Appelquist, Sannino, 98, 99

Ryttov, Sannino, 2008

Järvinen, Ryttov, Sannino, 2009

$$f_\Pi \simeq 246 \text{ GeV}$$


- ◆ Such that

Π^+ Π^- Π^0 Longitudinal W and Z

Π_{UD} $\Pi_{\overline{UD}}$ Goldstone DM & anti-DM, SM singlet

- ◆ Fermion mass generation sectors can yield mass to DM

Minimal TC Model & DM



- ◆ Gauge $SU_L(2) \times U_Y(1)$ in $SU(4)$

Appelquist, Sannino, 98, 99

Ryttov, Sannino, 2008

Järvinen, Ryttov, Sannino, 2009

$$f_\Pi \simeq 246 \text{ GeV}$$


- ◆ Such that

$\Pi^+ \quad \Pi^- \quad \Pi^0$ Longitudinal W and Z

$\Pi_{UD} \quad \Pi_{\overline{UD}}$ Goldstone DM & anti-DM, SM singlet

- ◆ Fermion mass generation sectors can yield mass to DM

Dark matter on Lattice




Lewis, Pica, Sannino 2012

Hietanen, Pica, Sannino, Søndergaard 2013

Hietanen, Lewis, Pica, Sannino 2013

Dark matter on Lattice




Lewis, Pica, Sannino 2012


Hietanen, Pica, Sannino, Søndergaard 2013

Hietanen, Lewis, Pica, Sannino 2013

Canadian-Danish collaboration




Dark Wilson action



$$\begin{aligned}
 S_W = & \frac{\beta}{2} \sum_{x,\mu,\nu} \left(1 - \frac{1}{2} \text{ReTr} U_\mu(x) U_\nu(x + \hat{\mu}) U_\mu^\dagger(x + \hat{\nu}) U_\nu^\dagger(x) \right) \\
 & + (4 + m_0) \sum_x \bar{\psi}(x) \psi(x) \\
 & - \frac{1}{2} \sum_{x,\mu} \left(\bar{\psi}(x) (1 - \gamma_\mu) U_\mu(x) \psi(x + \hat{\mu}) + \bar{\psi}(x + \hat{\mu}) (1 + \gamma_\mu) U_\mu^\dagger(x) \psi(x) \right)
 \end{aligned}$$

β	Volume	m_0	Therm.	Conf.
2.0	$16^3 \times 32$	-0.85, -0.9, -0.94, -0.945, -0.947, -0.949	320	680
2.0	32^4	-0.947	500	680
2.2	$16^3 \times 32$	-0.60, -0.65, -0.68, -0.70, -0.72, -0.75	320	680
2.2	$24^3 \times 32$	-0.75	500	~ 2000
2.2	32^4	-0.72, -0.735, -0.75	500	~ 2000

Table 1: Parameters used in the simulations. The thermalization column refers to the number of discarded initial configurations and the configuration column refers to the number of independent configurations used in measurements.




◆ Fermion mass via PCAC

$$m_q = \lim_{t \rightarrow \infty} \frac{1}{2} \frac{\partial_t V_\Pi}{V_{PP}}$$

$$V_\Pi(t_i - t_f) = a^3 \sum_{x_1, x_2, x_3} \langle \bar{u}(t_i) \gamma_0 \gamma_5 d(t_i) \bar{u}(t_f) \gamma_5 d(t_f) \rangle$$

$$V_{PP}(t_i - t_f) = a^3 \sum_{x_1, x_2, x_3} \langle \bar{u}(t_i) \gamma_5 d(t_i) \bar{u}_1(t_f) \gamma_5 d(t_f) \rangle$$



◆ Fermion mass via PCAC

$$m_q = \lim_{t \rightarrow \infty} \frac{1}{2} \frac{\partial_t V_\Pi}{V_{PP}}$$


$$V_\Pi(t_i - t_f) = a^3 \sum_{x_1, x_2, x_3} \langle \bar{u}(t_i) \gamma_0 \gamma_5 d(t_i) \bar{u}(t_f) \gamma_5 d(t_f) \rangle$$

$$V_{PP}(t_i - t_f) = a^3 \sum_{x_1, x_2, x_3} \langle \bar{u}(t_i) \gamma_5 d(t_i) \bar{u}_1(t_f) \gamma_5 d(t_f) \rangle$$

◆ Decay constant

$$f_\Pi = \frac{m_q}{m_\Pi^2} G_\Pi$$

$$V_{PP}(t_i - t_f) = -\frac{G_\Pi^2}{m_\Pi} \exp[-m_\Pi(t_i - t_f)]$$




◆ Fermion mass via PCAC

$$m_q = \lim_{t \rightarrow \infty} \frac{1}{2} \frac{\partial_t V_\Pi}{V_{\text{PP}}}$$

$$V_\Pi(t_i - t_f) = a^3 \sum_{x_1, x_2, x_3} \langle \bar{u}(t_i) \gamma_0 \gamma_5 d(t_i) \bar{u}(t_f) \gamma_5 d(t_f) \rangle$$

$$V_{\text{PP}}(t_i - t_f) = a^3 \sum_{x_1, x_2, x_3} \langle \bar{u}(t_i) \gamma_5 d(t_i) \bar{u}_1(t_f) \gamma_5 d(t_f) \rangle$$



◆ Decay constant


$$f_\Pi = \frac{m_q}{m_\Pi^2} G_\Pi$$


$$V_{\text{PP}}(t_i - t_f) = -\frac{G_\Pi^2}{m_\Pi} \exp[-m_\Pi(t_i - t_f)]$$

◆ 2 point functions

$$\begin{aligned} C_{\bar{u}d}^{(\Gamma)}(t_i - t_f) &= \sum_{\vec{x}_i, \vec{x}_f} \langle \mathcal{O}_{ud}^{(\Gamma)}(x_f) \mathcal{O}_{ud}^{(\Gamma)\dagger}(x_i) \rangle \\ &= \sum_{\vec{x}_i, \vec{x}_f} \text{Tr} \Gamma S_{d\bar{d}}(x_f, x_i) \gamma^0 \Gamma^\dagger \gamma^0 S_{u\bar{u}}(x_i, x_f) \end{aligned}$$


$$S_{u\bar{u}}(x, y) = \langle u(x) \bar{u}(y) \rangle$$





◆ Chiral symmetry breaks


$$m_\Pi^2 \propto m_q$$




◆ Chiral symmetry breaks

$$m_\Pi^2 \propto m_q$$

◆ Nonzero decay constant






◆ Chiral symmetry breaks


$$m_\Pi^2 \propto m_q$$

◆ Nonzero decay constant

◆ Perturbative estimate of Z_a

$$Z_a = 1 - \frac{g_0^2}{16\pi^2} \frac{N^2 - 1}{2N} 15.7 \stackrel{N=2}{=} 1 - 0.2983/\beta$$





◆ Chiral symmetry breaks


$$m_\Pi^2 \propto m_q$$

◆ Don't see yet the chiral logs

◆ Nonzero decay constant




◆ Perturbative estimate of Z_a

$$Z_a = 1 - \frac{g_0^2}{16\pi^2} \frac{N^2 - 1}{2N} 15.7 \stackrel{N=2}{=} 1 - 0.2983/\beta$$





Spin one spectrum

- ◆ 2 Volumes: $16^3 \times 32$ and 32^4 in lighter colors
- ◆ Smaller volume effects for beta = 2.0
- ◆ Larger volume effects for the most chiral points for beta = 2.2
- ◆ Volume corrections for lighter masses



Continuum estimates



$$m_\rho \simeq 2510(40)(280)(300) \text{ GeV}$$


$$f_\Pi \simeq 246 \text{ GeV}$$

$$m_A \simeq 3270(130)(370)(370) \text{ GeV}$$

(stat.) (continuum) (Z_a)


DM-photon interaction

- ◆ U_L and D_L form a weak doublet
- ◆ U_R and D_R fields are weak singlets
- ◆ U (D) have $1/2$ ($-1/2$) *electric charge*
- ◆ No gauge and Witten anomalies



DM-photon interaction

- ◆ U_L and D_L form a weak doublet
- ◆ U_R and D_R fields are weak singlets
- ◆ U (D) have $1/2$ ($-1/2$) *electric charge*
- ◆ No gauge and Witten anomalies




Dark coupling to photon via charge radius

$$\phi \equiv \Pi_{UD}$$

DM-photon interaction

- ◆ U_L and D_L form a weak doublet
- ◆ U_R and D_R fields are weak singlets
- ◆ U (D) have $1/2$ ($-1/2$) *electric charge*
- ◆ No gauge and Witten anomalies




Dark coupling to photon via charge radius

$$\phi \equiv \Pi_{UD}$$

$$\mathcal{L}_\gamma = ie \frac{d_B}{\Lambda^2} \phi^* \overleftrightarrow{\partial}_\mu \phi \partial_\nu F^{\mu\nu}$$

DM-photon interaction

- ◆ U_L and D_L form a weak doublet
- ◆ U_R and D_R fields are weak singlets
- ◆ U (D) have $1/2$ ($-1/2$) *electric charge*
- ◆ No gauge and Witten anomalies



Dark coupling to photon via charge radius


$$\phi \equiv \Pi_{UD}$$

$$\mathcal{L}_\gamma = ie \frac{d_B}{\Lambda^2} \phi^* \overleftrightarrow{\partial}_\mu \phi \partial_\nu F^{\mu\nu}$$

$$\frac{d_B}{\Lambda^2} = ?$$

DM-photon interaction

- ◆ U_L and D_L form a weak doublet
- ◆ U_R and D_R fields are weak singlets
- ◆ U (D) have $1/2$ ($-1/2$) *electric charge*
- ◆ No gauge and Witten anomalies



Dark coupling to photon via charge radius





$$\phi \equiv \Pi_{UD}$$

$$\mathcal{L}_\gamma = ie \frac{d_B}{\Lambda^2} \phi^* \overleftrightarrow{\partial}_\mu \phi \partial_\nu F^{\mu\nu}$$

$$\frac{d_B}{\Lambda^2} = ?$$

- ◆ Need to determine 3-point functions (GBs form factors)

GBs vector form factors




◆ Lattice electromagnetic current

$$V_\mu(x) = \frac{1}{2} V_\mu^U(x) - \frac{1}{2} V_\mu^D(x)$$

$$V_\mu^X(x) = \frac{1}{2} \bar{X}(x + \hat{\mu})(1 + \gamma_\mu) U_\mu^\dagger(x) X(x) - \frac{1}{2} \bar{X}(x)(1 - \gamma_\mu) U_\mu(x) X(x + \hat{\mu})$$

$$F_\Pi(Q^2) = \frac{C_{ud}^{(3)}(t_i, t, t_f, \vec{p}_i, \vec{p}_f) C_{ud}^{(2)}(t_i, t, \vec{p}_f)}{C_{ud}^{(2)}(t_i, t, \vec{p}_i) C_{ud}^{(2)}(t_i, t_f, \vec{p}_f)} \left(\frac{2E_\Pi(\vec{p}_f)}{E_\Pi(\vec{p}_i) + E_\Pi(\vec{p}_f)} \right)$$

Avoiding disconnected diagrams



- ◆ Lattice simulation expensive for different fermion masses
- ◆ But for $m_U = m_D$ form factors vanish
- ◆ Up and Down are related in QCD at large N [works for N=3]


$$F_{\pi^+}(Q^2) \approx \frac{2}{3} \left(\frac{m_\rho^2}{m_\rho^2 + Q^2} \right) + \frac{1}{3} \left(\frac{m_\rho^2}{m_\rho^2 + Q^2} \right)$$

$$F_{K^+}(Q^2) \approx \frac{2}{3} \left(\frac{m_\rho^2}{m_\rho^2 + Q^2} \right) + \frac{1}{3} \left(\frac{m_\phi^2}{m_\phi^2 + Q^2} \right)$$

$$F_{K^0}(Q^2) \approx -\frac{1}{3} \left(\frac{m_\rho^2}{m_\rho^2 + Q^2} \right) + \frac{1}{3} \left(\frac{m_\phi^2}{m_\phi^2 + Q^2} \right)$$

We will test VMD for SU(2)

GB form factor vs vector pole




Lattice prediction

- ◆ Form factor requires isospin breaking from ETC


Lattice determines

$$\Lambda = m_\rho, \quad d_B = \frac{m_{\rho_U} - m_{\rho_D}}{m_\rho}$$



Lattice prediction

- ◆ Form factor requires isospin breaking from ETC



Lattice determines


$$\Lambda = m_\rho, \quad d_B = \frac{m_{\rho_U} - m_{\rho_D}}{m_\rho}$$

- ◆ EM cross section with proton

$$\sigma_p^\gamma = \frac{\mu^2}{4\pi} \left(\frac{8\pi\alpha d_B}{\Lambda^2} \right)^2 \quad \mu = \frac{m_\phi m_N}{m_\phi + m_N} \quad |d_B| < 1 \quad m_\phi > m_p$$

Lattice prediction

- ◆ Form factor requires isospin breaking from ETC



Lattice determines

$$\Lambda = m_\rho, \quad d_B = \frac{m_{\rho_U} - m_{\rho_D}}{m_\rho}$$


- ◆ EM cross section with proton

$$\sigma_p^\gamma = \frac{\mu^2}{4\pi} \left(\frac{8\pi\alpha d_B}{\Lambda^2} \right)^2 \quad \mu = \frac{m_\phi m_N}{m_\phi + m_N} \quad |d_B| < 1 \quad m_\phi > m_p$$

$$\sigma_p^\gamma < 2.3 \times 10^{-44} \text{ cm}^2$$

First principle !

Composite Higgs ?




Composite Higgs ?

Basic interactions

$$\mathcal{L}_h = \frac{d_1}{\Lambda} h \partial_\mu \phi^* \partial^\mu \phi + \frac{d_2}{\Lambda} m_\phi^2 h \phi^* \phi$$

$$d_1 \approx d_2 \approx \mathcal{O}(1)$$




Composite Higgs ?

Basic interactions

$$\mathcal{L}_h = \frac{d_1}{\Lambda} h \partial_\mu \phi^* \partial^\mu \phi + \frac{d_2}{\Lambda} \boxed{m_\phi^2} h \phi^* \phi$$

$$d_1 \approx d_2 \approx \mathcal{O}(1)$$

DM is a GB



Composite Higgs ?

Basic interactions

$$\mathcal{L}_h = \frac{d_1}{\Lambda} h \partial_\mu \phi^* \partial^\mu \phi + \frac{d_2}{\Lambda} \boxed{m_\phi^2} h \phi^* \phi$$


$$d_1 \approx d_2 \approx \mathcal{O}(1)$$

DM is a GB

Cross section with the proton

$$\sigma_p = \frac{\mu^2}{4\pi} \left[\frac{(d_1 + d_2) f m_N m_\phi^2}{m_H^2 m_\phi v_{EW} \Lambda} + 8\pi\alpha \frac{d_B}{\Lambda^2} \right]^2$$

$$f \simeq 0.3$$




Composite Higgs ?

Basic interactions

$$\mathcal{L}_h = \frac{d_1}{\Lambda} h \partial_\mu \phi^* \partial^\mu \phi + \frac{d_2}{\Lambda} \boxed{m_\phi^2} h \phi^* \phi$$

$$d_1 \approx d_2 \approx \mathcal{O}(1)$$

DM is a GB



Cross section with the proton

$$\sigma_p = \frac{\mu^2}{4\pi} \left[\frac{(d_1 + d_2) f m_N m_\phi^2}{m_H^2 m_\phi v_{EW} \Lambda} + 8\pi\alpha \frac{d_B}{\Lambda^2} \right]^2$$

$$f \simeq 0.3$$

$$f_n = -\frac{(d_1 + d_2) f m_N m_\phi^2}{m_H^2 m_\phi v_{EW} \Lambda}$$

$$f_p = -\frac{(d_1 + d_2) f m_N m_\phi^2}{m_H^2 m_\phi v_{EW} \Lambda} - 8\pi\alpha \frac{d_B}{\Lambda^2}$$


Composite Higgs ?

Basic interactions

$$\mathcal{L}_h = \frac{d_1}{\Lambda} h \partial_\mu \phi^* \partial^\mu \phi + \frac{d_2}{\Lambda} \boxed{m_\phi^2} h \phi^* \phi$$

$$d_1 \approx d_2 \approx \mathcal{O}(1)$$

DM is a GB



Talks by Foadi and Kuti


Cross section with the proton

$$\sigma_p = \frac{\mu^2}{4\pi} \left[\frac{(d_1 + d_2) f m_N m_\phi^2}{m_H^2 m_\phi v_{EW} \Lambda} + 8\pi\alpha \frac{d_B}{\Lambda^2} \right]^2$$

$$f \simeq 0.3$$


$$f_n = -\frac{(d_1 + d_2) f m_N m_\phi^2}{m_H^2 m_\phi v_{EW} \Lambda}$$


$$f_p = -\frac{(d_1 + d_2) f m_N m_\phi^2}{m_H^2 m_\phi v_{EW} \Lambda} - 8\pi\alpha \frac{d_B}{\Lambda^2}$$



Experiments


Lattice





Experiments

Lattice




Summary

A natural avenue: Compositeness

Composite dark matter




Template for a composite Goldstone DM

First principle dark lattice simulations



Backup slides



LUX



Analysis by
 Del Nobile, Gelmini, Gondolo, Huh
 1311.4247


Taming volume effects

- ◆ 3 Volumes: $L^3 \times 32$
- ◆ Most chiral point for beta = 2.2



Approaching infinite volume

Form factors relations



$$C_{UD}^{(3)}(t_i, t, t_f, \vec{p}_i, \vec{p}_f) = T^U - T^D$$

$$C_{\overline{UD}}^{(3)}(t_i, t, t_f, \vec{p}_i, \vec{p}_f) = -T^U + T^D$$


$$C_{U\overline{D}}^{(3)}(t_i, t, t_f, \vec{p}_i, \vec{p}_f) = T^U + T^D$$

$$C_{\overline{UD}}^{(3)}(t_i, t, t_f, \vec{p}_i, \vec{p}_f) = -T^U - T^D$$

$$C_{\overline{UU}+\overline{DD}}^{(3)}(t_i, t, t_f, \vec{p}_i, \vec{p}_f) = 0$$

$$T^X = \sum_{\vec{x}_i, \vec{x}, \vec{x}_f} e^{-i(\vec{x}_f - \vec{x}) \cdot \vec{p}_f} e^{-i(\vec{x} - \vec{x}_i) \cdot \vec{p}_i} \left\langle 0 \left| \mathcal{O}_{UD}^{(\gamma_5)}(x_f) V_\mu^X(x) \mathcal{O}_{UD}^{(\gamma_5)\dagger}(x_i) \right| 0 \right\rangle$$

Continuum estimates



meson	β	m GeV	m^i GeV
vector	2.0	1750(10)	2060(12)
vector	2.2	1920(10)	2230(11)
vector	∞	2210(40)(290)(300)	2510(40)(280)(300)
axial vector	2.0	3280(50)	3850(50)
axial vector	2.2	3140(40)	3630(40)
axial vector	∞	2900(120)(230)(370)	3270(130)(370)(370)

- ◆ The estimates with an “i” account for the Z_a factor
- ◆ The table above is obtained taking the ratio of m over F_{π} first
- ◆ The table below are independent fits

meson	β	m GeV	m^i GeV
vector	2.0	1690(90)	1990(110)
vector	2.2	1860(60)	2120(70)
vector	∞	2160(270)(300)(270)	2430(300)(310)(300)
axial vector	2.0	3280(50)	3850(50)
axial vector	2.2	3140(40)	3630(40)
axial vector	∞	2900(120)(230)(370)	3270(130)(370)(370)