

Neutron-Antineutron Oscillation Matrix Elements on a Lattice

Michael Buchoff (University of Washington)

Chris Schroeder (Lawrence Livermore Natl. Lab)

Sergey Syritsyn (Brookhaven Natl. Lab)

Joe Wasem (Lawrence Livermore Natl. Lab)

*Lattice Meets Experiment 2013,
Brookhaven National Lab,
December 5 - 6, 2013*



Baryon Number Violation

- ✦ $\Delta B = 1$: Proton decay
- ✦ $\Delta B = 2$: Neutron-antineutron oscillation
(neutron/antineutron oscillation through $\Delta B = 1$ is possible, but suppressed)

Which one (if any) is realized in nature?

Searches for $n \rightarrow \bar{n}$ transitions

◆ In a nucleus

stability of a nucleus w.r.t $(n\bar{n})$ annihilation $T_d = R\tau_{n\bar{n}}^2$

iron atoms $T_d(Fe) > 7.2 \cdot 10^{31}$ year $\rightarrow \tau_{n\bar{n}} > 1.4 \cdot 10^8$ sec [Soudan 2]

oxygen atoms $T_d(O) > 1.77 \cdot 10^{32}$ year $\rightarrow \tau_{n\bar{n}} > 3.3 \cdot 10^8$ sec [Super-K]

◆ (Quasi) free neutrons

probability of event $P_{n \rightarrow \bar{n}} \sim \left(\frac{t}{\tau_{n\bar{n}}} \right)^2$

Reactor neutrons (Grenoble, 1990) $\tau_{n\bar{n}} > 0.86 \cdot 10^8$ sec

Proposed experiments:

- stored ultra-cold neutrons
- (focused, vertical) cold neutron beams

Neutron $\Leftrightarrow$ Antineutron Operators

Effective 6-quark operators *From Beyond (the Standard Model)* :
interaction with a massive Majorana lepton, unified theories, etc

[T.K.Kuo, S.T.Love, PRL45:93 (1980)]

[R.N.Mohapatra, R.E.Marshak, PRL44:1316 (1980)]

$$\mathcal{H}_{n\bar{n}} = \begin{pmatrix} E + V & \delta m \\ \delta m & E - V \end{pmatrix} \quad \tau_{n\bar{n}} = (2\delta m)^{-1}$$

$$\mathcal{L}_{\text{eff}} = \sum_i [c_i \mathcal{O}_i^{6q} + \text{h.c.}] \quad \delta m = -\langle \bar{n} | \int d^4x \mathcal{L}_{\text{eff}} | n \rangle = -\sum_i c_i \langle \bar{n} | \mathcal{O}_i^{6q} | n \rangle$$

Dimension-9 point-like operators suppressed by $(M_X)^{-5}$

*Sensitivity of matter to BN-violating terms is determined by
nuclear scale physics and non-perturbative QCD*

Neutron $\Leftrightarrow$ Antineutron Operators

Operators: pseudoscalar singlets w.r.t $SU(3)_c \otimes U(1)_{em} \left[\otimes SU(2)_L \right]$

$$\mathcal{O}_1 \chi_1 \{\chi_2 \chi_3\} = T_{ijklmn}^s [u_{\chi_1}^{iT} C u_{\chi_1}^j] [d_{\chi_2}^{kT} C d_{\chi_2}^l] [d_{\chi_3}^{mT} C d_{\chi_3}^n] \quad \chi_{1,2,3} = R, L$$

$$\mathcal{O}_2 \{\chi_1 \chi_2\} \chi_3 = T_{ijklmn}^s [u_{\chi_1}^{iT} C d_{\chi_1}^j] [u_{\chi_2}^{kT} C d_{\chi_2}^l] [d_{\chi_3}^{mT} C d_{\chi_3}^n]$$

$$\mathcal{O}_3 \{\chi_1 \chi_2\} \chi_3 = T_{ijklmn}^a [u_{\chi_1}^{iT} C d_{\chi_1}^j] [u_{\chi_2}^{kT} C d_{\chi_2}^l] [d_{\chi_3}^{mT} C d_{\chi_3}^n]$$

Computed using
MIT bag model

[S.Rao, R.Shrock, PLB116:238 (1982)]

Symmetry reduces # of operators: 24 $\rightarrow$ 14
(of which only 4 are $SU(2)_L$ symmetric)

$$3\mathcal{O}_3 \{\chi\chi\}\chi' = \mathcal{O}_2 \{\chi\chi\}\chi' - \mathcal{O}_1 \chi\{\chi\chi'\}$$

[W.Caswell, J.Milutinovic, G.Senjanovic, PLB122:373 (1983)]

On a lattice:

7 independent quantities
($R \leftrightarrow L$ symmetry)

$$\mathcal{O}_1 \chi\{\chi\chi\} \quad \mathcal{O}_1 \chi\{\chi\eta\} \quad \mathcal{O}_1 \chi\{\eta\eta\}$$


$$\mathcal{O}_2 \{\chi\chi\}\chi \quad \mathcal{O}_2 \{\chi\chi\}\eta \quad \mathcal{O}_2 \{\chi\eta\}\chi$$

$$\mathcal{O}_3 \{\chi\eta\}\chi \quad (\chi, \eta = R, L, \quad \chi \neq \eta)$$

$SU(2)_V$ and parity further reduce independent $\langle \bar{n} | \mathcal{O}_i^{6q} | n \rangle$ to 6 [B. Tiburzi]


Lattice Calculation

$$\langle N_{\uparrow}^{(+)}(t_2) \mathcal{O}^{6q}(0) N_{\downarrow}^{(-)}(-t_1) \rangle \underset{t_1, t_2, t_1+t_2 \rightarrow \infty}{\sim} e^{-M_n(t_2+t_1)} \langle n_{\uparrow} | \mathcal{O}^{6q} | \bar{n}_{\uparrow} \rangle$$



No quark-disconnected contractions!

Single propagator $\longrightarrow \forall t_1, t_2$



$$\begin{aligned} \langle n | \mathcal{O} | \bar{n} \rangle \Big|_{\text{lat}} &= \langle n | \mathcal{O} | \bar{n} \rangle \\ &+ O(e^{-\Delta E_{\text{exc}} t_1}, e^{-\Delta E_{\text{exc}} t_2}, e^{-\Delta E_{\text{exc}}(t_1+t_2)}) \end{aligned}$$

Complete set of correlators for sophisticated exc.state analysis:

- Exponential fits
- Variational (GPoF)

Preliminary Results

- ◆ $20^3 \times 256$ anisotropic lattice, clover-improved Wilson fermions
- ◆ pion mass 390 MeV
- ◆ lattice spacing $a = 0.123 \text{ fm}$

[Hadron Spectrum collaboration]


- ◆ no renormalization yet, only *bare* lattice quantities
- ◆ 886 gauge configurations x 9 samples per lattice

PRELIMINARY ANALYSIS !

Simple ratios to inspect signal for matrix elements





$$\langle n | \mathcal{O} | \bar{n} \rangle \sim \frac{C_{n\mathcal{O}\bar{n}}(t_2, 0, -t_1)}{\sqrt{C_{nn}(t_2, 0)C_{\bar{n}\bar{n}}(0, -t_1)}}$$

Effective Mass: Gauging Excited States




$$m_{eff}(t) = \log \frac{C_2(t)}{C_2(t+1)}$$

Bare Matrix Elements on a Lattice



Comparison to MIT Bag Model Calculation



(values normalized by $\mathcal{O}_{1\chi\{\chi\chi\}}$)

MIT Bag Model calc.: [S.Rao, R.Shrock, PLB116:238 (1982)]

- Operators have different anomalous dimensions
- Some operators mix
- Chiral symmetry breaking effects?

Next Step: Renormalization

Non-perturbative lattice renormalization: bare quantities $\longrightarrow$ MOM-scheme

Perturbative 1-loop scaling is known [\[W.Caswell et al PLB122:373 \(1983\)\]](#)

- need higher loop calculations?
- matching coefficients for conversion to MSbar?

In the chiral limit, mixing between operators is restricted by flavor $SU(2)_{L,R}$ symmetry:

$\mathcal{O}_{L(LL)}^1 + 4\mathcal{O}_{(LL)L}^2$	$(\mathbf{3}, -1)_L \otimes (\mathbf{0}, 0)_R$	$(\alpha_S/4\pi)(-12)$
$\mathcal{O}_{(LL)L}^2 - \mathcal{O}_{L(LL)}^1$	$(\mathbf{1}, -1)_L \otimes (\mathbf{0}, 0)_R$	$(\alpha_S/4\pi)(-2)$
$\mathcal{O}_{(LL)R}^2 - \mathcal{O}_{L(LR)}^1$	$(\mathbf{0}, 0)_L \otimes (\mathbf{1}, -1)_R$	0
$\mathcal{O}_{(LR)L}^3$	$(\mathbf{1}, -1)_L \otimes (\mathbf{0}, 0)_R$	$(\alpha_S/4\pi)(+2)$
$\mathcal{O}_{L(LR)}^1 + 2\mathcal{O}_{(LL)R}^2$	$(\mathbf{2}, 0)_L \otimes (\mathbf{1}, -1)_R$	$(\alpha_S/4\pi)(-6)$
$\mathcal{O}_{(LR)L}^2$	$(\mathbf{2}, -1)_L \otimes (\mathbf{1}, 0)_R$	$(\alpha_S/4\pi)(-6)$
$\mathcal{O}_{R(LL)}^1$	$(\mathbf{2}, -2)_L \otimes (\mathbf{1}, +1)_R$	$(\alpha_S/4\pi)(-6)$

(+ $L \leftrightarrow R$ counterparts)

Lattice: chiral symmetry is broken, renormalize $(L \pm R)$ multiplets instead
or switch to chiral (domain wall) fermions

Summary & Outlook

- ◆ Clear lattice signal for non-zero $\langle n | \mathcal{O}^{6q} | \bar{n} \rangle$ with modest statistics
- ◆ Qualitative agreement with the MIT Bag Model is reassuring
- ◆ Expect only moderate renormalization and mixing effects

Outlook

- ◆ Non-perturbative renormalization
- ◆ Improve analysis to extract ground state M.E.
- ◆ Additional volumes, pion masses, possibly with chiral quarks