

Energy-Energy Correlators (EEC) in DIS

a novel probe to TMDPDFs

Yiannis Makris
INFN (Pavia)




Supported by the FELLINI project and the European Commission



[arXiv:2102.05669](https://arxiv.org/abs/2102.05669) (H. Li, Y. Makris, I. Vitev)


Motivation

jet-TMDs




Perturbative effects dominate the final state contribution gives cleaner access to TMDPDF Introduces jet-clustering effects into the non-perturbative corrections (NGLs)

SIDIS




Strong final state effects from fragmentations functions. Couples FF and PDF effects in a non-trivial way

EEC_{DIS}



Every particle in the event is considered but with its weight is proportional to the “energy fraction”: $z_i = p_i \cdot P / q \cdot P$

In this talk



EEC in lepton colliders

$$\text{EEC} = \sum_{a,b} \int \frac{d\sigma(e^+e^- \rightarrow a+b+X)}{\sigma} \frac{2E_a E_b}{s} \delta(\cos \theta_{ab} - \cos(\chi))$$

PRL. 41 (1978) 1585 (C. L. Basham, L. S. Brown, S. D. Ellis, and S.T. Love)

Soft radiation (direct) contribution
is suppressed. Recoil from soft radiation
needs to be included.




Figure from arXiv:1801.02627 (I. Moulton and H.X. Zhu)

EEC in lepton colliders

$$\text{EEC} = \sum_{a,b} \int \frac{d\sigma(e^+e^- \rightarrow a + b + X)}{\sigma} \frac{2E_a E_b}{s} \delta(\cos \theta_{ab} - \cos(\chi))$$

PRL. 41 (1978) 1585 (C. L. Basham, L. S. Brown, S. D. Ellis, and S.T. Love)

Soft radiation (direct) contribution
is suppressed. Recoil from soft radiation
needs to be included.

EEC measurements/extractions of α_s

PLUTO, Phys. Lett. B 99, 292 (1981)

MARKII, Phys. Rev. Lett. 49, 521 (1982)

MARKJ, Phys. Rev. Lett. 50, 2051 (1983)

CELLO, Phys. Lett. B 138, 311 (1984)

JADE, Z. Phys. C 25, 231 (1984)

MAC, Phys. Rev. D 31, 2724 (1985)

TASSO, Z. Phys. C 36, 349 (1987)

TOPAZ, Phys. Lett. B 227, 495 (1989)

DELPHI, Phys. Lett. B 252, 149 (1990)

OPAL, Phys. Lett. B 252, 159 (1990)

ALEPH, Phys. Lett. B 257, 479 (1991)

L3, Phys. Lett. B 257, 469 (1991)

SLD, Phys. Rev. D 51, 962 (1995)

Transverse-EEC for hadronic colliders

$$\text{TEEC} = \sum_{a,b} \int \frac{d\sigma(pp \rightarrow a + b + X)}{\sigma} \frac{2E_{T,a}E_{T,b}}{(E_T^{\text{total}})^2} \delta(\cos \phi_{ab} - \cos \phi)$$

Phys. Lett. 141B, 447 (1984) (A. Ali, E. Pietarinen, and W. J. Stirling)

Soft and Beam (direct) contribution is suppressed. Recoil from soft radiation and ISR needs to be included.






Figure from arXiv:1901.04497 (A. Gao, H.T. Li, I. Moult, and H.X. Zhu)

Transverse-EEC for hadronic colliders

$$\text{TEEC} = \sum_{a,b} \int \frac{d\sigma(pp \rightarrow a + b + X)}{\sigma} \frac{2E_{T,a}E_{T,b}}{(E_T^{\text{total}})^2} \delta(\cos \phi_{ab} - \cos \phi)$$

arXiv: 1508.01579 (ATLAS)


arXiv: 1707.02562 (ATLAS)



Adaptations of TEEC in DIS


arXiv: 2008.00271 (A. Ali, G. Li, W. Wang, Z-P. Xing)

Breit Frame (end-point: dijets)



arXiv: 2006.02437 (H.T. Li, I. Vitev, Y.J. Zhu)

Laboratory Frame (single jet)



The definition is the same as before:

$$\text{TEEC} = \sum_{a,b} \int \frac{d\sigma(ep \rightarrow a + b + X)}{\sigma} \frac{2E_{T,a}E_{T,b}}{(E_T^{\text{total}})^2} \delta(\cos \phi_{ab} - \cos \phi)$$

Defined as lepton-hadron correlation:


$$\text{TEEC}_{\text{Lab.}} = \sum_a \int \frac{d\sigma(ep \rightarrow e + a + X)}{\sigma} \frac{2E_{T,l}E_{T,a}}{E_{T,l}E_T^{\text{hadr.}}} \delta(\cos \phi_{ab} - \cos \phi)$$

An EEC definition for DIS



$$\text{EEC}_{\text{DIS}} = \sum_a \int \frac{d\sigma_{ep \rightarrow e+a+X}}{\sigma} z_a \delta(\cos \theta_{ap} - \cos \theta)$$

$z_a \equiv \frac{P \cdot p_a}{P \cdot q} \Big|_{\text{B.F.}} \simeq \frac{p_a^+}{Q}$
 Soft and anti-collinear radiation is suppressed (also true in $\text{TEEC}_{\text{Lab.}}$)

Geometrical separation of: **collinear**, **soft**, and **anti-collinear**



Insensitive to (pseudo)rapidity constraints



EEC as a probe to TMDs

$$\text{EEC} = \sum_{a,b} \int \frac{d\sigma(e^+e^- \rightarrow a+b+X)}{\sigma} \frac{2E_a E_b}{s} \delta(\cos \theta_{ab} - \cos(\chi))$$

back-to-back limit:
TMD observable




EEC(back-to-back)

~

weighted TMD factorization



SLD, Phys. Rev. D 51, 962 (1995)



$$\text{EEC}_{\text{DIS}} = \sum_a \int \frac{d\sigma_{ep \rightarrow e+a+X}}{\sigma} z_a \delta(\cos \theta_{ap} - \cos \theta)$$

Connection to universal TMDs (unpolarized case)

$$\cos \theta \sim -1$$

$$\text{EEC}_{\text{DIS}} = \sigma_0(Q, x, y) H_{ij}(Q, x, \mu) \int d\mathbf{q}_T \delta\left(\cos \theta - \frac{2|\mathbf{q}_T|^2}{Q^2} + 1\right) \int \frac{d\mathbf{b}}{(2\pi)^2} \exp(-i\mathbf{b} \cdot \mathbf{q}_T) f_{1, j \leftarrow P}^{\text{sub.}}(x, b, \mu, \nu) J_i^{\text{sub.}}(b, \mu, \nu)$$


* universal = back-to-back
four Wilson-line soft function
incorporated into the definition.

TEEC case: **Not** back-to-back

Does not involve the variable
 z_h included TMD studies in SIDIS

TMD PDF

The **universal*** TMD
parton distribution
function, unpolarized
in this case.

For Trans. pol. beam
modification:
replace with **Sivers**.

EEC Jet function

The same jet function that appears
in the original EEC observable. Can
be written as the weighed integral of
the **universal*** TMD fragmentation
functions.

$$J_i(b, \mu, \nu) = \sum_a \int_0^1 dz \, z \, D_{i \rightarrow a}(z, b, \mu, \nu)$$

EEC jet function: hadronization effects

$$J_i(b, \mu, \nu) = \sum_a \int_0^1 dz \, z \, D_{i \rightarrow a}(z, b, \mu, \nu)$$

perturbative matching
onto collinear-FFs



$$D_{i \rightarrow a}(z, b, \mu, \nu) = \sum_j \int_z^1 \frac{du}{u} \mathcal{I}_{ij} \left(b, \frac{z}{u}, \mu, \nu \right) d_{j \rightarrow a}(u, \mu) + \mathcal{O} \left(\Lambda_{\text{QCD}}^2 b^2, \frac{\Lambda_{\text{QCD}}^2}{Q^2} \right)$$

$$J_i^{\text{OPE}}(b, \mu, \nu) = \sum_j \int_0^1 dz \, z \, \mathcal{I}_{ij}(b, z, \mu, \nu)$$




Sum rules for FF

$N^3LL + \mathcal{O}(\alpha_s^2)$ predictions

$$\cos \theta \sim -1$$

$$EEC_{DIS} = \sigma_0(Q, x, y) H_{ij}(Q, x, \mu) \int d\mathbf{q}_T \delta\left(\cos \theta - \frac{2|\mathbf{q}_T|^2}{Q^2} + 1\right) \int \frac{d\mathbf{b}}{(2\pi)^2} \exp(-i\mathbf{b} \cdot \mathbf{q}_T) f_{1,j \leftarrow P}^{\text{sub.}}(x, b, \mu, \nu) J_i^{\text{sub.}}(b, \mu, \nu)$$



EEC jet function: hadronization effects

$$J_i(b, \mu, \nu) = \sum_a \int_0^1 dz \, z \, D_{i \rightarrow a}(z, b, \mu, \nu)$$

perturbative matching
onto collinear-FFs



$$D_{i \rightarrow a}(z, b, \mu, \nu) = \sum_j \int_z^1 \frac{du}{u} \mathcal{I}_{ij} \left(b, \frac{z}{u}, \mu, \nu \right) d_{j \rightarrow a}(u, \mu) + \mathcal{O} \left(\Lambda_{\text{QCD}}^2 b^2, \frac{\Lambda_{\text{QCD}}^2}{Q^2} \right)$$

$$J_i^{\text{OPE}}(b, \mu, \nu) = \sum_j \int_0^1 dz \, z \, \mathcal{I}_{ij}(b, z, \mu, \nu)$$



Sum rules for FF

Ansatz used in extractions of TMD-FF

$$\sqrt{S} D_{i/a}(z, b; \mu_0, \nu_0) = \sqrt{S^{\text{pert.}}} D_{i/a}^{\text{OPE}}(z, b; \mu_0, \nu_0) D_{i/a}^{\text{NP}}(z, b)$$

$$\sqrt{S} J_i(b, \mu_0, \nu_0) = \sqrt{S^{\text{pert.}}} J_i^{\text{OPE}}(b, \mu_0, \nu_0) j_i(b)$$

Challenges


Prone to experimental uncertainties due to the inclusion of neutral particles

Handle over flavor separation

$$j_i(b) = \left[J_i^{\text{OPE}}(b, \mu_0, \nu_0) \right]^{-1} \sum_a \int_0^1 du dy (uy) \, \mathcal{I}_{ij} \left(\frac{b}{u}, u; \mu_0, \nu_0 \right) d_{i/a}(y, \mu_0) D_{i/a}^{\text{NP}}(uy, b)$$

EEC jet function: hadronization effects



$$j_i(b) = \left[J_i^{\text{OPE}}(b, \mu_0, \nu_0) \right]^{-1} \sum_a \int_0^1 du dy (uy) \mathcal{I}_{ij} \left(\frac{b}{u}, u; \mu_0, \nu_0 \right) d_{i/a}(y, \mu_0) D_{i/a}^{\text{NP}}(uy, b)$$



arXiv:0707.1506 (D. de Florian, R. Sassot, M. Stratmann)
arXiv:hep-ph/0703242 (D. de Florian, R. Sassot, M. Stratmann)

$$D_{i/a}^{\text{NP}}(y, b) = \exp \left(-0.042 \frac{b^2}{y^2} \right)$$

arXiv:1406.3073 (P. Sun, J. Isaacson, C.-P. Yuan, F. Yuan)
arXiv:1505.05588 (A. Prokudin, P. Sun, F. Yuan)




+
LO perturbative
components


Best Fit: $j_i(b) = \exp \left(-0.59b - 0.03b^2 \right)$

EEC jet function: hadronization effects

Effect of total NP
(TMDPDF + EECjet + Rapidity AD)





Decomposition of NP elements



$$j_i(b) = \exp\left(-0.59b - 0.03b^2\right)$$

$N^3\text{LL} + \mathcal{O}(\alpha_s^2)$ predictions



Charged particles and flavor separation: $\sum_a \rightarrow \sum_{a \in \mathbb{S}}$

$$J_i(b, \mu, \nu) \rightarrow \mathcal{G}_{i \rightarrow \mathbb{S}}(b, \mu, \nu) = \sum_{a \in \mathbb{S}} \int_0^1 dz \, z D_{i/a}(z, \mathbf{b}; \mu, \nu)$$

perturbative matching
onto collinear-FFs



$$\mathcal{G}_{i \rightarrow \mathbb{S}}^{\text{OPE}}(b, \mu, \nu) = \sum_j \mathcal{P}_{j \rightarrow \mathbb{S}} \int_0^1 dz \, z \mathcal{I}_{ij}(b, z, \mu, \nu)$$

Sum rules for FF



$$D_{i \rightarrow a}(z, b, \mu, \nu) = \sum_j \int_z^1 \frac{du}{u} \mathcal{I}_{ij}\left(b, \frac{z}{u}, \mu, \nu\right) d_{j \rightarrow a}(u, \mu) + \mathcal{O}\left(\Lambda_{\text{QCD}}^2 b^2, \frac{\Lambda_{\text{QCD}}^2}{Q^2}\right)$$

Ansatz used in extractions of TMD-FF

$$\sqrt{S} \mathcal{G}_{i \rightarrow \mathbb{S}}(\mathbf{b}, \mu, \nu) = \sqrt{S^{\text{pert}}} R(\mathbf{b}, \mu, \nu) \mathcal{G}_{i \rightarrow \mathbb{S}}^{\text{OPE}}(\mathbf{b}, \mu_0, \nu_0) g_{i \rightarrow \mathbb{S}}(\mathbf{b})$$

$$\mathcal{G}_{i \rightarrow \mathbb{S}}^{\text{OPE}}(\mathbf{b}, \mu, \nu) = \sum_j \mathcal{F}_{j \rightarrow \mathbb{S}}(\mu) \int_0^1 dw \, w \mathcal{I}_{ij}\left(\frac{\mathbf{b}}{w}, w; \mu, \nu\right)$$

Interesting cases

$\mathbb{S} = \text{all charge hadrons}$

$\mathbb{S} = h$

$$g_{i \rightarrow \mathbb{S}}(\mathbf{b}) = \left[\mathcal{G}_{i \rightarrow \mathbb{S}}^{\text{OPE}}(\mathbf{b}, \mu_0, \nu_0) \right]^{-1} \sum_{a \in \mathbb{S}} \int_0^1 dz \, z D_{i/a}(z, \mathbf{b}; \mu_0, \nu_0)$$

Summary and Outlook

Introduce definition of EEC in DIS

Provide a factorized cross-section in terms of TMD elements

Obtain $N^3LL + \mathcal{O}(\alpha_s^2)$ predictions

Evaluate NP component of EEC jet function from past TMDFF extractions

Expand the definition to charge particles only!

Data on EEC in e^+e^- could be utilized for sim. extraction of rapidity AD EEC jet function and strong coupling

Spin effects: Sivers!

Break flavor degeneracy with FSR particle selection