

Dark Interactions:

PERSPECTIVES FROM THEORY AND EXPERIMENT

June 11-13, 2014 • Brookhaven National Laboratory

Dark Matter in ATLAS SUSY searches

Estel Perez Codina (TRIUMF)

On behalf of the ATLAS Collaboration





Introduction

- Overwhelming evidence for dark matter (DM)
- In SUSY models with RPC, the lightest supersymmetric particle (**LSP**) is an **excellent WIMP candidate**
- DM at collider experiments [ATLAS]:
 - **Escapes detector** $\Rightarrow$ imbalance of visible momentum (E_T^{miss})
 - Favorite SUSY DM candidates:
 - Neutralino
 - Gravitino
 - **Direct production of LSPs** not accessible due to **low cross-section**
 - In SUSY the LSP is typically produced in the **decay chain of heavier SUSY particles**

Outline


- SUSY production
- Analysis strategy
- Neutralino LSP results (*)
- Gravitino LSP results (*)
- Summary & Outlook

*Focus on latest SUSY results relevant to dark matter



Signature: $E_T^{\text{miss}} + \text{SM particles}$ 2


SUSY Production



<https://twiki.cern.ch/twiki/bin/view/LHCPhysics/SUSYCrossSections>

arXiv:1206.2892

- SUSY searches strategy driven by cross section and luminosity
- Inclusive searches:
 - Strong (gluino and 1st&2nd gen. squark) production
- Exclusive searches:
 - Strong (3rd gen. squark) production
 - Weak production (Direct gaugino / sleptons)




Constrains on the LSP mass are relative to the spectrum considered (masses of strongly/weakly produced particles)

Analysis strategy

SUSY searches rely on understanding the SM backgrounds

Strategy for a “typical” SUSY analysis:



Large number of SR optimized for each scenario



Account for potential signal contamination in control regions when setting limits

Neutralino LSP

- The super-partners of:
 - gauge bosons (**winos**, **bino**)
 - higgs boson (**higgsinos**)


mix to give **charginos and neutralinos**: $\tilde{\chi}^{+/-}, \tilde{\chi}^0$.

- Lightest Neutralino ($\tilde{\chi}_1^0$) likely to be the LSP
DM candidate $\Rightarrow$ **Constrains on its composition from observed relic density**



Well-tempered

Z- and h-funnel




Slepton-coannihilation or A-funnel

Each point represents a pMSSM model (fixed values of the mass parameters) surviving the DM constraints

In the following results, the LSP is the **bino-like** $\tilde{\chi}_1^0$, unless otherwise stated

Strong production (gluinos, 1st/2nd gen. squarks)



$\tilde{q}\tilde{q}, \tilde{g}\tilde{g}, \tilde{q}\tilde{g}$ production



0-lepton + 2-6 jets

Use m_{eff} as discriminant

$E_T^{\text{miss}}/m_{\text{eff}}$ reduce multijets bkg.




Strong production (stop)

$\Delta m(\tilde{t}, \tilde{\chi}_1^0)$ small
 $\Rightarrow$ b not accessible


$\tilde{t} \rightarrow c \tilde{\chi}_1^0$

ISR + charm tagged jet



ATLAS-CONF-2013-068

Several analysis target
stop pair production:
 two-step or one-step
 decay



Status, Moriond 2014

Binary

SS

3.4171]


- 0L, $\tilde{t}_1 \rightarrow t \tilde{\chi}_1^0$
- 1L, $\tilde{t}_1 \rightarrow t \tilde{\chi}_1^0$
- 2L, $\tilde{t}_1 \rightarrow t \tilde{\chi}_1^0$
- 2L, $\tilde{t}_1 \rightarrow W b \tilde{\chi}_1^0$
- 0L, mono-jet/c-tag, $\tilde{t}_1 \rightarrow c \tilde{\chi}_1^0$
- 0L, $m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_1^0} + 5 \text{ GeV}$
- 1-2L, $\tilde{t}_1 \rightarrow b \tilde{\chi}_1^\pm, m_{\tilde{\chi}_1^\pm} = 106 \text{ GeV}$
- 1L, $\tilde{t}_1 \rightarrow b \tilde{\chi}_1^\pm, m_{\tilde{\chi}_1^\pm} = 150 \text{ GeV}$
- 2L, $\tilde{t}_1 \rightarrow b \tilde{\chi}_1^\pm, m_{\tilde{\chi}_1^\pm} = m_{\tilde{t}_1} - 10 \text{ GeV}$
- 1-2L, $\tilde{t}_1 \rightarrow b \tilde{\chi}_1^\pm, m_{\tilde{\chi}_1^\pm} = 2 \times m_{\tilde{\chi}_1^0}$

$L_{\text{int}} = 20 - 21 \text{ fb}^{-1}$ $\sqrt{s} = 8 \text{ TeV}$

0L ATLAS-CONF-2013-024
 1L ATLAS-CONF-2013-037
 2L [1403.4853]
 2L [1403.4853]
 0L mono-jet/c-tag, CONF-2013-068
 0L [1308.2631]
 2L [1403.4853]
 1L CONF-2013-037, 0L [1308.2631]
 2L [1403.4853]
 1L CONF-2013-037, 2L [1403.4853]

$L_{\text{int}} = 4.7 \text{ fb}^{-1}$ $\sqrt{s} = 7 \text{ TeV}$

0L [1208.1447]
 1L [1208.2590]
 2L [1209.4186]
 -
 -
 2L [1208.4305], 1-2L [1209.2102]
 -
 -
 1-2L [1209.2102]



$m_{\tilde{t}_1} [\text{GeV}]$

Electroweak production


Cover as many final states as possible

Exploiting Higgs in decay chain for the first time!

Multileptons: Use m_{T2} , m_{ll} , lepton charge and flavor


3lep, 2lep, 2tau, 1lep+b $\bar{b}$

ATLAS Preliminary 20.3-20.7 fb $^{-1}$, $\sqrt{s}=8$ TeV Status: Moriond 2014




$\tilde{\chi}_1^+ \tilde{\chi}_1^-$ production

2L




$\tilde{\chi}^{+/-}_1 \tilde{\chi}^0_2$ production


2L



3L



3L



Compressed scenarios




Electroweak production

Intermediate **right-handed sleptons**

$\Rightarrow \tilde{\chi}_1^+$ has to decay via W (lower BR)

production of higgsino-like $\tilde{\chi}_2^0 \tilde{\chi}_3^0$ interesting

4lep + MET , Z veto





Electroweak production (pMSSM)

Phenomenological model: **pMSSM** (with $\tilde{\chi}_1^0$ LSP)

Composition of electroweakinos controlled by **wino (M2)** and **higgsino (μ)** mass parameters

Bino mass **M1=50 GeV** $\Rightarrow$ Z “funnel” $\Rightarrow$ high annihilation rate $\Rightarrow$ can **satisfy DM relic abundance** constraints



Electroweak production (AMSB)




AMBS model: $\tilde{\chi}_1^0, \tilde{\chi}_1^{+/-}$ wino-like co-LSPs $\Rightarrow$ degenerate

$\Delta m < 1 \text{ GeV} \Rightarrow \tilde{\chi}_1^{+/-} \rightarrow \tilde{\chi}_1^0 \pi^{+/-}$

$\Rightarrow \tilde{\chi}_1^{+/-}$ long-lived $\Rightarrow (0.1 - 10 \text{ ns})$ decay in tracker

Disappearing track

$\tilde{\chi}_1^0 \tilde{\chi}_1^{+/-}$ or $\tilde{\chi}_1^+ \tilde{\chi}_1^-$ production (+ ISR jet)




Gravitino LSP

- **GMSB/GGM** scenarios:
 - **Gravitino** ($\tilde{G}$, supersymmetric partner of the graviton) **is always the LSP**
 - **very light** ($< \text{keV}$) $\rightarrow$ harder spectrum (w.r.t. $\tilde{\chi}_1^0$ LSP case)
 - **Phenomenology** driven by the nature of the **NLSP**:
 - Typically $\tilde{\chi}_1^0$, staus, ...
 - NLSP $\rightarrow \tilde{G} + \text{SM particle}$

Stau NLSP

Stop pair production





Squark / gluino pair production

NLSP mostly stau at high $\tan\beta$
(hadronic tau decay)

2 taus




Uses m_T , H_T , m_{eff} as discriminants



$\tilde{\chi}_1^0$ NLSP

$$\tilde{\chi}_1^0 \text{ bino-like} \Rightarrow \tilde{\chi}_1^0 \rightarrow \gamma \tilde{G}$$

2 photons




$$\tilde{\chi}_1^0 \text{ higgsino-like} \Rightarrow \tilde{\chi}_1^0 \rightarrow Z/h \tilde{G}$$


Stop pair production

$$\tilde{t} \rightarrow b \tilde{\chi}_1^{\pm} \text{ or } \tilde{t} \rightarrow t \tilde{\chi}_{1/2}^0$$

Z+b-jets



4lep (Z)




Squark/gluino NLSP

Direct production of $\tilde{G} + \tilde{q}/\tilde{g}$


Decay $\tilde{q}/\tilde{g} \rightarrow \tilde{G} + q/g$

$\Rightarrow$ **Mono-jet** signature

Probe **gravitino mass**



Analysis covered by Steven Schramm in his talk "Dark Matter Searches at ATLAS"



$m(\tilde{q}/\tilde{g}) < 1250 \text{ GeV}$

$\Rightarrow m(\tilde{G}) < 10^{-4} \text{ eV}$ excluded

ATLAS SUSY Searches* - 95% CL Lower Limits

Status: Moriond 2014

ATLAS Preliminary

$$\int \mathcal{L} dt = (4.6 - 22.9) \text{ fb}^{-1} \quad \sqrt{s} = 7, 8 \text{ TeV}$$

Model	e, μ, τ, γ	Jets	E_T^{miss}	$\int \mathcal{L} dt [\text{fb}^{-1}]$	Mass limit	Reference
Inclusive Searches	MSUGRA/CMSSM	0	2-6 jets	Yes	20.3	$\tilde{q}, \tilde{g}$ 1.7 TeV $m(\tilde{q})=m(\tilde{g})$ ATLAS-CONF-2013-047
	MSUGRA/CMSSM	1 e, μ	3-6 jets	Yes	20.3	$\tilde{g}$ 1.2 TeV any $m(\tilde{q})$ ATLAS-CONF-2013-062
	MSUGRA/CMSSM	0	7-10 jets	Yes	20.3	$\tilde{g}$ 1.1 TeV any $m(\tilde{q})$ 1308.1841
	$\tilde{q}\tilde{q}, \tilde{q} \rightarrow q\tilde{\chi}_1^0$	0	2-6 jets	Yes	20.3	$\tilde{q}$ 740 GeV $m(\tilde{\chi}_1^0)=0 \text{ GeV}$ ATLAS-CONF-2013-047
	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow q\tilde{q}\tilde{\chi}_1^0$	0	2-6 jets	Yes	20.3	$\tilde{g}$ 1.3 TeV $m(\tilde{\chi}_1^0)=0 \text{ GeV}$ ATLAS-CONF-2013-047
	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow q\tilde{q}\tilde{\chi}_1^{\pm} \rightarrow q\tilde{q}W^{\pm}\tilde{\chi}_1^0$	1 e, μ	3-6 jets	Yes	20.3	$\tilde{g}$ 1.18 TeV $m(\tilde{\chi}_1^0)<200 \text{ GeV}, m(\tilde{\chi}^{\pm})=0.5(m(\tilde{\chi}_1^0)+m(\tilde{g}))$ ATLAS-CONF-2013-062
	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow q\tilde{q}(\ell\ell/\ell\nu/\nu\nu)\tilde{\chi}_1^0$	2 e, μ	0-3 jets	-	20.3	$\tilde{g}$ 1.12 TeV $m(\tilde{\chi}_1^0)=0 \text{ GeV}$ ATLAS-CONF-2013-089
	GMSB ($\tilde{\ell}$ NLSP)	2 e, μ	2-4 jets	Yes	4.7	$\tilde{g}$ 1.24 TeV $\tan\beta<15$ 1208.4688
	GMSB ($\tilde{\ell}$ NLSP)	1-2 τ	0-2 jets	Yes	20.7	$\tilde{g}$ 1.4 TeV $\tan\beta>18$ ATLAS-CONF-2013-026
	GGM (bino NLSP)	2 γ	-	Yes	20.3	$\tilde{g}$ 1.28 TeV $m(\tilde{\chi}_1^0)>50 \text{ GeV}$ ATLAS-CONF-2014-001
3 rd gen. $\tilde{g}$ med.	GGM (wino NLSP)	1 $e, \mu + \gamma$	-	Yes	4.8	$\tilde{g}$ 619 GeV $m(\tilde{\chi}_1^0)>50 \text{ GeV}$ ATLAS-CONF-2012-144
	GGM (higgsino-bino NLSP)	γ	1 b	Yes	4.8	$\tilde{g}$ 900 GeV $m(\tilde{\chi}_1^0)>220 \text{ GeV}$ 1211.1167
	GGM (higgsino NLSP)	2 e, μ (Z)	0-3 jets	Yes	5.8	$\tilde{g}$ 690 GeV $m(\tilde{H})>200 \text{ GeV}$ ATLAS-CONF-2012-152
	Gravitino LSP	0	mono-jet	Yes	10.5	$\tilde{F}^{1/2}$ scale 645 GeV $m(\tilde{g})>10^{-4} \text{ eV}$ ATLAS-CONF-2012-147
	$\tilde{g} \rightarrow b\tilde{b}\tilde{\chi}_1^0$	0	3 b	Yes	20.1	$\tilde{g}$ 1.2 TeV $m(\tilde{\chi}_1^0)<600 \text{ GeV}$ ATLAS-CONF-2013-061
	$\tilde{g} \rightarrow t\tilde{t}\tilde{\chi}_1^0$	0	7-10 jets	Yes	20.3	$\tilde{g}$ 1.1 TeV $m(\tilde{\chi}_1^0)<350 \text{ GeV}$ 1308.1841
	$\tilde{g} \rightarrow t\tilde{t}\tilde{\chi}_1^{\pm}$	0-1 e, μ	3 b	Yes	20.1	$\tilde{g}$ 1.34 TeV $m(\tilde{\chi}_1^0)<400 \text{ GeV}$ ATLAS-CONF-2013-061
	$\tilde{g} \rightarrow b\tilde{t}\tilde{\chi}_1^{\pm}$	0-1 e, μ	3 b	Yes	20.1	$\tilde{g}$ 1.3 TeV $m(\tilde{\chi}_1^0)<300 \text{ GeV}$ ATLAS-CONF-2013-061
	$\tilde{b}_1\tilde{b}_1, \tilde{b}_1 \rightarrow b\tilde{\chi}_1^0$	0	2 b	Yes	20.1	$\tilde{b}_1$ 100-620 GeV $m(\tilde{\chi}_1^0)<90 \text{ GeV}$ 1308.2631
	$\tilde{b}_1\tilde{b}_1, \tilde{b}_1 \rightarrow t\tilde{\chi}_1^{\pm}$	2 e, μ (SS)	0-3 b	Yes	20.7	$\tilde{b}_1$ 275-430 GeV $m(\tilde{\chi}_1^0)=2 m(\tilde{\chi}_1^0)$ ATLAS-CONF-2013-007
3 rd gen. squarks direct production	$\tilde{t}_1\tilde{t}_1$ (light), $\tilde{t}_1 \rightarrow b\tilde{\chi}_1^{\pm}$	1-2 e, μ	1-2 b	Yes	4.7	$\tilde{t}_1$ 110-167 GeV $m(\tilde{\chi}_1^0)=55 \text{ GeV}$ 1208.4305, 1209.2102
	$\tilde{t}_1\tilde{t}_1$ (light), $\tilde{t}_1 \rightarrow Wb\tilde{\chi}_1^0$	2 e, μ	0-2 jets	Yes	20.3	$\tilde{t}_1$ 130-210 GeV $m(\tilde{\chi}_1^0)=m(\tilde{t}_1)-m(W)-50 \text{ GeV}, m(\tilde{t}_1)<m(\tilde{\chi}_1^0)$ 1403.4853
	$\tilde{t}_1\tilde{t}_1$ (medium), $\tilde{t}_1 \rightarrow t\tilde{\chi}_1^0$	2 e, μ	2 jets	Yes	20.3	$\tilde{t}_1$ 215-530 GeV $m(\tilde{\chi}_1^0)=1 \text{ GeV}$ 1403.4853
	$\tilde{t}_1\tilde{t}_1$ (medium), $\tilde{t}_1 \rightarrow b\tilde{\chi}_1^{\pm}$	0	2 b	Yes	20.1	$\tilde{t}_1$ 150-580 GeV $m(\tilde{\chi}_1^0)<200 \text{ GeV}, m(\tilde{\chi}_1^{\pm})-m(\tilde{\chi}_1^0)=5 \text{ GeV}$ 1308.2631
	$\tilde{t}_1\tilde{t}_1$ (heavy), $\tilde{t}_1 \rightarrow t\tilde{\chi}_1^0$	1 e, μ	1 b	Yes	20.7	$\tilde{t}_1$ 200-610 GeV $m(\tilde{\chi}_1^0)=0 \text{ GeV}$ ATLAS-CONF-2013-037
	$\tilde{t}_1\tilde{t}_1$ (heavy), $\tilde{t}_1 \rightarrow t\tilde{\chi}_1^{\pm}$	0	2 b	Yes	20.5	$\tilde{t}_1$ 320-660 GeV $m(\tilde{\chi}_1^0)=0 \text{ GeV}$ ATLAS-CONF-2013-024
	$\tilde{t}_1\tilde{t}_1, \tilde{t}_1 \rightarrow c\tilde{\chi}_1^0$	0	mono-jet/c-tag	Yes	20.3	$\tilde{t}_1$ 90-200 GeV $m(\tilde{t}_1)-m(\tilde{\chi}_1^0)<85 \text{ GeV}$ ATLAS-CONF-2013-068
	$\tilde{t}_1\tilde{t}_1$ (natural GMSB)	2 e, μ (Z)	1 b	Yes	20.3	$\tilde{t}_1$ 150-580 GeV $m(\tilde{\chi}_1^0)>150 \text{ GeV}$ 1403.5222
	$\tilde{t}_2\tilde{t}_2, \tilde{t}_2 \rightarrow \tilde{t}_1 + Z$	3 e, μ (Z)	1 b	Yes	20.3	$\tilde{t}_2$ 290-600 GeV $m(\tilde{\chi}_1^0)<200 \text{ GeV}$ 1403.5222
	$\tilde{\ell}_L\tilde{\ell}_L, \tilde{\ell} \rightarrow \ell\tilde{\chi}_1^0$	2 e, μ	0	Yes	20.3	$\tilde{\ell}$ 90-325 GeV $m(\tilde{\chi}_1^0)=0 \text{ GeV}$ 1403.5294
EW direct	$\tilde{\chi}_1^{\pm}\tilde{\chi}_1^{\mp}, \tilde{\chi}_1^{\pm} \rightarrow \tilde{\ell}\nu(\ell\bar{\nu})$	2 e, μ	0	Yes	20.3	$\tilde{\chi}_1^{\pm}$ 140-465 GeV $m(\tilde{\chi}_1^0)=0 \text{ GeV}, m(\tilde{\ell}, \bar{\nu})=0.5(m(\tilde{\chi}_1^{\pm})+m(\tilde{\chi}_1^0))$ 1403.5294
	$\tilde{\chi}_1^{\pm}\tilde{\chi}_1^{\mp}, \tilde{\chi}_1^{\pm} \rightarrow \tilde{\tau}\nu(\tau\bar{\nu})$	2 τ	-	Yes	20.7	$\tilde{\chi}_1^{\pm}$ 180-330 GeV $m(\tilde{\chi}_1^0)=0 \text{ GeV}, m(\tilde{\tau}, \bar{\nu})=0.5(m(\tilde{\chi}_1^{\pm})+m(\tilde{\chi}_1^0))$ ATLAS-CONF-2013-028
	$\tilde{\chi}_1^{\pm}\tilde{\chi}_1^{\mp} \rightarrow \tilde{\ell}_L\nu\tilde{\ell}_L\ell(\bar{\nu}\nu), \ell\bar{\nu}\tilde{\ell}_L\ell(\bar{\nu}\nu)$	3 e, μ	0	Yes	20.3	$\tilde{\chi}_1^{\pm}, \tilde{\chi}_1^{\mp}$ 700 GeV $m(\tilde{\chi}_1^0)=m(\tilde{\chi}_1^0), m(\tilde{\chi}_1^0)=0, m(\tilde{\ell}, \bar{\nu})=0.5(m(\tilde{\chi}_1^{\pm})+m(\tilde{\chi}_1^0))$ 1402.7029
	$\tilde{\chi}_1^{\pm}\tilde{\chi}_1^{\mp} \rightarrow W\tilde{\chi}_1^0Z\tilde{\chi}_1^0$	2-3 e, μ	0	Yes	20.3	$\tilde{\chi}_1^{\pm}, \tilde{\chi}_1^{\mp}$ 420 GeV $m(\tilde{\chi}_1^0)=m(\tilde{\chi}_1^0), m(\tilde{\chi}_1^0)=0$, sleptons decoupled 1403.5294, 1402.7029
	$\tilde{\chi}_1^{\pm}\tilde{\chi}_1^{\mp} \rightarrow W\tilde{\chi}_1^0h\tilde{\chi}_1^0$	1 e, μ	2 b	Yes	20.3	$\tilde{\chi}_1^{\pm}, \tilde{\chi}_1^{\mp}$ 285 GeV $m(\tilde{\chi}_1^0)=m(\tilde{\chi}_1^0), m(\tilde{\chi}_1^0)=0$, sleptons decoupled ATLAS-CONF-2013-093
	Direct $\tilde{\chi}_1^{\pm}\tilde{\chi}_1^{\mp}$ prod., long-lived $\tilde{\chi}_1^{\pm}$	Disapp. trk	1 jet	Yes	20.3	$\tilde{\chi}_1^{\pm}$ 270 GeV $m(\tilde{\chi}_1^{\mp})-m(\tilde{\chi}_1^0)=160 \text{ MeV}, \tau(\tilde{\chi}_1^{\pm})=0.2 \text{ ns}$ ATLAS-CONF-2013-069
	Stable, stopped $\tilde{g}$ R-hadron	0	1-5 jets	Yes	22.9	$\tilde{g}$ 832 GeV $m(\tilde{\chi}_1^0)=100 \text{ GeV}, 10 \mu\text{s}<\tau(\tilde{g})<1000 \text{ s}$ ATLAS-CONF-2013-057
	GMSB, stable $\tilde{\tau}, \tilde{\chi}_1^0 \rightarrow \tilde{\tau}(\tilde{e}, \tilde{\mu})+\tau(e, \mu)$	1-2 μ	-	-	15.9	$\tilde{\chi}_1^0$ 475 GeV $10<\tan\beta<50$ ATLAS-CONF-2013-058
	GMSB, $\tilde{\chi}_1^0 \rightarrow \gamma\tilde{G}$, long-lived $\tilde{\chi}_1^0$	2 γ	-	Yes	4.7	$\tilde{\chi}_1^0$ 230 GeV $0.4<\tau(\tilde{\chi}_1^0)<2 \text{ ns}$ 1304.6310
	$\tilde{q}\tilde{q}, \tilde{\chi}_1^0 \rightarrow q\tilde{q}\mu$ (RPV)	1 μ , displ. vtx	-	-	20.3	$\tilde{q}$ 1.0 TeV $1.5<\tau<156 \text{ mm}, \text{BR}(\mu)=1, m(\tilde{\chi}_1^0)=108 \text{ GeV}$ ATLAS-CONF-2013-092
RPV	LFV $pp \rightarrow \tilde{\nu}_\tau + X, \tilde{\nu}_\tau \rightarrow e + \mu$	2 e, μ	-	-	4.6	$\tilde{\nu}_\tau$ 1.61 TeV $\lambda'_{311}=0.10, \lambda'_{132}=0.05$ 1212.1272
	LFV $pp \rightarrow \tilde{\nu}_\tau + X, \tilde{\nu}_\tau \rightarrow \tau + e$	2 e, μ	-	-	4.6	$\tilde{\nu}_\tau$ 1.61 TeV $\lambda'_{311}=0.10, \lambda'_{132}=0.05$ 1212.1272
	Bill $\tilde{\chi}_1^{\pm}\tilde{\chi}_1^{\mp}$	0	6-7 jets	-	20.3	$\tilde{\chi}_1^{\pm}$ 916 GeV $\text{BR}(\tilde{t})=\text{BR}(\tilde{b})=\text{BR}(\tilde{c})=0\%$ ATLAS-CONF-2012-140
	$\tilde{\chi}_1^{\pm}\tilde{\chi}_1^{\mp}$	0	6-7 jets	-	20.3	$\tilde{\chi}_1^{\pm}$ 880 GeV ATLAS-CONF-2013-036
	$\tilde{\chi}_1^{\pm}\tilde{\chi}_1^{\mp}$	0	6-7 jets	-	20.3	$\tilde{\chi}_1^{\pm}$ 916 GeV ATLAS-CONF-2013-036
	$\tilde{g} \rightarrow q\tilde{q}q$	0	6-7 jets	-	20.3	$\tilde{g}$ 916 GeV $\text{BR}(\tilde{t})=\text{BR}(\tilde{b})=\text{BR}(\tilde{c})=0\%$ ATLAS-CONF-2013-091
	$\tilde{g} \rightarrow \tilde{t}_1 t, \tilde{t}_1 \rightarrow b s$	2 e, μ (SS)	0-3 b	Yes	20.7	$\tilde{g}$ 880 GeV ATLAS-CONF-2013-007
	Scalar gluon pair, sgluon $\rightarrow q\tilde{q}$	0	4 jets	-	4.6	sgluon 100-287 GeV incl. limit from 1110.2693 1210.4826
	Scalar gluon pair, sgluon $\rightarrow t\tilde{t}$	2 e, μ (SS)	2 b	Yes	14.3	sgluon 350-800 GeV ATLAS-CONF-2013-051
	WIMP interaction (D5, Dirac χ)	0	mono-jet	Yes	10.5	M^* scale 704 GeV $m(\chi)<80 \text{ GeV}$, limit of $<687 \text{ GeV}$ for D8 ATLAS-CONF-2012-147

<https://twiki.cern.ch/twiki/bin/view/AtlasPublic/SupersymmetryPublicResults>


$\sqrt{s} = 7 \text{ TeV}$ full data
 $\sqrt{s} = 8 \text{ TeV}$ partial data
 $\sqrt{s} = 8 \text{ TeV}$ full data

10⁻¹ 1 Mass scale [TeV]

Summary and Outlook


- **LSP** is an excellent **candidate for DM**
- Most of the SUSY scenarios predict the **LSP** to be either $\tilde{\chi}_1^0$ or $\tilde{G}$
- ATLAS has **excellent sensitivity** to dark matter produced in **cascade decays** of susy particles
- No significant excess is seen $\Rightarrow$ **limits** set on mass and cross section
- **Explore new regions** of parameter and phase space with the LHC Run2 data at **~ 14 TeV** to be collected **starting in 2015**

Stay tuned!



Backup

Upgrade Timeline and luminosity



The $m_{T2}^{\ell\ell}$ variable is calculated as

$$m_{T2}(\mathbf{p}_{T,1}, \mathbf{p}_{T,2}, \mathbf{p}_T^{\text{miss}}) = \min_{\mathbf{q}_{T,1} + \mathbf{q}_{T,2} = \mathbf{p}_T^{\text{miss}}} \{ \max[m_T(\mathbf{p}_{T,1}, \mathbf{q}_{T,1}), m_T(\mathbf{p}_{T,2}, \mathbf{q}_{T,2})] \}, \quad (1)$$

where m_T indicates the transverse mass, $\mathbf{p}_{T,1}$ and $\mathbf{p}_{T,2}$ are the transverse momentum vectors of the two leptons, and $\mathbf{q}_{T,1}$ and $\mathbf{q}_{T,2}$ are transverse vectors which satisfy $\mathbf{q}_{T,1} + \mathbf{q}_{T,2} = \mathbf{p}_T^{\text{miss}}$. The minimisation is performed over all the possible decompositions of $\mathbf{p}_T^{\text{miss}}$.




Figure 3: Efficiency of large $m_{T2}^{\ell\ell}$ selection: fraction of signal events that are selected by at least one SR of Ref. [30] as a function of top squark and τ slepton masses.