

NONTRIVIAL HOLONOMY AND COLLISIONAL ENERGY LOSS

Vladimir Skokov



WESTERN MICHIGAN UNIVERSITY

Midwest Critical Mass


MOTIVATION: QCD AT FINITE TEMPERATURE

- Commonly accepted language for description of matter in heavy-ion collisions: **finite temperature Quantum ChromoDynamics**
- Method for finite temperature field theory and QCD in particular:
 - Lattice numerical simulations (LQCD): non-perturbative first-principle approach
Limitations:
 - real-time properties (viscosity, conductivity, etc.) are very difficult to calculate (maximal entropy methods are usually applied),
 - only zero baryon density: sign problem
 - Functional methods (functional renormalization group, Schwinger-Dyson)
Limitations:
 - based on non-systematic truncations
 - AdS/CFT duality
Limitations:
 - infinite N_c
 - strong coupling limit only
 - Perturbative methods:
Limitations:
 - applicable only at large T and μ

Our goal: to include important non-perturbative effects in perturbative QCD.

Hydrodynamics is quite successful in describing two-particle correlations in $A + A$ collisions

Elliptic flow



$$v_n \sim \frac{1}{N} \int d\phi \frac{dN}{d\phi} \cos(\pi n)$$


$A+A$ RHIC and LHC: $v_2 \sim 0.1$

This is consistent with an assumption of strongly interacting matter with small viscosity

Small viscosity! Strongly interacting system!

Viscosity is proportional to mean free path (inversely proportional to scattering cross-section)


HEAVY-QUARK/JET ENERGY LOSS IN QUAR-GLUON PLASMA



- A particle traveling in the medium scatters on its constituents and loses its energy.
- Small viscosity of the matter implies strong interaction.
- Thus it is expected that the energy loss is dramatically large.
- This is not the case!


Suppression of single inclusive high $p_{\perp}$ hadrons

$$R_{AA} = \frac{d^2N/dp_{\perp}dy}{T_{AA}(0)d^2\sigma^{NN}/dp_{\perp}dy}$$




While η/s is close to its “minimal” AdS/CFT value, the energy loss seems to be overestimated in AdS/CFT.


IONIZATION IN QCD PLASMA



Neutral state $\leadsto$ confined phase,
color neutrality $>$ hadronic scale




Completely ionized plasma $\leadsto$
perturbative QGP with freely moving
charges



Partially ionized plasma $\leadsto$ *partial* ionization of color: hadrons and color charges;
semi-QGP, nontrivial holonomy

ORDER PARAMETER FOR DECONFINEMENT PHASE TRANSITION



- Wilson line $\mathbb{L} = \exp\left(ig \int_0^{1/T} A_0 d\tau\right)$;
trace of Wilson line: the Polyakov loop, $L = \text{tr}\mathbb{L}/N_c$, is gauge invariant
- Free energy of static quark, in pure glue $F(T) = F_1(r \rightarrow \infty, T)/2$
- Polyakov loop $L = \exp(-F(T)/T)$


POLYAKOV LOOP AS A MEASURE OF PARTIAL IONIZATION: PURE GLUE

$$\text{Polyakov loop: } \langle L \rangle \sim e^{-F_{\text{test qk}}/T}$$

Confined: $F_{\text{test qk}} \rightarrow \infty$,
 $\langle L \rangle \rightarrow 0$

Semi QGP: $0 < \langle L \rangle < 1$
 $\langle L \rangle$ measures degree of ionization

Perturbative QGP:
 $F_{\text{test qk}}/T \rightarrow 0$, $\langle L \rangle \rightarrow 1$




POLYAKOV LOOP AS A MEASURE OF PARTIAL IONIZATION: LQCD

$$\text{Polyakov loop: } \langle L \rangle \sim e^{-F_{\text{test qk}}/T}$$

Confined: $F_{\text{test qk}} \rightarrow \infty$,
 $\langle L \rangle \rightarrow 0$

Intermediate regime: $0 < \langle L \rangle < 1$
 $\langle L \rangle$ measures degree of ionization

Deconfined:
 $F_{\text{test qk}}/T \rightarrow 0$, $\langle L \rangle \rightarrow 1$



Non-trivial holonomy

$$\text{tr } \mathbb{L} = \exp \left(i \textcolor{red}{g} \int_0^{1/T} \textcolor{blue}{A}_0 d\tau \right)$$

can be represented by eigenvalues of Wilson line.

In mean-field approximation:


$$A_0^{\text{cl}} = \frac{i}{\textcolor{red}{g}} \text{diag}(\textcolor{blue}{Q}^1, \textcolor{blue}{Q}^2, \dots, \textcolor{blue}{Q}^{N_c})$$

The background field A_0^{cl} is manifestly non-perturbative ($\propto 1/\textcolor{red}{g}$).

**Punchline: transition region (“semi”-QGP):
must exhibit partial ionization of color
shear viscosity, energy loss... must depend upon the degree of ionization**

Energy loss

$$S_i = \frac{\text{energy loss in semi-QGP}}{\text{energy loss in perturbative QGP}}$$



- S_q - Coulomb scattering (scattering off of **light quark**, t channel)
 - $S_g^{t,u}$ - Compton scattering (scattering off of **gluons**, t and u channels)
- S_i increases as color is ionized

Usual argument of kinetic theory

Majumder, Muller and Wang, hep-ph/0703082

Liao and Shuryak, 0810.4116

Asakawa, Bass, and Muller, hep-ph/0603092, 1208.2426

- Viscosity $\eta \sim \rho^2 / \sigma$
 - ρ - density of color charges
 $\rho \sim 1$
 - σ - cross section: $\sigma \sim g^4$,
 g - coupling
 - large $g \leadsto$ small η

- Energy loss $\frac{dE}{dx} \sim g^2 \rho^2$
 - large $g \leadsto$ large $\frac{dE}{dx}$

Semi-QGP

Y. Hidaka, R. Pisarski 0912.0940

R. Pisarski, V. Skokov proceedings of QM2013

- Viscosity $\eta \sim \rho^2 / \sigma$
 - ρ - density of color charges, $\rho \sim \langle L \rangle^2$
 - σ - cross section: $\sigma \sim \langle L \rangle^2$
 - $\eta \sim \langle L \rangle^2$, small in semi-QGP

- Energy loss (large N_c)
 - $\frac{dE}{dx} \sim \langle L \rangle \cdot \frac{dE}{dx}$ on light quarks
 $+ \langle L \rangle^2 \cdot \frac{dE}{dx}$ off gluons

AdS/CFT

- small viscosity (conjectured lowest limit) $\eta/s = 1/4\pi$
- large energy loss ($\propto l^3$) (here l = length)

Collection of results

- AdS/sQCD: energy loss $\propto l^3$
- pQCD, energy loss $\propto l^2$
- **experiment**, energy loss $< l^2$
- **semi QGP** provides suppression of pQCD results

Details

- Collisional energy loss in large N_c limit
- Collisional energy loss due to scattering off of light quark, $N_c = 3$
- Collisional energy loss due to scattering off of gluons, $N_c = 3$
- Outlook: radiative energy loss, photon and dilepton production

NON-ZERO POLYAKOV LOOP $\leadsto$ NON-TRIVIAL HOLONOMY

- Finite temperature field theory: $S(1) \times R(3)$
- Polyakov loop $L = \text{Tr } \mathcal{P} \exp \left(ig \int_0^{1/T} \mathbf{A}_0 d\tau \right)$
- Ansatz for $[\mathbf{A}_0]_{ab} = \delta_{ab} \frac{Q^a}{g}$, for the sake of simplicity $Q^a = 2\pi T \cdot q^a$
- Tracelessness $\text{tr } \mathbf{A}_0 = 0 \leadsto \sum_a Q^a = \sum_a q^a = 0$
- Classical approximation: zero action for $\mathbf{A}_0$
- One loop about $\mathbf{A}_0$: Gross, Pisarski, Yaffe '81 (no functional modification to two loop order, see A. Dumitru et al, 2013)

$$U_{\text{pert}} = -2\pi^2 T^4 \left[\frac{N^2 - 1}{45} - \frac{1}{3} \sum_{a,b} (q_a - q_b)^2 (1 - |q_a - q_b|)^2 \right]$$

Gives only trivial $\mathbf{A}_0$

- Non-perturbative contribution are modeled by (R. Pisarski et al)

$$U_{\text{non-pert}} = T^2 T_d^2 \left[c_1 \sum_{a,b} |q_a - q_b| (1 - |q_a - q_b|) + c_2 \sum_{a,b} (q_a - q_b)^2 (1 - |q_a - q_b|)^2 + c_3 \right]$$

- c_i are fixed to get transition at $T = T_d$, and describe lattice data
- three colors: $q_1 = -q_2 = q, q_3 = 0$. Confining at $q = 1/3$ and perturbative $q = 0$.

DISTRIBUTION FUNCTION FOR QUARKS

- Lagrangian for quarks in background field A_0
 $\mathcal{L} = \bar{\psi} (\gamma_\mu i \mathbf{D}^\mu - m) \psi$ with $\mathbf{D}^\mu = \partial^\mu + ig \mathbf{A}^\mu$ and $\mathbf{A}^\mu = \delta^{\mu 0} A^0$, $A_4 = iA_0$
- Logarithm of partition function:

$$\begin{aligned} \ln Z &= \ln \int \mathcal{D}\psi \exp \left(- \int d^4 x_E \mathcal{L}_E \right) = \sum_a \ln \det \left(\gamma^\mu \partial_\mu + 2\pi T q_a \delta^{\mu 4} + im \right) \\ &= 2 \sum_a \ln \det \left[(\partial_4 + 2\pi q_a)^2 - \partial^2 + m^2 \right] \\ &= -2T \sum_a \int \frac{d^3 p}{(2\pi)^3} \left[\ln \left(1 + \exp \left(-E_p/T - i2\pi q_a \right) \right) \right] + \text{c.c.} \\ &= \sum_a \int \frac{d^3 p}{(2\pi)^3} \frac{p^2}{E_p} n^q(E_p - 2\pi i T q_a) + \text{c.c.}, \quad E_p = \sqrt{p^2 + m^2} \end{aligned}$$

- Quark distributing function

$$n^q(E_p - 2\pi i T q_a) = \frac{1}{\exp(E_p/T - 2\pi i q_a) + 1}$$

DISTRIBUTION FUNCTION FOR QUARKS & GLUONS IN A_0

Quark and gluon propagator in a background A_0 field: Hidaka, Pisarski 0906.1751

- Distribution function for quarks ($Q_a = 2\pi T q_a$)

$$n_a^q(p, Q) = \left[\exp\left(\frac{p - iQ_a}{T}\right) + 1 \right]^{-1}$$

- Compare the previous to usual Fermi-Dirac distribution at finite chemical potential

$$n_a^q(p) = \left[\exp\left(\frac{p - \mu}{T}\right) + 1 \right]^{-1}$$

$$\mu \rightarrow iQ_a$$

- Distribution function for gluons

$$n_{a,b}^g(p, Q) = \left[\exp\left(\frac{p - i(Q_a - Q_b)}{T}\right) - 1 \right]^{-1}$$

DISTRIBUTION FUNCTION FOR QUARKS & GLUONS: LIMITS

- Perturbative QGP (trivial holonomy), $Q = 0$

$$n^g(k, Q = 0) = \left[\exp\left(\frac{k}{T}\right) - 1 \right]^{-1} \quad n^q(k, Q = 0) = \left[\exp\left(\frac{k}{T}\right) + 1 \right]^{-1}$$

- In confined regime: $\text{tr } \mathbb{L}^n = \sum_a \exp(2\pi n i q_a) = 0$ unless $n = jN_c, j \in \mathbb{Z}$, when $\text{tr } \mathbb{L}^n = (-1)^{k(N_c+1)} N_c$ therefore

$$\sum_a n_a^q = \sum_a \sum_{n=1}^{\infty} \exp(-nk/T) \text{tr } \mathbb{L}^n = \frac{N_c}{1 + \exp(N_c \beta k)}$$

Number of quarks in confined state ($k \rightarrow N_c k$)

$$N_{\text{conf}}^q = \int [dk] \sum_a n_a^q = \frac{1}{N_c^{d-1}} \int [dk] \frac{1}{1 + \exp(\beta k)},$$

where d is the number of spatial dimensions.

To be compared with number of particles in deconfined regime N_{dc}


$$N_{\text{dc}}^q = \sum_a \int [dk] n^q(Q = 0) = N_c \int [dk] \frac{1}{1 + \exp(\beta k)}$$

In the large N_c limit, $N_{\text{conf}} \rightarrow 0$ unless $d < 2$.

- Heavy, energetic particle in the initial state: $E \gg M \gg T$.
- Collisional energy loss per unit length is proportional to the density of particles.
Scattering off of light quarks:

$$\frac{dE}{dx} \sim \sum_a \int_k n(k + 2\pi i T q_a) \times \text{Matrix element}$$

LARGE N LIMIT FOR SCATTERING OFF LIGHT QUARK: BIRDTRACKS



$$\left(\text{diagram 1} - \frac{1}{N} \text{diagram 2} \right) \left(\text{diagram 3} - \frac{1}{N} \text{diagram 4} \right) =$$

$$\begin{aligned} & \left(\begin{array}{cc} c & d \\ | & | \\ a & b \end{array} - \frac{1}{N} \begin{array}{cc} c & d \\ | & | \\ a & b \end{array} \right) \left(\begin{array}{cc} c & d \\ | & | \\ a & b \end{array} - \frac{1}{N} \begin{array}{cc} c & d \\ | & | \\ a & b \end{array} \right) + \frac{1}{N^2} \text{diagram 5} \bigcirc_N = \\ & \begin{array}{cc} c & d \\ | & | \\ a & b \end{array} - \frac{1}{N} \begin{array}{cc} c & d \\ | & | \\ a & b \end{array} \end{aligned}$$

large N $\rightarrow$

Generators of $SU(N_c)$: **'t Hooft's double line basis** (N_c^2 generators)

$$(t^{ab})_{cd} = \frac{1}{\sqrt{2}} \left(\delta_c^a \delta_d^b - \frac{1}{N_c} \delta^{ab} \delta_{cd} \right) \quad \text{Tr } t^{ab} t^{cd} = \frac{1}{2} P^{(ab)(cd)}$$




$$\begin{aligned} \frac{dE}{dx} &\propto \sum_{a,b}^N \int [dk][dk'] [dp'] f(p, k, k', p') n(E_k + iQ_a) [1 - n(E_{k'} + iQ_b)] \\ &\quad \sum_a n(E_k + iQ_a) = \sum_a [\exp(\beta E_k + i\beta Q_a) + 1]^{-1} \\ &= \sum_{n=1}^{\infty} (-1)^n \exp(-n\beta E_k) \sum_a \exp(in\beta Q_a) = \sum_{n=1}^{\infty} (-1)^n \exp(-n\beta E_k) \text{tr } L^n \end{aligned}$$

for small $\text{tr } L$ scattering off **light quarks**

$$\frac{dE}{dx} \propto \text{tr } L \cdot \left(\frac{dE}{dx} \right)_{\text{pert.}}$$

Similar argument for scattering off **gluons** gives

$$\frac{dE}{dx} \propto (\text{tr } L)^2 \cdot \left(\frac{dE}{dx} \right)_{\text{pert.}}$$




As in perturbative QCD, but taking into account modification of distribution function in final/initial state.

N=3: SCATTERING OFF LIGHT QUARKS

$$\frac{dE}{dx} = \sum_i \int_k \frac{n_i(k)}{2k} \int_{k'} \frac{1 \pm n_i(k')}{2k'} \int_{p'} \frac{(2\pi)^4 \delta^{(4)}(P + K - P' - K') |\mathcal{M}_i|^2 \omega}{4EE'd}$$

$$\frac{1}{d} \sum_i |\mathcal{M}_i|^2 = \frac{8N_f g^4}{N_c} \frac{2(s - M^2)^2 + (u - M^2)^2 + 2M^2 t}{t^2} \times \frac{N_c^2 - 1}{4N_c}$$



Keeping only leading log terms

$$\left. \frac{dE(A_0^{\text{cl}})}{dx} \right|_q = S_q(A_0^{\text{cl}}) \left(\frac{dE}{dx} \right)_q^{\text{pert}}; \quad \left(\frac{dE}{dx} \right)_q^{\text{pert}} = \alpha_s^2 T^2 N_f \frac{N_c^2 - 1}{12N_c} \pi \ln \frac{ET}{m_D^2}$$

with semi QGP suppression factor

$$S_q(A_0^{\text{cl}}) = \frac{12}{\pi^2} \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^2} \frac{\text{tr} \mathbb{L}^n}{N_c}$$


- pert. limit $A_0 = 0$: $S_q(A_0^{\text{cl}} = 0) = 1$
- confining limit: $S_q(A_0^{\text{cl}}) = 1/N_c^2$

N=3: SCATTERING OFF GLUONS

$$S_i = \left(\frac{dE}{dx} \right)_i \bigg/ \left(\frac{dE}{dx} \right)_{i,\text{pert.}}$$


- t channel

$$S_g^t(A_0^{\text{cl}}) = \frac{1}{N_c^2 - 1} \left(\frac{6}{\pi^2} \sum_{n=1}^{\infty} \frac{|\text{tr} \mathbb{L}^n|^2}{n^2} - 1 \right); \quad \left. \frac{dE}{dx} \right|_g^{t,\text{pert}} = \alpha_s^2 T^2 (N_c^2 - 1) \frac{\pi}{6} \ln \frac{ET}{m_D^2}$$




- u channel, the same suppression factor

$$\left. \frac{dE}{dx} \right|_g^{u,\text{pert}} = \alpha_s^2 T^2 C_F^2 \frac{\pi}{6} \ln \frac{ET}{M^2}, \quad C_F = (N_c^2 - 1)/(2N_c)$$



- Limits: perturbative: $S_g(A_0^{\text{cl}} = 0) = 1$
confining background field: $S_g = 0$

POLYAKOV LOOP: LQCD



POLYAKOV LOOP AS A FUNCTION OF T : PERTURBATIVE CORRECTIONS

- One needs to extract the eigenvalues of the Wilson line, q_a
- One can use models or first principle **lattice QCD** calculations
- **Lattice QCD**: not only classical field contributions to **Polyakov Loop**.
Polyakov loop receives the perturbative corrections
(Burnier, Laine, Veplalainen 2009)





- Perturbative corrections exponentiate, i.e. for the case of the non-trivial holonomy

$$L_{w/pert}(A_0^{cl}) = L(A_0^{cl}) \exp \left[\delta L_{pert}(A_0^{cl}) \right]$$

The dependence of the perturbative corrections on the background field A_0

$$\delta L_{pert}(A_0^{cl}) \approx \delta L_{pert}(0)$$

POLYAKOV LOOP AS A FUNCTION OF T : $N_c = 3$



S. Lin, R. Pisarski and V.S. arXiv:1312.3340





$$\mathbf{q} = (-q, 0, q); \quad L = \frac{1}{3} [1 + 2 \cos(2\pi q)]$$

$\text{tr} \mathbb{L}$ is known, $\text{tr} \mathbb{L}^n$ are needed

Lattice: Karsch et al

$$\text{tr} \mathbb{L}^n = 1 + 2 \cos(2\pi n q)$$

NUMERICAL RESULTS



$$S_i = \left(\frac{dE}{dx} \right)_i \bigg/ \left(\frac{dE}{dx} \right)_{i,\text{pert.}}$$

For three colors analytical result:



$$S_q^{N_c=3} = 1 - 8 q^2$$

$$S_g^{N_c=3} = 1 - 3 q (2 - 3 q)$$

- Different processes are suppressed differently
- Processes with gluons are suppressed stronger than those with quarks
- for quarks:
 $S_q \rightarrow 1/N_c^2$ at low temperatures

S. Lin, R. Pisarski and V.S. arXiv:1312.3340

RADIATIVE ENERGY LOSS: PRELIMINARY LARGE N RESULT



Suppressed by


$$n_{-a}(1 - n_c)(1 + n_{-c,-b}) \sim \text{tr } L$$

Correct treatment includes ladder diagram resummation, work in progress...

PHOTON PRODUCTION

Based on very similar arguments

Naive calculations include only



Suppression: $\text{tr } L$

Suppression: at given T the density of quarks is smaller.

However, complete leading order results include collinear emission, nonlinear quark annihilation and LPM effects, Arnold Moore Yaffe (AMY) '00


AMY in nontrivial background A_0 : work in progress...

Color singlet in the final state, no suppression at leading order.

Therefore high T photons are suppressed, but not high T dileptons.


Naive argument:

- photon emission $\propto T^4$ and thus the highest in quark-gluon phase
- the flow of matter is the smallest in initial state
- thus, it was expected that photon azimuthal anisotropy is smaller than hadronic one (from final state) by a factor of ~ 5




PHOTON AZIMUTHAL ANISOTROPY

- Direct photons:



- Hadrons:



- experiment (PHENIX at RHIC and ALICE at LHC) showed that hadron and photon azimuthal anisotropy is approximately the same
- there is either an alternative mechanism for photon azimuthal anisotropy (e.g. conformal anomaly and magnetic field G. Bazar, D. Kharzeev and V.S. Phys.Rev.Lett. 109 (2012) 202303) or there is a **suppression** of photon production in QGP state

Our mechanism provides this suppression for photon, but not for dileptons!

Dileptons: no suppression! As a consequence $v_2^{l^+l^-} < v_2^\gamma$

CONCLUSIONS: PERTURBATIVE QCD MEETS HEAVY-ION PHENOMENOLOGY

- semiQGP: nontrivial holonomy as an instrument to include non-perturbative confinement effects into perturbative QCD
- calculations in a non-trivial background classical fields: quite tedious, but doable
- collisional energy loss: suppression factor for scattering off of quarks is proportional to the Polyakov loop, scattering off of gluons is proportional to the second power of the Polyakov loop
- this approach may help to solve two puzzles of HIC phenomenology:
 - strong-coupling paradox: small viscosity and moderate energy loss
 - large azimuthal anisotropy of photons

Thank you!