

# Determination of $K\pi$ scattering lengths at physical kinematics

Tadeusz Janowski  
Supervisor: Prof. C. T. Sachrajda

RBC-UKQCD collaboration  
SHEP  
University of Southampton  
tj1g11@soton.ac.uk

June 23, 2014

## UKQCD

Rudy Arthur (Odense)  
Peter Boyle (Edinburgh)  
Luigi Del Debbio (Edinburgh)  
Shane Drury (Southampton)  
Jonathan Flynn (Southampton)  
Julien Frison (Edinburgh)  
Nicolas Garron (Dublin)  
Jamie Hudspith (Toronto)  
Tadeusz Janowski (Southampton)  
Andreas Juettner (Southampton)  
Ava Kamseh (Edinburgh)  
Richard Kenway (Edinburgh)  
Andrew Lytle (TIFR)  
Marina Marinkovic (Southampton)  
Brian Pendleton (Edinburgh)  
Antonin Portelli (Southampton)  
Thomas Rae (Mainz)  
Chris Sachrajda (Southampton)  
Francesco Sanfilippo (Southampton)  
Matthew Spraggs (Southampton)  
Tobias Tsang (Southampton)

## RBC

Ziyuan Bai (Columbia)  
Thomas Blum (UConn/RBRC)  
Norman Christ (Columbia)  
Xu Feng (Columbia)  
Tomomi Ishikawa (RBRC)  
Taku Izubuchi (RBRC/BNL)  
Luchang Jin (Columbia)  
Chulwoo Jung (BNL)  
Taichi Kawanai (RBRC)  
Chris Kelly (RBRC)  
Hyung-Jin Kim (BNL)  
Christoph Lehner (BNL)  
Jasper Lin (Columbia)  
Meifeng Lin (BNL)  
Robert Mawhinney (Columbia)  
Greg McGlynn (Columbia)  
David Murphy (Columbia)  
Shigemi Ohta (KEK)  
Eigo Shintani (Mainz)  
Amarjit Soni (BNL)  
Sergey Syritsyn (RBRC)  
Oliver Witzel (BU)  
Hantao Yin (Columbia)  
Jianglei Yu (Columbia)  
Daiqian Zhang (Columbia)

# Scattering phase shifts

For a system with two (pseudo)scalar particles, the S-matrix in centre-of-mass frame can be written in partial wave basis as:

$$\langle E', p', l', m' | S | E, 0, l, m \rangle = \delta(E' - E) \delta(p) \delta_{ll'} \delta_{mm'} S_l(E)$$

which is required by Lorentz invariance of the S-matrix, specifically  $[H, S] = 0$ ,  $[P, S] = 0$ ,  $[J^2, S] = 0$ ,  $[J_3, S] = 0$  and  $[J_{\pm}, S] = 0$ . Furthermore, unitarity of the S-matrix implies  $S^\dagger S = SS^\dagger = 1$  gives

$$S_l(E) = e^{2i\delta_l(E)}$$

This means that the two (pseudo)scalar particle scattering can be expressed in terms of a single real parameter  $\delta_l(E)$  called the *phase shift*.

# Scattering length

At low energies (or equivalently momenta,  $k$ ), phase shifts have the following threshold behaviour:

$$\delta_l(k) \sim k^{l+1}$$

- The dominant contribution will come from the s-wave ( $l = 0$ ).
- We can define a constant called the *scattering length*:

$$(\delta_0(k)/k)^{-1} = \frac{1}{a_0} + \frac{r_{\text{eff}}}{2}k^2 + \mathcal{O}(k^4)$$

# $K\pi$ scattering

With  $m_u = m_d \equiv m_{ud}$  and QCD interactions only, isospin becomes a good quantum number.

Pions have  $I = 1$ , kaons have  $I = 1/2$ , so  $K\pi$  can be in  $I = 3/2$  or  $I = 1/2$  state. Specifically:

$$|I = 3/2; I_z = 3/2\rangle = |K^+\pi^+\rangle$$

$$|I = 3/2; I_z = 1/2\rangle = \frac{1}{\sqrt{3}} |K^0\pi^+\rangle + \sqrt{\frac{2}{3}} |K^+\pi^0\rangle$$


$$|I = 3/2; I_z = -1/2\rangle = \frac{1}{\sqrt{3}} |K^+\pi^-\rangle + \sqrt{\frac{2}{3}} |K^0\pi^0\rangle$$

$$|I = 3/2; I_z = -3/2\rangle = |K^0\pi^-\rangle$$

$$|I = 1/2; I_z = 1/2\rangle = \frac{1}{\sqrt{3}} |K^+\pi^0\rangle - \sqrt{\frac{2}{3}} |K^0\pi^+\rangle$$

$$|I = 1/2; I_z = -1/2\rangle = -\frac{1}{\sqrt{3}} |K^0\pi^0\rangle + \sqrt{\frac{2}{3}} |K^+\pi^-\rangle$$

# Results so far



Plot courtesy of G. Colangelo.

	$a_0^{3/2} m_\pi$	$a_0^{1/2} m_\pi$
NPLQCD <sup>1</sup>	$-0.0574(16) \begin{pmatrix} +24 \\ -58 \end{pmatrix}$	$0.1725(13) \begin{pmatrix} +23 \\ -156 \end{pmatrix}$
Fu <sup>2</sup>	$-0.0512(18)$	$0.1819(35)$
PACS-CS <sup>3</sup>	$-0.0602(31)(26)$	$0.183(18)(35)$

Calculation also done by Lang et. al. <sup>4</sup>, but without extrapolation to physical point.

This work: evaluation of scattering length **directly at physical point**.

---

<sup>1</sup>S. R. Beane et al. [NPLQCD Collaboration], Phys. Rev. D 74 (2006) 114503


<sup>2</sup>Z. Fu, Phys. Rev. D 85 (2012) 074501

<sup>3</sup>Kiyoshi Sasaki (Tokyo Inst. Tech.) et al. [PACS-CS Collaboration], Phys.Rev. D89 (2014) 054502

<sup>4</sup>C. B. Lang, L. Leskovec, D. Mohler and S. Prelovsek, Phys. Rev. D 86 (2012) 054508

# S-wave phase shifts from a lattice - Lüscher's formula

In infinite volume, the two meson energies can be visualised on the complex energy plane as a branch cut starting at  $m_1 + m_2$ . In finite volume this is replaced by series of poles.



Position of these poles can be related to the s-wave phase shifts by Lüscher's condition <sup>5</sup>:

$$\delta(k) + \phi(q) = n\pi$$

$$\tan \phi(q) = \frac{q\pi^{3/2}}{Z_{00}(1; q)}$$

$$Z_{00}(1; q) = \frac{1}{\sqrt{4\pi}} \sum_{k \in \mathbb{Z}^3} \frac{1}{q^2 - k^2}$$

<sup>5</sup>M. Lüscher, Nucl. Phys. B354 (1991) 531-578



# Two meson correlation function - choice of interpolators

$$\begin{aligned} C_{K\pi}^{'ij}(t) &\equiv \langle 0 | O_{K\pi}^{i\dagger}(t) O_{K\pi}^j(0) | 0 \rangle \\ &= \sum_n \langle 0 | O_{K\pi}^{i\dagger} | n \rangle \langle n | O_{K\pi}^j | 0 \rangle e^{-E_n t} \\ &\xrightarrow{t \rightarrow \infty} \langle 0 | O_{K\pi}^{i\dagger} | K\pi \rangle \langle K\pi | O_{K\pi}^j | 0 \rangle e^{-E_{K\pi} t} \end{aligned}$$

We use:

$$O_{K\pi}^{\pm}(t) = \bar{s}(t \pm 2) \gamma^5 l(t \pm 2) \bar{l}(t) \gamma^5 l(t)$$

# Noise sources

Trick to get mesons with point source: invert the quark propagator with a stochastic source  $\eta$ , so that its propagator becomes:

$$S_{\eta}(x_{snk}) = \sum_{x_{src}} S(x_{snk}, x_{src}) \eta(x_{src})$$

where  $\eta$  are independent random variables that satisfy:

$$\langle \eta(x) \eta^{\dagger}(y) \rangle = \delta(x - y),$$

where the angle bracket denotes an average over number of 'hits' (independent random vectors).

For 2 propagators at the same time slice:

$$\langle S_{\eta}(x) S_{\eta}^{\dagger}(y) \rangle = \sum_{x_0, y_0} S(x, x_0) \langle \eta(x_0) \eta^{\dagger}(y_0) \rangle S^{\dagger}(y_0, x_0) = \sum_z S(x, z) S^{\dagger}(z, y)$$

We use  $\eta(x) = \pm 1 \pm i$  with number of hits equal to 1.

$K\pi$  at  $l=1/2$  has a scalar resonance  $K_0^*(800)$  (called  $\kappa$ ). At physical kinematics,  $\kappa$  mass is larger than  $E_{K\pi}^{l=1/2}$ , so the ground state still corresponds to  $E_{K\pi}^{l=1/2}$ .

Previous studies have shown that the two-meson interpolators have a good overlap with  $K\pi$  states for light pion masses. Another viable choice would be the scalar meson interpolator:




$$O(t) = \bar{s}(t)d(t). \quad (1)$$

This operator however was shown<sup>6</sup> to only have a good overlap with  $K\pi$  state for heavy pion masses ( $> 700$  MeV) and was not considered in our run.

---

<sup>6</sup>Kiyoshi Sasaki (Tokyo Inst. Tech.) et al. [PACS-CS Collaboration], Phys.Rev. D89 (2014) 054502

# $K\pi$ $I=3/2$ contractions




$$D = \text{Tr} \left( S^\dagger(t; 2) L(t; 2) \right) \text{Tr} \left( L(t + 2; 0)^\dagger L(t + 2; 0) \right)$$

$$C = \text{Tr} \left( S^\dagger(t; 2) L(2; 0) L^\dagger(t + 2; 0) L(t + 2; 2) \right)$$

$$C_{K\pi}^{I=3/2}(t) = D - C$$

# Rectangle graph for $l=1/2$ correlator




$$C_{K\pi}^{l=1/2}(t) = D + 0.5C - 1.5R$$


# $\pi\pi$ $I=2$ scattering

For  $\pi\pi$  scattering the energy difference can be accurately predicted from:

$$\frac{C_{\pi\pi}}{C_{\pi}^2} \approx N e^{-(E_{\pi\pi} - 2m_{\pi})t} \approx N(1 - (E_{\pi\pi} - 2m_{\pi})t)$$



# $K\pi$ $I=3/2$ scattering



# Around-the-world effects


In lattice simulations we often use (anti)periodic boundary conditions in the time direction. This means that, rather than e.g.  $\langle 0 | O_1(t_1) O(t_2) | 0 \rangle$  for some operators  $O_1, O_2$ , we're actually calculating:

$$\sum_n \langle n, t = T \sim 0 | e^{-H(T-t_1)} O_1 e^{-H(t_1-t_2)} O_2 e^{-Ht_2} | n, t = 0 \rangle$$

This sum will contain not only the desired term, but also other contributions, referred to as 'around-the-world effects'.



# Around-the-world effects




## 5 parameter fit

$$\begin{aligned}C_{K\pi}(t) = & Ae^{-E_{K\pi}(t-2)} \\& + Be^{-E_{K\pi}(T-t-2)} \\& + Ce^{-m_K(t-2)}e^{-m_\pi(T-t-2)} \\& + De^{-m_\pi(t-2)}e^{-m_K(T-t-2)}\end{aligned}$$

with:

$$\begin{aligned}A &= \langle 0 | O_{snk} | K\pi \rangle \langle K\pi | O_{src} | 0 \rangle \\B &= \langle K\pi | O_{snk} | 0 \rangle \langle 0 | O_{src} | K\pi \rangle \\C &= \langle \pi | O_{snk} | K \rangle \langle K | O_{src} | \pi \rangle \\D &= \langle K | O_{snk} | \pi \rangle \langle \pi | O_{src} | K \rangle\end{aligned}$$

# 5-parameter fit - test results



# Physical run parameters

Lattice size	$48^3 \times 96$
Gauge action	Iwasaki
Fermion action	Möbius DWF
$L_s$	24
M	1.8
$\beta$	2.13
$am_s$	0.0362
$am_l$	0.00078
$a^{-1}$	1.73(3) GeV
$am_K$	0.08079(24)
$am_\pi$	0.28886(35)

- quark sources every second time slice (48 per configuration)
- antiperiodic boundary conditions in time direction only

# Preliminary results

All results shown are **PRELIMINARY**, based on 27 gauge configurations.

$I=3/2$

$$E_{K\pi} = 0.36978(52)$$

$$E_{K\pi} - m_K - m_\pi = 0.00045(31)$$

$$a_0^{3/2} m_\pi = -0.039(26)$$

$I=1/2$

$$E_{K\pi} = 0.36760(62)$$

$$E_{K\pi} - m_K - m_\pi = -0.00173(53)$$

$$a_0^{1/2} m_\pi = 0.174(60)$$

# Comparison

	$a_0^{3/2} m_\pi$	$a_0^{1/2} m_\pi$
NPLQCD	$-0.0574(16) \left( \begin{array}{c} +24 \\ -58 \end{array} \right)$	$0.1725(13) \left( \begin{array}{c} +23 \\ -156 \end{array} \right)$
Fu	$-0.0512(18)$	$0.1819(35)$
PACS-CS	$-0.0602(31)(26)$	$0.183(18)(35)$
RBC-UKQCD ( <b>preliminary</b> )	$-0.039(26)$	$0.174(60)$

- We are able to generate ensembles with physical pion and kaon masses.
- Calculation of  $K\pi$  energies at low values of  $m_\pi T$  and  $m_K T$  suffers from significant around-the world effects.
- Around-the-world effects can be treated reliably using a 5-parameter fit.
- First calculation of scattering lengths that does not rely on chiral perturbation theory.
- Although low statistics prevent us from obtaining an accurate  $I = 3/2$  result, we can get a good estimate for  $I = 1/2$ , which has been dominated by  $\chi^{PT}$  errors in previous calculations.

Thank you for your attention!