32nd International Symposium on Lattice Field Theory (Lattice 2014)



Contribution ID: 180

Type: Talk

Renormalization constants for $N_{\rm f}=2+1+1$ twisted mass QCD

Tuesday, 24 June 2014 16:30 (20 minutes)

We summarize recent non-perturbative results obtained for renormalization constants computed in the RI'-MOM scheme for $N_{\rm f} = 2 + 1 + 1$ twisted mass QCD. Our implementation employs the Iwasaki gauge action and four dynamical degenerate twisted mass fermions. Renormalization constants for scalar, pseudo-scalar, vector and axial operators, as well as for the quark propagator, are computed at three different values of the lattice spacing, two different volumes and several values of the twisted mass. Our method allows for a precise cross-check of the running, because of the particular proper treatment of the hypercubic artifacts. Preliminary results for twist-2 operators are also presented.

Primary author: Dr ZAFEIROPOULOS, Savvas (LPC Universite Blaise Pascal CNRS)

Co-authors: Dr BLOSSIER, Benoit (LPT Orsay); Dr RODRIGUEZ-QUINTERO, Jose (Universidad de Huelva); Dr BRINET, Mariane (LPSC Grenoble); Dr PENE, Olivier (LPT-Orsay); Dr GUICHON, Pierre (CEA-Saclay); Dr MORE-NAS, Vincent (LPC Universite Blaise Pascal)

Presenter: Dr ZAFEIROPOULOS, Savvas (LPC Universite Blaise Pascal CNRS)

Session Classification: Standard model parameters and renormalization

Track Classification: Standard Model Parameters and Renormalization