



Contribution ID: 10

Type: **Talk**

Study of axial magnetic effect

Thursday, 26 June 2014 14:35 (20 minutes)

The axial magnetic field, which couples to left- and right-handed fermions with opposite signs, may generate an equilibrium dissipationless energy flow of fermions in the direction of the field even in the presence of interactions. In this report numerical observation of the Axial Magnetic Effect in SU(2) lattice gauge theory is presented. The temperature behavior of the Axial Magnetic Effect is studied. It is shown that in the confinement (hadron) phase the effect is absent. In the deconfinement transition region the conductivity quickly increases, reaching the asymptotic T^2 behavior in a deep deconfinement (quark-gluon plasma) phase.

Primary author: Mr BRAGUTA, Victor (IHEP)

Co-authors: Mr MOLOCHKOV, Alexander (Far Eastern Federal University); Mr LANDSTEINER, Karl (Universidad Autonoma de Madrid); Mr CHERNODUB, Maxim (Department of Physics and Astronomy, University of Gent); Mr ULYBYSHEV, Maxim (ITEP); Mr GOY, Vladimir (Far Eastern Federal University)

Presenter: Mr BRAGUTA, Victor (IHEP)

Session Classification: Vacuum Structure and Confinement

Track Classification: Vacuum Structure and Confinement