32nd International Symposium on Lattice Field Theory (Lattice 2014)



Contribution ID: 203

Type: Talk

## The Chiral Condensate of One-Flavor QCD and the Dirac Spectrum at $\theta = 0$

Wednesday, 25 June 2014 09:00 (20 minutes)

In a sector of fixed topological charge, the chiral condensate has a discontinuty given by the Banks-Casher formula also in the case of one-flavor QCD. However, at fixed  $\theta$  angle, the chiral condensate remains constant when the quark mass crosses zero. To reconcile these contradictory observations, we have evaluated the spectral density of one-flavor QCD at  $\theta = 0$ . For negative quark mass, it becomes a strongly oscillating function with a period given by the inverse space-time volume and an amplitude that increases expoentially with the space-time volume. As we have learned from QCD at nonzero chemical potential, if this is the case, an alternative to the Banks-Casher formula applies, and we will demonstrate that for one-flavor QCD, this results into a continuous chiral condensate. A special role is played by the topological zero modes which have to be taken into account exactly in order to get a finite chiral condensate in the thermodynamic limit.

**Primary authors:** Prof. VERBAARSCHOT, Jacobus (Stony Brook University); Prof. WETTIG, Tilo (University of Regensburg)

Presenter: Prof. VERBAARSCHOT, Jacobus (Stony Brook University)

Session Classification: Chiral Symmetry

Track Classification: Chiral Symmetry