

## Abstract

Neutrino decay modifies neutrino propagation in a unique way; not only is there flavor changing as there is in neutrino oscillations, there is also energy transport from initial to final neutrinos. The most sensitive direct probe of neutrino decay is currently IceCube which can measure the energy and flavor of neutrinos traveling over extragalactic distances. For the first time we calculate the flavor transition probability for the cases of visible and invisible neutrino decay, including the effects of the expansion of the universe, and consider the implications for IceCube. As an example, we demonstrate how neutrino decay addresses a tension in the IceCube data.

# Astrophysical Neutrino Decay

Peter B. Denton

HET Lunch Discussion

May 22, 2020

1805.05950 with Irene Tamborra  
and

2005.07200 with Asli Abdullahi

[github.com/PeterDenton/Astro-Nu-Decay](https://github.com/PeterDenton/Astro-Nu-Decay)

[peterdenton.github.io/Data/Visible\\_Decay/index.html](https://peterdenton.github.io/Data/Visible_Decay/index.html)



# Overview

1. The global neutrino decay picture
2. How to calculate visible neutrino decay for astrophysics
3. The impact of the different parameters
4. Hints of neutrino decay at IceCube

# Neutrino Decay

Since neutrinos have different masses, they decay

- ▶ Loop suppressed
- ▶ Long lifetime:  $\tau \gtrsim 10^{35}$  years

Test this!

Typical Lagrangian for  $\nu_i \rightarrow \nu_j + \phi$  with  $m_i > m_j$

$$\mathcal{L} \supset \frac{g_{ij}}{2} \bar{\nu}_j \nu_i \phi + \frac{g'_{ij}}{2} \bar{\nu}_j i \gamma_5 \nu_i \phi$$

# Neutrino Decay Phenomenology

Neutrino decay is phenomenologically classified into:

- ▶ **Invisible decay:**

- ▶ The decay products are sterile or too low energy to be detected
- ▶ Results in a *depletion* of the flux below the relevant energy

- ▶ **Visible decay:**

- ▶ Decay products are detected
- ▶ In addition to depletion, there is *regeneration*
- ▶ Regeneration happens at a lower energy than depletion


# Decay Regimes

The decay width in *lab frame* is  $\Gamma_i$

$\nu_i$  has lifetime  $\tau_i = E/m_i\Gamma_i$

- ▶ **No decay (SM):**  $\Gamma_i L \ll 1$
- ▶ **Partial decay:**  $\Gamma_i L \sim 1$
- ▶ **Full decay:**  $\Gamma_i L \gg 1$

# Invisible $\nu$ Decay Constraints and Evidence



M. Gonzalez-Garcia and M. Maltoni [0802.3699](#)

J. Berryman, A. de Gouvea, D. Hernandez [1411.0308](#)

G. Pagliaroli, et al. [1506.02624](#)

PBD, I. Tamborra [1805.05950](#)

Kamiokande-II, PRL 58 1490 (1987)

S. Ando [hep-ph/0307169](#)

S. Hannestad, G. Raffelt [hep-ph/0509278](#)

A. Long, C. Lunardini, E. Sabancila [1405.7654](#)

# Why IceCube for Neutrino Decay

- ▶ DSNB and  $C\nu B$  are still some time off
- ▶ The next galactic supernova could come tomorrow, or in fifty years
- ▶ If  $\nu_1$  is stable SN1987A isn't too relevant (25 events + theory uncertainties)
  - ▶ Mass ordering looks to be normal at  $\sim 3 - 3.5 \sigma$
  - ▶ Texture in the mixing matrix
  - ▶ If  $m_1 \gtrsim m_\phi$
- ▶ Early universe constraints don't constrain *decay* just the typical decay diagram  
G. Dvali and L. Funcke [1602.03191](#)
- ▶ IceCube measures **all three flavors** over  $> 1$  decade in energy
- ▶ Astrophysical uncertainties seem like a problem, aren't really



# How to Calculate Visible Neutrino Decay

## Ingredients:

1. Oscillation averaged/decohered SM contribution
2. Depletion component
  - ▶ This takes us to invisible decay
3. Regeneration component at lower energies
  - ▶ This takes us to visible decay

## Steps:

1. Integrate over decay location
2. Integrate over initial energy spectrum due to regeneration
3. Include multiple decays
4. Include cosmology
5. Mix thoroughly, let bake for an hour

## SM Contribution: How to Calculate

First we define a “probability”

$$P_{\alpha\beta}(E_f) \equiv \frac{\Phi_{\alpha\beta}^E(E_f)}{\Phi_{\alpha}^S(E_f)}$$

Not actually a probability as it can be more than 1, but is probability-like and is useful  
Over large distances the mass states decohere and/or the wave packets separate

$$\frac{\Delta m^2 L}{E} \gg 1$$

This is easily satisfied for extragalactic sources

Wave packet separation results in an identical flux to oscillation averaging:

$$\sin^2\left(\frac{\Delta m^2 L}{4E}\right) \rightarrow \frac{1}{2}$$

## SM Contribution: The Probability

Given the usual Hamiltonian,

$$H = U_{\text{PMNS}} \begin{pmatrix} 0 & & \\ & \frac{\Delta m_{21}^2}{2E} & \\ & & \frac{\Delta m_{31}^2}{2E} \end{pmatrix} U_{\text{PMNS}}^\dagger$$

The SM oscillation probability is:

$$P_{\alpha\beta}^{\text{SM}} = \left| U_{\alpha 1}^* U_{\beta 1} + U_{\alpha 2}^* U_{\beta 2} e^{-i \frac{\Delta m_{21}^2 L}{2E}} + U_{\alpha 3}^* U_{\beta 3} e^{-i \frac{\Delta m_{31}^2 L}{2E}} \right|^2$$

When averaged/decohered:

$$\bar{P}_{\alpha\beta}^{\text{SM}} = \sum_{i=1}^3 |U_{\alpha i}|^2 |U_{\beta i}|^2$$

No interference terms.

## Depletion Component: How to Calculate

$$H = U_{\text{PMNS}} \begin{pmatrix} 0 & & \\ & \frac{\Delta m_{21}^2}{2E} - \frac{i}{2}\Gamma_2 & \\ & & \frac{\Delta m_{31}^2}{2E} - \frac{i}{2}\Gamma_3 \end{pmatrix} U_{\text{PMNS}}^\dagger$$

Assume here and throughout that  $\nu_1$  is stable (no lighter sterile neutrino) and the normal ordering

# Depletion Component: How to Calculate

$$H = U_{\text{PMNS}} \begin{pmatrix} 0 & & \\ & \frac{\Delta m_{21}^2}{2E} - \frac{i}{2}\Gamma_2 & \\ & & \frac{\Delta m_{31}^2}{2E} - \frac{i}{2}\Gamma_3 \end{pmatrix} U_{\text{PMNS}}^\dagger$$

Assume here and throughout that  $\nu_1$  is stable (no lighter sterile neutrino) and the normal ordering

The partial width including scalar and pseudo-scalar as well as  $\nu \rightarrow \nu$  and  $\nu \rightarrow \bar{\nu}$

$$\Gamma_{ij} = \frac{m_i m_j}{16\pi E_i} \{ g_{ij}^2 [f(x_{ij}) + k(x_{ij})] + g'_{ij}{}^2 [h(x_{ij}) + k(x_{ij})] \}$$

$$f(x) = \frac{x}{2} + 2 + \frac{2}{x} \log x - \frac{2}{x^2} - \frac{1}{2x^3}$$

$$h(x) = \frac{x}{2} - 2 + \frac{2}{x} \log x + \frac{2}{x^2} - \frac{1}{2x^3}$$

$$k(x) = \frac{x}{2} - \frac{2}{x} \log x - \frac{1}{2x^3}$$

$$\begin{aligned} \Gamma_i &= \sum_j \Gamma_{ij} \\ \tau_i &= m_i / E_i \Gamma_i \\ x_{ij} &\equiv m_i / m_j \end{aligned}$$

See slide 28 on  $\nu/\bar{\nu}$

## Depletion Component: The Probability

$$\bar{P}_{\alpha\beta}^{\text{dep}}(E, L) = -|U_{\alpha 2}|^2 |U_{\beta 2}|^2 (1 - e^{-\Gamma_2 L}) - |U_{\alpha 3}|^2 |U_{\beta 3}|^2 (1 - e^{-\Gamma_3 L})$$

The invisible probability is:

$$\bar{P}_{\alpha\beta}^{\text{inv}} = \bar{P}_{\alpha\beta}^{\text{SM}} + \bar{P}_{\alpha\beta}^{\text{dep}}$$

$$\bar{P}_{\alpha\beta}^{\text{inv}} = |U_{\alpha 1}|^2 |U_{\beta 1}|^2 + |U_{\alpha 2}|^2 |U_{\beta 2}|^2 e^{-\Gamma_2 L} + |U_{\alpha 3}|^2 |U_{\beta 3}|^2 e^{-\Gamma_3 L}$$

# Regeneration Component: How to Calculate

Steps:

1. Shift to mass basis
2. Integrate amplitude squared over decay position

Assume everything is incoherent, see slide 20

3. Integrate over initial neutrino energy weighted by initial neutrino flux
4. Add double decays and cosmology

This is where the meat in the recipe is.

## Regeneration Component: Amplitude

The  $\nu_i \rightarrow \nu_j$  amplitude contains

1. the survival of  $\nu_i$  over a distance  $L_1$ ,  $e^{-\frac{1}{2}\Gamma_i L_1}$ ,
2. the phase accumulation of  $\nu_i$  from the source to  $L_1$ ,  $e^{-iE_i L_1}$ ,
3. the decay of  $\nu_i$  and appearance of  $\nu_j$ ,  $\sqrt{\Gamma_{ij} W_{ij}}$ ,
4. for unstable  $\nu_j$ , survival until Earth,  $e^{-\frac{1}{2}\Gamma_j(L-L_1)}$ ,
5. and the phase accumulation of  $\nu_j$  until Earth,  $e^{-iE_f(L-L_1)}$ .



## Regeneration Component: Amplitude

The  $\nu_i \rightarrow \nu_j$  amplitude contains

1. the survival of  $\nu_i$  over a distance  $L_1$ ,  $e^{-\frac{1}{2}\Gamma_i L_1}$ ,
2. the phase accumulation of  $\nu_i$  from the source to  $L_1$ ,  $e^{-iE_i L_1}$ ,
3. the decay of  $\nu_i$  and appearance of  $\nu_j$ ,  $\sqrt{\Gamma_{ij} W_{ij}}$ ,
4. for unstable  $\nu_j$ , survival until Earth,  $e^{-\frac{1}{2}\Gamma_j(L-L_1)}$ ,
5. and the phase accumulation of  $\nu_j$  until Earth,  $e^{-iE_f(L-L_1)}$ .

After removing phases which are irrelevant for averaging,

$$\bar{\mathcal{A}}_{ij}^{\text{reg}} = e^{-\frac{1}{2}\Gamma_i L_1} \sqrt{\Gamma_{ij} W_{ij}} e^{-\frac{1}{2}\Gamma_j(L-L_1)}$$

Note that  $\Gamma_1 = 0$

For  $\nu \rightarrow \nu$  and  $\nu \rightarrow \bar{\nu}$ :

$$\Gamma_{ij} W_{ij} = \frac{m_i m_j}{16\pi E_i^2} \left[ g_{ij}^2 \left( \frac{1}{x_{ij}} + x_{ij} + 2 \right) + g'_{ij}{}^2 \left( \frac{1}{x_{ij}} + x_{ij} - 2 \right) \right]$$

Significantly simpler than expected or in either  $\nu \rightarrow \nu$  or  $\nu \rightarrow \bar{\nu}$  cases!

## Regeneration Component: Integrals

$$\bar{P}_{ij}^{\text{reg}}(E_f, L) = \frac{1}{\Phi_i^S(E_f)} \int_0^L dL_1 \int_{E_f}^{x_{ij}^2 E_f} dE_i |\bar{\mathcal{A}}_{ij}^{\text{reg}}(E_i, E_f, L, L_1)|^2 \Phi_i^S(E_i)$$

$$x_{ij} \equiv m_i/m_j$$

## Regeneration Component: Integrals

$$\bar{P}_{ij}^{\text{reg}}(E_f, L) = \frac{1}{\Phi_i^S(E_f)} \int_0^L dL_1 \int_{E_f}^{x_{ij}^2 E_f} dE_i |\bar{\mathcal{A}}_{ij}^{\text{reg}}(E_i, E_f, L, L_1)|^2 \Phi_i^S(E_i)$$

$$x_{ij} \equiv m_i/m_j$$

After the  $L_1$  integral,

$$\bar{P}_{ij}^{\text{reg}}(E_f, L) = \frac{1}{\Phi_i^S(E_f)} \int_{E_f}^{x_{ij}^2 E_f} dE_i \frac{\Gamma_{ij} W_{ij}}{\Gamma_i - \Gamma_j} \left[ 1 - e^{-(\Gamma_i - \Gamma_j)L} \right] \Phi_i^S(E_i)$$

## Regeneration Component: Integrals

$$\bar{P}_{ij}^{\text{reg}}(E_f, L) = \frac{1}{\Phi_i^S(E_f)} \int_0^L dL_1 \int_{E_f}^{x_{ij}^2 E_f} dE_i |\bar{\mathcal{A}}_{ij}^{\text{reg}}(E_i, E_f, L, L_1)|^2 \Phi_i^S(E_i)$$

$$x_{ij} \equiv m_i/m_j$$

After the  $L_1$  integral,

$$\bar{P}_{ij}^{\text{reg}}(E_f, L) = \frac{1}{\Phi_i^S(E_f)} \int_{E_f}^{x_{ij}^2 E_f} dE_i \frac{\Gamma_{ij} W_{ij}}{\Gamma_i - \Gamma_j} \left[ 1 - e^{-(\Gamma_i - \Gamma_j)L} \right] \Phi_i^S(E_i)$$

After the  $E_i$  integral,

$$\bar{P}_{ij}^{\text{reg}}(E_f, L) = \frac{z(x)}{\gamma y(x)} \left\{ 1 - \frac{1}{x^{2\gamma}} + \gamma T^{-\gamma} \left[ \Gamma(\gamma, T) - \Gamma\left(\gamma, \frac{T}{x^2}\right) \right] \right\}$$

$$T \equiv \frac{m_i m_j L}{16\pi E_f} y(x)$$

$$y(x) = g_{ij}^2 [f(x) + k(x)] + g'_{ij}{}^2 [h(x) + k(x)]$$

$$z(x) = g_{ij}^2 \left(\frac{1}{x} + x + 2\right) + g'_{ij}{}^2 \left(\frac{1}{x} + x - 2\right)$$

## Regeneration Component: Analytic Limits

- ▶ Verify that as  $E_f \rightarrow \infty$ ,  $\bar{P}_{ij}^{\text{reg}} \rightarrow 0$  as expected

SM

## Regeneration Component: Analytic Limits

- ▶ Verify that as  $E_f \rightarrow \infty$ ,  $\bar{P}_{ij}^{\text{reg}} \rightarrow 0$  as expected
- ▶ As  $E_f \rightarrow 0$ :

SM

Full decay

$$\lim_{E_f \rightarrow 0} \bar{P}_{ij}^{\text{reg}}(E_f, L) = \frac{z(x)}{\gamma y(x)} \left( 1 - \frac{1}{x^{2\gamma}} \right)$$

## Regeneration Component: Analytic Limits

- ▶ Verify that as  $E_f \rightarrow \infty$ ,  $\bar{P}_{ij}^{\text{reg}} \rightarrow 0$  as expected

SM

- ▶ As  $E_f \rightarrow 0$ :

Full decay

$$\lim_{E_f \rightarrow 0} \bar{P}_{ij}^{\text{reg}}(E_f, L) = \frac{z(x)}{\gamma y(x)} \left( 1 - \frac{1}{x^{2\gamma}} \right)$$

- ▶ As  $m_1 \rightarrow \infty$

Degenerate masses

$$\lim_{\substack{E_f \rightarrow 0 \\ m_1 \rightarrow \infty}} \bar{P}_{ij}^{\text{reg}}(E_f, L) = 1$$

## Regeneration Component: Analytic Limits

- ▶ Verify that as  $E_f \rightarrow \infty$ ,  $\bar{P}_{ij}^{\text{reg}} \rightarrow 0$  as expected

SM

- ▶ As  $E_f \rightarrow 0$ :

Full decay

$$\lim_{E_f \rightarrow 0} \bar{P}_{ij}^{\text{reg}}(E_f, L) = \frac{z(x)}{\gamma y(x)} \left(1 - \frac{1}{x^{2\gamma}}\right)$$

- ▶ As  $m_1 \rightarrow \infty$

Degenerate masses

$$\lim_{\substack{E_f \rightarrow 0 \\ m_1 \rightarrow \infty}} \bar{P}_{ij}^{\text{reg}}(E_f, L) = 1$$

- ▶ As  $m_1 \rightarrow 0$

Depends on nature of interaction!

$$\lim_{\substack{E_f \rightarrow 0 \\ m_1 \rightarrow 0}} \bar{P}_{ij}^{\text{reg}}(E_f, L) = \frac{1}{\gamma}$$



# Cosmology

Two main changes:

1.  $L \rightarrow L(z_a, z_b)$

$$L(z_a, z_b) = L_H \int_{z_a}^{z_b} \frac{dz'}{(1+z')h(z')}$$

$$h(z) \equiv \sqrt{(1+z)^3 \Omega_m + \Omega_\Lambda}$$

2.  $E \rightarrow E(1+z)$

▶  $\Gamma \rightarrow \Gamma/(1+z)$

▶  $\Gamma W \rightarrow \Gamma W/(1+z)^2$

▶  $\exp(-\Gamma L) \rightarrow \exp[-L_H \int_{z_a}^{z_b} \frac{dz'}{(1+z')^2 h(z')}]$

By definition, the “probability” now is:

$$\bar{P}_{ij}^{\text{SM}} = \delta_{ij}(1+z)^{-\gamma}$$

Affects both depletion and regeneration!

# Regeneration Component: Analytic Limits with Cosmology

► As  $m_1 \rightarrow \infty$

$$\lim_{\substack{E_f \rightarrow 0 \\ m_1 \rightarrow \infty}} \bar{P}_{ij}^{\text{reg}}(E_f, L) = (1+z)^{-2\gamma}$$

$$\lim_{\substack{E_f \rightarrow 0 \\ m_1 \rightarrow \infty}} \bar{P}_{\alpha\beta}^{\text{vis}} = (1+z)^{-\gamma} [ |U_{\alpha 1}|^2 |U_{\beta 1}|^2 + |U_{\alpha 2}|^2 |U_{\beta 2}|^2 + (1+z)^{-\gamma} |U_{\alpha 3}|^2 |U_{\beta 1}|^2 ]$$

# Regeneration Component: Analytic Limits with Cosmology

- ▶ As  $m_1 \rightarrow \infty$

$$\lim_{\substack{E_f \rightarrow 0 \\ m_1 \rightarrow \infty}} \bar{P}_{ij}^{\text{reg}}(E_f, L) = (1+z)^{-2\gamma}$$

$$\lim_{\substack{E_f \rightarrow 0 \\ m_1 \rightarrow \infty}} \bar{P}_{\alpha\beta}^{\text{vis}} = (1+z)^{-\gamma} [ |U_{\alpha 1}|^2 |U_{\beta 1}|^2 + |U_{\alpha 2}|^2 |U_{\beta 2}|^2 + (1+z)^{-\gamma} |U_{\alpha 3}|^2 |U_{\beta 1}|^2 ]$$

- ▶ As  $m_1 \rightarrow 0$

$$\lim_{\substack{E_f \rightarrow 0 \\ m_1 \rightarrow 0}} \bar{P}_{ij}^{\text{reg}}(E_f, L) = \frac{(1+z)^{-2\gamma}}{\gamma}$$

$$\lim_{\substack{E_f \rightarrow 0 \\ m_1 \rightarrow 0}} \bar{P}_{\alpha\beta}^{\text{vis}} = (1+z)^{-\gamma} \left[ |U_{\alpha 1}|^2 |U_{\beta 1}|^2 + |U_{\alpha 2}|^2 |U_{\beta 2}|^2 + \frac{(1+z)^{-\gamma}}{\gamma} |U_{\alpha 3}|^2 |U_{\beta 1}|^2 \right]$$

## Regeneration Component: Multiple Decays

If  $g_{32}$  and  $g_{21}$  are nonzero, there is another way to get from  $\nu_3 \rightarrow \nu_1$ :

1. Decay from  $\nu_3 \rightarrow \nu_2$  at  $z_1$  from  $E_i \rightarrow E_{\text{int}}$
2. Decay from  $\nu_2 \rightarrow \nu_1$  at  $z_2$  from  $E_{\text{int}} \rightarrow E_f$

## Regeneration Component: Multiple Decays

If  $g_{32}$  and  $g_{21}$  are nonzero, there is another way to get from  $\nu_3 \rightarrow \nu_1$ :

1. Decay from  $\nu_3 \rightarrow \nu_2$  at  $z_1$  from  $E_i \rightarrow E_{\text{int}}$
2. Decay from  $\nu_2 \rightarrow \nu_1$  at  $z_2$  from  $E_{\text{int}} \rightarrow E_f$

$$\bar{P}_{31}^{\text{reg},2} = \frac{L_H^2}{\Phi_i^S(E_f)} \int_0^z dz_2 \int_{z_2}^z dz_1 \int_{E_f(1+z_2)}^{E_f x_{21}^2(1+z_2)} dE_{\text{int}} \int_{E_{\text{int}}(1+z_1)}^{E_{\text{int}} x_{32}^2(1+z_1)} dE_i$$
$$\frac{\Gamma_{32} W_{32} \Gamma_{21} W_{21} e^{-\Gamma_3 L(z_1, z) - \Gamma_2 L(z_2, z_1)}}{(1+z_1)^2 h(z_1) (1+z_2)^2 h(z_2)} \Phi_i^S(E_i(1+z_1))$$

Where

$$\Gamma_3 \rightarrow \Gamma_3(E_i) \quad \text{and} \quad \Gamma_2 \rightarrow \Gamma_2(E_{\text{int}})$$
$$\Gamma_{32} W_{32} \rightarrow \Gamma_{32} W_{32}(E_i, E_{\text{int}}) \quad \text{and} \quad \Gamma_{21} W_{21} \rightarrow \Gamma_{21} W_{21}(E_{\text{int}}, E_f)$$

## Regeneration Component: Coherency

Now coherency is a problem:

1. If the decay happens before they decohere/separate the interference term should be included
2. If the decay happens after they decohere/separate it shouldn't be included

In principle there should be a  $\exp[-(L/L_{\text{COH}})^2]$  type term

A. de Gouvea, V. De Romeri, C. Ternes [2005.03022](#)

We ignore this

1. For partial decay this effect is negligible, only matters for full decay
2. Production regions could be anywhere in the  $10^9 - 10^{17}$  cm region

Backup slide [47](#) has a plot

## Results: SM

Assuming an initial flavor ratio of  $(1 : 2 : 0)$  (pion decay):

$$(1 : 2 : 0) \rightarrow (1 : 1 : 1)$$

This is a result of several coincidences

- ▶ The neutrino energies from pion decay are all roughly the same
- ▶ The mixing matrix is approximately tri-bimaximal
  - ▶ The deviations we know that exist from TBM don't affect this flavor ratio much

Expect the same flux of each flavor

See slide [43](#) for deviations from this

## Results: Benchmark Values

- ▶ Include both scalar and pseudo-scalar interactions with equal couplings
- ▶ Include both  $\nu \rightarrow \nu$  and  $\nu \rightarrow \bar{\nu}$  channels
- ▶ Assume a power law spectrum  $\Phi_{\alpha}^S(E_i) = \Phi_{\alpha,0}^S E^{-\gamma}$  with  $\gamma = 2$
- ▶ Assume  $m_1 = 0$  eV

Anything less than  $\sim 10^{-3}$  eV is equivalent to zero

- ▶ Assume they are all coming from  $z = 1$

See backup slide 48


- ▶ Assume all six couplings are  $10^{-6}$

Puts the partial decay feature in IceCube's view


Turn all the knobs!




# Results: Invisible Decay




# Results: Visible Decay




# Results: Visible Decay: Neutrino mass scale




# Results: Visible Decay: Spectral Index



# Results: Visible Decay: Scalar vs. Pseudo-scalar




# Results: Visible Decay: Neutrinos vs. Anti-neutrinos



# Summary of Parameters

1.  $\gamma$ : harder spectra  $\Rightarrow$  large regeneration component
2.  $m_1$ : higher mass scale  $\gtrsim 0.1$  eV  $\Rightarrow$  large regeneration component
3.  $g_{ij}$ : different features depending on the texture
4. Redshift evolution  $\Rightarrow$  small effect
5. Scalar/Pseudo-scalar  $\Rightarrow$  small effect
6.  $\nu \rightarrow \nu, \nu \rightarrow \bar{\nu} \Rightarrow$  small effect

# Results: Visible Decay: Flavors






## IceCube Measures:


- ▶ Energy
- ▶ Direction
- ▶ Flavor(ish)

# IceCube is Great for This!

IceCube has measured the extragalactic high energy (100 TeV - 1 PeV) flux!



IC ICRC 2017







Galactic component is negligible  
PBD, D. Marfatia, T. Weiler [1703.09721](#)

# IceCube Can Detect Flavor (sort of)

High enough energy to be well about  $\nu_\tau$  threshold

Y. Jeong, M. Reno [1007.1966](#)




Cascade topology  
 $\nu_e$  CC,  $\nu_\tau$  CC\*, NC

Track topology  
 $\nu_\mu$  CC,  $\nu_\tau$  CC  $\rightarrow \tau \rightarrow \mu + 2\nu$

# Tension

$$\Phi(E) = \Phi_0 E^{-\gamma}$$



$$\Delta\gamma = +0.54$$

“The p-value for obtaining the combined fit result and the result reported here from an unbroken powerlaw flux is  $3.3\sigma$ , and is therefore in significant **tension**.”

IC 1607.08006

“This [cascade] fit [is] in **tension** with previous results based on through-going muons”

IC 1808.07629

## Conventional Wisdom

- ▶ High energy neutrinos are produced from full  $\pi$  decay
- ▶ Flavor ratio at source of 1:2:0 converts to 1:1:1\* at Earth
- ▶ All neutrinos have the same energy<sup>†</sup>

\*the fact that this ratio is 1:1:1 is coincidental not fundamental

<sup>†</sup>also a coincidence; kinematic corrections are small

## Conventional Wisdom

- ▶ High energy neutrinos are produced from full  $\pi$  decay
- ▶ Flavor ratio at source of 1:2:0 converts to 1:1:1\* at Earth
- ▶ All neutrinos have the same energy<sup>†</sup>

Some of these *must* be incorrect.

\*the fact that this ratio is 1:1:1 is coincidental not fundamental


<sup>†</sup>also a coincidence; kinematic corrections are small

Need a phenomenon that non-trivially depends on  
**energy** and **flavor** at the same time

# Muon Cooling

$$\pi \rightarrow \nu_\mu + \mu$$

$$\mu \rightarrow \nu_\mu + \nu_e + e$$




- ▶ E.g. synchrotron
- ▶ More  $\nu_\mu$  at high energy
- ▶  $E_b$  determined by  $B$  field



# Muon Cooling

$$\pi \rightarrow \nu_\mu + \mu$$

$$\mu \rightarrow \nu_\mu + \nu_e + e$$



- ▶ E.g. synchrotron
- ▶ More  $\nu_\mu$  at high energy
- ▶  $E_b$  determined by  $B$  field
  
- ▶ This doesn't work at all!
- ▶ Oscillations kill this
  - ▶  $\mu - \tau$  symmetry
- ▶  $\max \Delta\gamma \simeq 0.2$

## Other Options

Neutron decay:  $n \rightarrow p + e + \bar{\nu}_e$

- ▶ Produces extra  $\nu_e$ 's
- ▶ Produced with pions in  $p\gamma$  interactions
- ▶ Also come from photodissociation of heavy ions

A. Palladino [1902.08630](#)

L. Anchordoqui [1411.6457](#)

## Other Options

Neutron decay:  $n \rightarrow p + e + \bar{\nu}_e$

- ▶ Produces extra  $\nu_e$ 's
- ▶ Produced with pions in  $p\gamma$  interactions
- ▶ Also come from photodissociation of heavy ions

A. Palladino [1902.08630](#)

L. Anchordoqui [1411.6457](#)

**But**

- ▶ Neutrino energies are  $\sim 2$ -3 orders of magnitude less for  $p\gamma$
- ▶ Neutrino flux from heavy ions is also suppressed

D. Biehl, et al. [1705.08909](#)

X. Rodrigues, et al. [1711.02091](#)

# New Physics!

We need a stronger effect, so we look to new physics.

- ▶ NSI with ultra-light mediators ( $m \ll 1$  eV)

weak

A. Joshipura, S. Mohanty [hep-ph/0310210](#)

M. Bustamante, S. Agarwalla [1808.02042](#)

- ▶ Pseudo-dirac neutrinos

weak

L. Wolfenstein [Nucl. Phys. B186, 147 \(1981\)](#)


S. Pakvasa, A. Joshipura, S. Mohanty [1209.5630](#)

- ▶ Electrophilic dark matter decay


- ▶ Neutrino decay

strong,  $\sim 3.3 \sigma$


strong but CMB



# Spectral Indices at IceCube



# Spectral Indices at IceCube



## Some Track to Cascade with Decay Observations

- ▶ Decay usually hardens the spectrum
  - ▶ Only  $\bar{P}_{\mu e}^{\text{vis}} > \bar{P}_{\mu e}^{\text{SM}}$  for  $m_1 \sim 0$  and  $\gamma \sim 2$
  - ▶ While  $\bar{P}_{\tau\beta}^{\text{vis}} > \bar{P}_{\tau\beta}^{\text{SM}}$ , no  $\nu_\tau$ 's are produced at the sources

See backup slide 53

- ▶ The effect is larger for tracks than cascades


$\max \Delta\gamma$	$g_{21}^{(l)}$	$g_{31}^{(l)}$	$g_{32}^{(l)}$	All
Invisible	0.006	0.200	0.200	0.438
Visible	0.042	0.227	0.172	0.400

$$\min \Delta\gamma = -0.01$$

$$\Delta\gamma \equiv \gamma_c - \gamma_t$$

- ▶ This is the same direction of the IceCube data!
- ▶ The other sign (cascades harder than tracks) requires the inverted ordering

# Preferred Region: Visible





# Uncertainties

or “How to muck it all up with astrophysics”

## What doesn't work:

- ▶ Multiple classes of sources with different spectra
- ▶  $pp$  vs.  $p\gamma$  sources
- ▶ Different redshift evolution  $\Rightarrow$  shift the  $g_{ij}$
- ▶ Neutron decay sources
- ▶ Varying the oscillation parameters
- ▶ IceCube track or cascade normalization

## What could work: (other than neutrino decay)

- ▶ Muon damped  $\Rightarrow \Delta\gamma \sim 0.2$
- ▶ Track and cascade spectra are fit over slightly different energy ranges  $\Rightarrow$  broken power law can help
- ▶ Energy misreconstruction (tracks could be susceptible to this)
- ▶ Dark matter?


## Key Points

- ▶ Neutrino decay pheno can be probed in a broad range of experimental regions
- ▶ Visible neutrino decay contains depletion and regeneration terms
- ▶ Varying the initial spectrum, mass scale, and couplings leads to a range of spectra
- ▶ IceCube is uniquely sensitive to this
- ▶ IceCube's track/cascade spectrum can be well described by neutrino decay
- ▶ Neutrino decay within the NO predicts the same kind of tension that IceCube sees


Thanks!

# Backups


# The effect of decoherence




# Visible Decay for Different Redshift Evolution Functions



# IceCube Track and Cascade Spectral Index Difference: Invisible Decay




# IceCube Track and Cascade Spectral Index Difference: Visible Decay






# Preferred Region: Invisible



# Analytic Validation



# Results: Visible Decay: Flavors with $\nu_\tau$

