

Inclusive jets and dynamical mass effects

Andrea Signori

University of Pavia and Jefferson Lab

Jets for 3D imaging workshop

November 24, 2020



Outline

Introduction

Inclusive jets

Hadronization and mass generation

Semi-inclusive processes



Outline

Introduction

Inclusive jets

Hadronization and mass generation

Semi-inclusive processes



Some references:

A [selection of references](#) related to the topics discussed in this talk:

- ▶ On the connection between quark propagation and hadronization

A. Accardi, [A. Signori](#) - 2005.11310 - EPJC

- ▶ Quark fragmentation as a probe of dynamical mass generation

A. Accardi, [A. Signori](#) - 1903.04458 - PLB

Some references:

A [selection of references](#) related to the topics discussed in this talk:

- ▶ On the connection between quark propagation and hadronization
A. Accardi, [A. Signori](#) - 2005.11310 - EPJC
- ▶ Quark fragmentation as a probe of dynamical mass generation
A. Accardi, [A. Signori](#) - 1903.04458 - PLB
- ▶ Accessing the nucleon transverse structure in deep-inelastic scattering
A. Accardi, A. Bacchetta - 1706.02000 - PLB

Some references:

A [selection of references](#) related to the topics discussed in this talk:

- ▶ On the connection between quark propagation and hadronization
A. Accardi, [A. Signori](#) - 2005.11310 - EPJC
- ▶ Quark fragmentation as a probe of dynamical mass generation
A. Accardi, [A. Signori](#) - 1903.04458 - PLB
- ▶ Accessing the nucleon transverse structure in deep-inelastic scattering
A. Accardi, A. Bacchetta - 1706.02000 - PLB
- ▶ Collinear factorization for deep inelastic scattering structure functions at large Bjorken x_B
A. Accardi, J.W. Qiu - 0805.1496 - PRD

Hadron physics

Hadron physics \leftrightarrow (non-)perturbative QCD

The two faces of confinement:

- ▶ hadron structure (tomography): hadron \rightarrow quark/gluon transition
- ▶ hadronization: hadron \leftarrow quark/gluon transition

Hadron physics



Hadron physics \leftrightarrow (non-)perturbative QCD

The two faces of confinement:

- ▶ **hadron structure (tomography):** hadron \rightarrow quark/gluon transition
- ▶ **hadronization:** hadron \leftarrow quark/gluon transition

Motivations:

- ▶ **conceptual:** understand QCD, in particular confinement, dynamical breaking of chiral symmetry
- ▶ **practical:** reactions involving hadrons in the initial and/or final state




Fragmentation functions

Hadronization: dynamical generation of hadronic properties from quarks/gluons
→ fundamental topic

It follows any QCD hard scattering event and populates the final states with hadrons.

Maps of hadronization in momentum space: fragmentation functions (FFs)




Fragmentation functions

Hadronization: dynamical generation of hadronic properties from quarks/gluons
→ fundamental topic

It follows any QCD hard scattering event and populates the final states with hadrons.

Maps of hadronization in momentum space: fragmentation functions (FFs)




Momentum distributions:

Fragmentation functions

Hadronization: dynamical generation of hadronic properties from quarks/gluons
→ fundamental topic

It follows any QCD hard scattering event and populates the final states with hadrons.

Maps of hadronization in momentum space: fragmentation functions (FFs)



Momentum distributions:


$D_1^{a \rightarrow h}(z)$: collinear FF

Fragmentation functions

Hadronization: dynamical generation of hadronic properties from quarks/gluons
→ fundamental topic

It follows any QCD hard scattering event and populates the final states with hadrons.

Maps of hadronization in momentum space: fragmentation functions (FFs)



Momentum distributions:

$D_1^{a \rightarrow h}(z)$: collinear FF


$D_1^{a \rightarrow h}(z, P_T^2)$: TMD FF

Fragmentation functions

Hadronization: dynamical generation of hadronic properties from quarks/gluons
→ fundamental topic

It follows any QCD hard scattering event and populates the final states with hadrons.

Maps of hadronization in momentum space: fragmentation functions (FFs)



Momentum distributions:

$D_1^{a \rightarrow h}(z)$: collinear FF

$D_1^{a \rightarrow h}(z, P_T^2)$: TMD FF


$J_a(s)$: inclusive jet function

Fragmentation functions

Hadronization: dynamical generation of hadronic properties from quarks/gluons
→ fundamental topic

It follows any QCD hard scattering event and populates the final states with hadrons.

Maps of hadronization in momentum space: **fragmentation functions (FFs)**



Momentum distributions:

$D_1^{a \rightarrow h}(z)$: collinear FF

$D_1^{a \rightarrow h}(z, P_T^2)$: TMD FF

$J_a(s)$: inclusive jet function


$\mathcal{G}^{a \rightarrow h}(s, z)$: fragmenting jet function

Fragmentation functions

Hadronization: dynamical generation of hadronic properties from quarks/gluons
→ fundamental topic

It follows any QCD hard scattering event and populates the final states with hadrons.

Maps of hadronization in momentum space: **fragmentation functions (FFs)**



Momentum distributions:

$D_1^{a \rightarrow h}(z)$: collinear FF

$D_1^{a \rightarrow h}(z, P_T^2)$: TMD FF

$J_a(s)$: inclusive jet function

$\mathcal{G}^{a \rightarrow h}(s, z)$: fragmenting jet function

...

The generation of mass in QCD

What generates the masses of partons and hadrons?

- ▶ Higgs mechanism, only quark masses: $m_q \sim \text{MeV} \ll M_{p/n} \sim 1 \text{ GeV}$



The generation of mass in QCD

What generates the masses of partons and hadrons?

- ▶ Higgs mechanism, only quark masses: $m_q \sim \text{MeV} \ll M_{p/n} \sim 1 \text{ GeV}$
- ▶ the rest comes from the **dynamics of QCD** \rightarrow **dynamical mass**




The generation of mass in QCD

What generates the masses of partons and hadrons?

- ▶ Higgs mechanism, only quark masses: $m_q \sim \text{MeV} \ll M_{p/n} \sim 1 \text{ GeV}$
- ▶ the rest comes from the **dynamics of QCD** \rightarrow **dynamical mass**

The dynamical generation of mass in QCD can be addressed in different ways:

- ▶ gap equation
e.g. in the NJL model of QCD: $M_q = m_q - 4G_\pi \langle \bar{q}q \rangle \gg m_q$




The generation of mass in QCD

What generates the masses of partons and hadrons?

- ▶ Higgs mechanism, only quark masses: $m_q \sim \text{MeV} \ll M_{p/n} \sim 1 \text{ GeV}$
- ▶ the rest comes from the **dynamics of QCD** \rightarrow **dynamical mass**

The dynamical generation of mass in QCD can be addressed in different ways:

- ▶ gap equation
e.g. in the NJL model of QCD: $M_q = m_q - 4G_\pi \langle \bar{q}q \rangle \gg m_q$
- ▶ Energy Momentum Tensor \rightarrow hadron mass decomposition




The generation of mass in QCD

What generates the masses of partons and hadrons?

- ▶ Higgs mechanism, only quark masses: $m_q \sim \text{MeV} \ll M_{p/n} \sim 1 \text{ GeV}$
- ▶ the rest comes from the **dynamics of QCD** \rightarrow **dynamical mass**

The dynamical generation of mass in QCD can be addressed in different ways:

- ▶ gap equation
e.g. in the NJL model of QCD: $M_q = m_q - 4G_\pi \langle \bar{q}q \rangle \gg m_q$
- ▶ Energy Momentum Tensor \rightarrow hadron mass decomposition
- ▶ “mass sum rule” for **fragmentation functions** - new and observable!



Outline

Introduction


Inclusive jets

Hadronization and mass generation

Semi-inclusive processes




Inclusive jets



Inclusive jet function $J_i(s)$:
sensitive to the jet virtuality s

Inclusive jets




Inclusive jet function $J_i(s)$:
sensitive to the jet virtuality s

“Composition” of the jet:

- ▶ perturbative radiation (large s , wiggles)

Inclusive jets




Inclusive jet function $J_i(s)$:
sensitive to the jet virtuality s

“Composition” of the jet:

- ▶ perturbative radiation (large s , wiggles)
- ▶ non-perturbative radiation (low s , dashed lines)

Inclusive jets




Inclusive jet function $J_i(s)$:
sensitive to the jet virtuality s

“Composition” of the jet:

- ▶ perturbative radiation (large s , wiggles)
- ▶ non-perturbative radiation (low s , dashed lines)
- ▶ non-perturbative quark propagation (yellow blob)

Inclusive jets




Inclusive jet function $J_i(s)$:
sensitive to the jet virtuality s

“Composition” of the jet:

- ▶ perturbative radiation (large s , wiggles)
- ▶ non-perturbative radiation (low s , dashed lines)
- ▶ non-perturbative quark propagation (yellow blob) ← today's focus

Inclusive jets and FJFs


Procura, Stewart 0911.4980 - PRD
Jain, Procura, Waalewijn 1101.4953 - JHEP




Inclusive jet function $J_i(s)$:
sensitive to the jet virtuality s

Inclusive jets and FJFs

Procura, Stewart 0911.4980 - PRD
Jain, Procura, Waalewijn 1101.4953 - JHEP




Inclusive jet function $J_i(s)$:
sensitive to the jet virtuality s



Fragmenting jet function (FJF) $\mathcal{G}^{i \rightarrow h}(s, z)$:
sensitive to jet virtuality s
and hadron momentum fraction z
(less inclusive)


Inclusive jets and FJFs

Procura, Stewart 0911.4980 - PRD
Jain, Procura, Waalewijn 1101.4953 - JHEP



Inclusive jets and FJFs

Procura, Stewart 0911.4980 - PRD
Jain, Procura, Waalewijn 1101.4953 - JHEP




$$J_i(s) = \frac{1}{2(2\pi)^3} \sum_h \int dz z \mathcal{G}^{i \rightarrow h}(s, z)$$

Connection between the unpolarized jet function and FJFs :
jet as the “inclusive” limit of the in-jet fragmentation


FJFs and 1h-FFs

Procura, Stewart 0911.4980 - PRD
Jain, Procura, Waalewijn 1101.4953 - JHEP



FJFs and 1h-FFs

Procura, Stewart 0911.4980 - PRD
Jain, Procura, Waalewijn 1101.4953 - JHEP



$$\mathcal{G}^{i \rightarrow h}(s, z) = \sum_j \mathcal{J}_{ij}(s, z) \otimes D_1^{j \rightarrow h}(z) + \mathcal{O}(\Lambda_{qcd}^2 s^{-1})$$


Large- s expansion of the unpolarized FJF \mathcal{G}
on the single-hadron collinear FF D_1

Inclusive jets and 1h-FFs

Accardi, Signori 1903.04458 - PLB

Accardi, Signori 2005.11310 - EPJC


In a similar way, one can interpret the gauge-invariant quark propagator (aka *inclusive jet correlator*) as the *inclusive limit* of the *1h-fragmentation correlator*:



Inclusive jets and 1h-FFs

Accardi, Signori 1903.04458 - PLB
Accardi, Signori 2005.11310 - EPJC

In a similar way, one can interpret the gauge-invariant quark propagator (aka *inclusive jet correlator*) as the *inclusive limit* of the *1h-fragmentation correlator*:




$$k^\mu \Xi^i(k) = \sum_{h, S_h} \int \frac{d^4 P_h}{(2\pi)^3} \delta(P_h^2 - M_h^2) P_h^\mu \Delta^{i \rightarrow h}(k, P_h, S_h)$$

Inclusive jets and 1h-FFs

Accardi, Signori 1903.04458 - PLB
Accardi, Signori 2005.11310 - EPJC

In a similar way, one can interpret the gauge-invariant quark propagator (aka *inclusive jet correlator*) as the *inclusive limit* of the *1h-fragmentation correlator*:




$$k^\mu \Xi^i(k) = \sum_{h, S_h} \int \frac{d^4 P_h}{(2\pi)^3} \delta(P_h^2 - M_h^2) P_h^\mu \Delta^{i \rightarrow h}(k, P_h, S_h)$$

Unpolarized Dirac projection: momentum sum rule for $D_1(z)$

Inclusive jets and 1h-FFs

Accardi, Signori 1903.04458 - PLB
Accardi, Signori 2005.11310 - EPJC

In a similar way, one can interpret the gauge-invariant quark propagator (aka **inclusive jet correlator**) as the **inclusive limit** of the **1h-fragmentation correlator**:




$$k^\mu \Xi^i(k) = \sum_{h, S_h} \int \frac{d^4 P_h}{(2\pi)^3} \delta(P_h^2 - M_h^2) P_h^\mu \Delta^{i \rightarrow h}(k, P_h, S_h)$$

Unpolarized Dirac projection: momentum sum rule for $D_1(z)$
corresponds to sum rule between $J_i(s)$ and $\mathcal{G}^{i \rightarrow h}(s, z)$
at the **leading order** and **integrated** over the jet virtuality s .

Inclusive jets and 1h-FFs

Accardi, Signori 1903.04458 - PLB
Accardi, Signori 2005.11310 - EPJC

In a similar way, one can interpret the gauge-invariant quark propagator (aka **inclusive jet correlator**) as the **inclusive limit** of the **1h-fragmentation correlator**:



$$k^\mu \Xi^i(k) = \sum_{h, S_h} \int \frac{d^4 P_h}{(2\pi)^3} \delta(P_h^2 - M_h^2) P_h^\mu \Delta^{i \rightarrow h}(k, P_h, S_h)$$

Unpolarized Dirac projection: momentum sum rule for $D_1(z)$
corresponds to sum rule between $J_i(s)$ and $\mathcal{G}^{i \rightarrow h}(s, z)$
at the **leading order** and **integrated** over the jet virtuality s .

Novelty: consider **all the Dirac projections up to twist 3**
and connect **1h-FFs** to the **non-perturbative properties** of the **quark propagator**.

Outline

Introduction

Inclusive jets


Hadronization and mass generation

Semi-inclusive processes




Quark 1h-FFs

$$\Delta_{ij}(k, P_h, S_h) = \int \frac{d^4\xi}{(2\pi)^4} e^{ikx} \frac{\text{Tr}_c}{N_c} \langle \Omega | \hat{T} W_1(\infty, \xi) \psi_i(\xi) a^\dagger a \bar{\psi}_j(0) W_2(0, \infty) | \Omega \rangle$$



Quark 1h-FFs

$$\Delta_{ij}(k, P_h, S_h) = \int \frac{d^4\xi}{(2\pi)^4} e^{ikx} \frac{\text{Tr}_c}{N_c} \langle \Omega | \hat{T} W_1(\infty, \xi) \psi_i(\xi) a^\dagger a \bar{\psi}_j(0) W_2(0, \infty) | \Omega \rangle$$



		quark pol.		
		U	L	T
hadron pol.	U	D_1		H_1^\perp
	L		G_{1L}	H_{1L}^\perp
	T	D_{1T}^\perp	G_{1T}	H_1, H_{1T}^\perp

8 (TMD) fragmentation functions at leading twist

Quark higher twist 1h-FFs

Twist 3 transverse momentum dependent FFs $\mathcal{D}_{\dots}^{a \rightarrow h}(z, P_{h\perp}^2)$
for a quark hadronizing into a spin 1/2 hadron

		quark pol.		
hadron pol.		U	L	T
	U	D^\perp	G^\perp	E, H
	L	D_L^\perp	G_L^\perp	H_L, E_L
	T	D_T, D_T^\perp	G_T, G_T^\perp	$H_T, H_T^\perp, E_T, E_T^\perp$

Quark higher twist 1h-FFs

Twist 3 transverse momentum dependent FFs $\mathcal{D}_{\dots}^{a \rightarrow h}(z, P_{h\perp}^2)$
for a quark hadronizing into a spin 1/2 hadron

		quark pol.		
hadron pol.		U	L	T
	U	D^\perp	G^\perp	E, H
	L	D_L^\perp	G_L^\perp	H_L, E_L
	T	D_T, D_T^\perp	G_T, G_T^\perp	$H_T, H_T^\perp, E_T, E_T^\perp$

Black and magenta: survive transverse momentum integration

Red and magenta: T-odd

Blue: T-even, w/o collinear counterpart

Quark propagator

Källen-Lehman representation in terms of **spectral functions** $\rho_{1,3}$:

$$\Xi(k) \rightarrow S_F(k) = \int \frac{d\mu^2}{(2\pi)^4} \{ \not{k} \rho_3(\mu^2) + \sqrt{\mu^2} \rho_1(\mu^2) \mathbb{I} \} \frac{\theta(\mu^2)}{k^2 - \mu^2 + i\epsilon}$$

Quark propagator

Källen-Lehman representation in terms of **spectral functions** $\rho_{1,3}$:

$$\Xi(k) \rightarrow S_F(k) = \int \frac{d\mu^2}{(2\pi)^4} \{ \not{k} \rho_3(\mu^2) + \sqrt{\mu^2} \rho_1(\mu^2) \mathbb{I} \} \frac{\theta(\mu^2)}{k^2 - \mu^2 + i\epsilon}$$

Projecting the operator sum rule between Ξ^i and $\Delta^{i \rightarrow h}$ one obtains:

Quark propagator

Källen-Lehman representation in terms of **spectral functions** $\rho_{1,3}$:

$$\Xi(k) \rightarrow S_F(k) = \int \frac{d\mu^2}{(2\pi)^4} \{ \not{k} \rho_3(\mu^2) + \sqrt{\mu^2} \rho_1(\mu^2) \mathbb{I} \} \frac{\theta(\mu^2)}{k^2 - \mu^2 + i\epsilon}$$

Projecting the operator sum rule between Ξ^i and $\Delta^{i \rightarrow h}$ one obtains:

twist 2 (γ^-):

$$\sum_h \int_0^1 dz z D_1^h(z) = \int_0^{+\infty} d\mu^2 \rho_3(\mu^2) \equiv 1 \quad (\text{QFT!})$$

Quark propagator

Källen-Lehman representation in terms of **spectral functions** $\rho_{1,3}$:

$$\Xi(k) \rightarrow S_F(k) = \int \frac{d\mu^2}{(2\pi)^4} \{ \not{k} \rho_3(\mu^2) + \sqrt{\mu^2} \rho_1(\mu^2) \mathbb{I} \} \frac{\theta(\mu^2)}{k^2 - \mu^2 + i\epsilon}$$

Projecting the operator sum rule between Ξ^i and $\Delta^{i \rightarrow h}$ one obtains:

twist 2 (γ^-):

$$\sum_h \int_0^1 dz z D_1^h(z) = \int_0^{+\infty} d\mu^2 \rho_3(\mu^2) \equiv 1 \quad (\text{QFT!})$$

twist 3 (\mathbb{I}):

$$\sum_h \int_0^1 dz M_h E^h(z) = \int_0^{+\infty} d\mu^2 \sqrt{\mu^2} \rho_1(\mu^2) \equiv M_j$$

Quark propagator

Källen-Lehman representation in terms of **spectral functions** $\rho_{1,3}$:

$$\Xi(k) \rightarrow S_F(k) = \int \frac{d\mu^2}{(2\pi)^4} \{ \not{k} \rho_3(\mu^2) + \sqrt{\mu^2} \rho_1(\mu^2) \mathbb{I} \} \frac{\theta(\mu^2)}{k^2 - \mu^2 + i\epsilon}$$

Projecting the operator sum rule between Ξ^i and $\Delta^{i \rightarrow h}$ one obtains:

twist 2 (γ^-):

$$\sum_h \int_0^1 dz z D_1^h(z) = \int_0^{+\infty} d\mu^2 \rho_3(\mu^2) \equiv 1 \quad (\text{QFT!})$$

twist 3 (\mathbb{I}):

$$\sum_h \int_0^1 dz M_h E^h(z) = \int_0^{+\infty} d\mu^2 \sqrt{\mu^2} \rho_1(\mu^2) \equiv M_j$$

The **non-perturbative** structure of the **jet** is **trivial** at **twist 2**, **but not at twist 3**!

Hadronization and mass generation

“Mass sum rule” for twist 3 E fragmentation function:

$$\sum_h \int dz M_h E^h(z) = M_j$$

quark/jet dynamical mass M_j as the
average of produced hadron masses
weighted by chiral-odd E FF



Hadronization and mass generation

“Mass sum rule” for twist 3 E fragmentation function:

$$\sum_h \int dz M_h E^h(z) = M_j$$

quark/jet dynamical mass M_j as the
average of produced hadron masses
weighted by chiral-odd E FF

QCD equations of motions: $E^h = \tilde{E}^h + \frac{m_q}{M_h} z D_1^h$

Hadronization and mass generation

“Mass sum rule” for twist 3 E fragmentation function:

$$\sum_h \int dz M_h E^h(z) = M_j$$

quark/jet dynamical mass M_j as the average of produced hadron masses weighted by chiral-odd E FF

QCD equations of motions: $E^h = \tilde{E}^h + \frac{m_q}{M_h} z D_1^h$

Wandzura-Wilczek (WW) approximation: $\tilde{E}^h = 0 \implies M_j = m_q$

Hadronization and mass generation

“Mass sum rule” for twist 3 E fragmentation function:

$$\sum_h \int dz M_h E^h(z) = M_j$$

quark/jet dynamical mass M_j as the average of produced hadron masses weighted by chiral-odd E FF

QCD equations of motions: $E^h = \tilde{E}^h + \frac{m_q}{M_h} z D_1^h$

Wandzura-Wilczek (WW) approximation: $\tilde{E}^h = 0 \implies M_j = m_q$

Full QCD: $M_j = m_q + m_q^{corr}$ (current and dynamical components), where

$$\sum_h \int dz M_h \tilde{E}^h(z) = M_j - m_q = m_q^{corr}$$

\tilde{E} and m_q^{corr} probe quark-gluon-quark $\sim \langle 0 | \bar{\psi} A \psi | 0 \rangle$ dynamical correlations

Hadronization and mass generation



“Mass sum rule” for twist 3 E fragmentation function:

$$\sum_h \int dz M_h E^h(z) = M_j$$

quark/jet dynamical mass M_j as the average of produced hadron masses weighted by chiral-odd E FF

Quark/jet mass M_j and its dynamical component m_q^{corr} :

- ▶ related to the “mass function” from the QCD gap equation
- ▶ gauge invariant and observable
- ▶ contribute to physical observables at twist 3 in the chiral-odd sector



Outline

Introduction

Inclusive jets

Hadronization and mass generation

Semi-inclusive processes




Semi-inclusive processes

- ▶ We can study the phenomenology of the dynamical mass in (semi-) inclusive hard processes
- ▶ interesting but challenging: **chiral-odd** sector at least at **twist-3**
- ▶ working in collinear factorization :

Semi-inclusive processes

- ▶ We can study the phenomenology of the dynamical mass in (semi-) inclusive hard processes
- ▶ interesting but challenging: **chiral-odd** sector at least at **twist-3**
- ▶ working in collinear factorization :

$$\ell N^\uparrow \rightarrow \ell j X: h_1(x) \otimes m_q^{corr}$$




dynamical mass coupled to the transversity PDF

A. Accardi, A. Bacchetta - 1706.02000 - PLB

Semi-inclusive processes

- ▶ We can study the phenomenology of the dynamical mass in (semi-) inclusive hard processes
- ▶ interesting but challenging: **chiral-odd** sector at least at **twist-3**
- ▶ working in collinear factorization :

$$e^+e^- \rightarrow h^\uparrow j X: H_1(z) \otimes m_q^{corr}$$




(Accardi, Signori et al. - in progress)

$$\begin{aligned} \frac{d\sigma^L(e^+e^- \rightarrow h^\uparrow X)}{d\Omega dz} = & \frac{3\alpha^2}{Q^2} \lambda_e \sum_a e_a^2 \left\{ \frac{C(y)}{2} \lambda_h G_{1L}(z) \right. \\ & \left. + D(y) |\mathbf{S}_T| \cos(\phi_S) \frac{2M_h}{Q} \left(\frac{G_T(z)}{z} + \frac{m_q^{corr}}{M_h} H_1(z) \right) \right\} \end{aligned}$$

Semi-inclusive processes

- ▶ We can study the phenomenology of the dynamical mass in (semi-) inclusive hard processes
- ▶ interesting but challenging: **chiral-odd** sector at least at **twist-3**
- ▶ working in collinear factorization :

$$e^+e^- \rightarrow \{h_1 h_2\} X: H_1^{\triangleleft} \otimes m_q^{corr}$$



Also requires lepton polarization

Accardi, Signori et al. - work in progress

Conclusions

- ▶ hadronization \longrightarrow a **fundamental** aspect of QCD

Conclusions

- ▶ hadronization \longrightarrow a **fundamental** aspect of QCD
- ▶ the **non-perturbative structure of inclusive jets** can be related to the properties of the quark propagator, in particular the normalization of the **spectral functions**

Conclusions

- ▶ hadronization \longrightarrow a **fundamental** aspect of QCD
- ▶ the **non-perturbative structure of inclusive jets** can be related to the properties of the quark propagator, in particular the normalization of the **spectral functions**
- ▶ at **twist two** these non-perturbative effects are **trivial** ($\int d\mu^2 \rho_3(\mu^2) \equiv 1$)

Conclusions

- ▶ hadronization \longrightarrow a **fundamental** aspect of QCD
- ▶ the **non-perturbative structure of inclusive jets** can be related to the properties of the quark propagator, in particular the normalization of the **spectral functions**
- ▶ at **twist two** these non-perturbative effects are **trivial** ($\int d\mu^2 \rho_3(\mu^2) \equiv 1$)
- ▶ at **twist three** the non-perturbative structure emerges as a **mass term** with a current and a **dynamical** component (explicit and dynamical CSB)

Conclusions

- ▶ hadronization \longrightarrow a **fundamental** aspect of QCD
- ▶ the **non-perturbative structure of inclusive jets** can be related to the properties of the quark propagator, in particular the normalization of the **spectral functions**
- ▶ at **twist two** these non-perturbative effects are **trivial** ($\int d\mu^2 \rho_3(\mu^2) \equiv 1$)
- ▶ at **twist three** the non-perturbative structure emerges as a **mass term** with a current and a **dynamical** component (explicit and dynamical CSB)
- ▶ this mass is gauge-invariant, and the dynamical component **can be accessed** at twist three in scattering experiments and via momentum sum rules for twist three FFs

Backup



Useful references/1:

A selection of useful references related to inclusive jets and dynamical mass effects:

- ▶ Fully unintegrated parton correlation functions and factorization in lowest order hard scattering
J.C. Collins, T.C. Rogers, A.M. Stasto - 0708.2833
- ▶ Collinear factorization for deep inelastic scattering structure functions at large Bjorken x_B
A. Accardi, J.W. Qiu - 0805.1496
- ▶ Quark fragmentation as a probe of dynamical mass generation
A. Accardi, A. Signori - 1903.04458
- ▶ On the connection between quark propagation and hadronization
A. Accardi, A. Signori - 2005.11310
- ▶ Accessing the nucleon transverse structure in deep-inelastic scattering
A. Accardi, A. Bacchetta - 1706.02000

Useful references/2:

A selection of useful references dealing with fragmentation functions, inclusive jets in pQCD, e^+e^- annihilation:

- ▶ Parton fragmentation functions (review)

A. Metz, A. Vossen - 1607.02521

- ▶ Quark fragmentation within an identified jet

M. Procura, I. Stewart - 0911.4980

- ▶ Parton fragmentation within an identified jet at NNLL

A. Jain, M. Procura, W. Waalewijn - 1101.4953

- ▶ Asymmetries in polarized hadron production in e^+e^- annihilation up to order $1/Q$


D. Boer, R. Jakob, P.J. Mulders - hep-ph/9702281

- ▶ Angular dependences in inclusive two-hadron production at Belle

D. Boer - 0804.2408

The cut quark propagator


$$\Xi_{ij}(k; v) = \text{Disc} \int \frac{d^4\xi}{(2\pi)^4} e^{ikx} \frac{\text{Tr}_c}{N_c} \langle \Omega | \hat{T} W_1(\infty, \xi; v) \psi_i(\xi) \bar{\psi}_j(0) W_2(0, \infty; v) | \Omega \rangle$$



- ▶ **Partonic picture:** gauge invariant dressed quark correlator
 - ▶ only the discontinuity is considered \rightarrow on-shellness
 - ▶ the color is neutralized

The cut quark propagator


$$\Xi_{ij}(k; v) = \text{Disc} \int \frac{d^4\xi}{(2\pi)^4} e^{ikx} \frac{\text{Tr}_c}{N_c} \langle \Omega | \hat{T} W_1(\infty, \xi; v) \psi_i(\xi) \bar{\psi}_j(0) W_2(0, \infty; v) | \Omega \rangle$$



- ▶ **Partonic picture:** gauge invariant dressed quark correlator
 - ▶ only the discontinuity is considered \rightarrow on-shellness
 - ▶ the color is neutralized
- ▶ **Hadronic picture:** “fully inclusive jet” correlator
 - ▶ X : the complete set of hadronization products crossing the cut
 - ▶ no hadrons are measured
 - ▶ the scale is defined by the end-point kinematics

The cut quark propagator

$$\Xi_{ij}(k; v) = \text{Disc} \int \frac{d^4\xi}{(2\pi)^4} e^{ikx} \frac{\text{Tr}_c}{N_c} \langle \Omega | \hat{T} W_1(\infty, \xi; v) \psi_i(\xi) \bar{\psi}_j(0) W_2(0, \infty; v) | \Omega \rangle$$



- ▶ **Partonic picture:** gauge invariant dressed quark correlator
 - ▶ only the discontinuity is considered \rightarrow on-shellness
 - ▶ the color is neutralized
- ▶ **Hadronic picture:** “fully inclusive jet” correlator
 - ▶ X : the complete set of hadronization products crossing the cut
 - ▶ no hadrons are measured
 - ▶ the scale is defined by the end-point kinematics
- ▶ insights into **dynamical generation** of mass and momentum and **chiral symmetry** breaking

The quark/jet mass

$$M_j(k^-) \sim \int dk^+ \text{Tr}_D [\Xi \mathbb{I}]$$




Mass associated with the scalar term (**chiral-odd**) of the cut quark propagator:

- inclusive “**jet mass**” or color-screened dressed **quark mass**

The quark/jet mass

$$M_j(k^-) \sim \int dk^+ \text{Tr}_D [\Xi \mathbb{I}]$$



Mass associated with the scalar term (**chiral-odd**) of the cut quark propagator:

- inclusive “**jet mass**” or color-screened dressed **quark mass**

In the light-cone gauge we can relate it to the chiral-odd spectral function for the quark propagator:

$$M_j = \int_0^{+\infty} d\mu^2 \sqrt{\mu^2} \rho_1^{l_{cg}}(\mu^2)$$



The quark/jet mass

$$M_j(k^-) \sim \int dk^+ \text{Tr}_D [\Xi \mathbb{I}]$$



Mass associated with the scalar term (**chiral-odd**) of the cut quark propagator:

- ▶ inclusive “**jet mass**” or color-screened dressed **quark mass**

In the light-cone gauge we can relate it to the chiral-odd spectral function for the quark propagator:

$$M_j = \int_0^{+\infty} d\mu^2 \sqrt{\mu^2} \rho_1^{l_{cg}}(\mu^2)$$

This mass term:

- ▶ gauge-invariant
- ▶ renormalization scale dependent
- ▶ calculable via the spectral functions of the cut quark propagator
- ▶ **accessible via momentum sum rules for twist-3 FFs**

Semi-inclusive processes

- ▶ We can study the phenomenology of the dynamical mass in (semi-) inclusive hard processes
- ▶ interesting but challenging: **chiral-odd** sector at least at **twist-3**
- ▶ working in collinear factorization :

$$\text{▶ } (?) \quad pp^\uparrow \rightarrow h_1 h_2 j X \quad \xrightarrow[\text{sum rule}]{\text{mass}} \quad f_1(x_1) \otimes h_1(x_2) \otimes D_1(z) \otimes m_q^{\text{corr}}$$

(fixed-target configuration at LHC)

- ▶ (?) potentially also TMD factorization
- ▶ in order to make quantitative predictions and extractions the factorization of these processes has to be addressed