


Detector Maps from Inclusive Group

Barak, Nobuo and Renee

May 6, 2020


Critical Channels for Detector Development

- The inclusive group covers a large set of physics interests, ranging from gluon helicity to nuclear PDFs and non-linear dynamics.
- Parity violating (A_{PV}) and charged current (CC) channels were singled out, both for high scientific interest and stringent detector requirements.
 - I. Inclusive A_{PV} - Sensitive to $\Delta s^+(Q^2)$ and $s^+(Q^2)$. Small signal requires high precision measurement. Electron PID will be critical.
 - II. σ_r^{CC} - Allows flavor separation at high x for sbar. Reduced cross-section requires reconstruction of Q^2 , x from hadronic recoil. Pushing down thresholds for ECAL/HCAL will be critical.




Inclusive e^- maps

- Work of Barak Schmookler
- θ_e defined WRT +z
- Yields represent 1 fb^{-1}



Inclusive e^- maps

- Work of Barak Schmookler
- θ_e defined WRT +z
- Yields represent 1 fb^{-1}




Hadronic Recoil Maps

- This is the work of Xiaoxuan Chu. The full details are presented here <https://indico.bnl.gov/event/8389/>
- Use Django, e- beam = 18 GeV, p beam = 275 GeV e-p, $\sqrt{s} = 141$ GeV. Radiative corrections turned on. $L = 10 \text{ fb}^{-1}$, $0.01 < y < 0.95$ and $100 < Q^2 < 10^5 \text{ GeV}^2$.
- Reduced cross-section is extracted using Jacquet-Blondel kinematic reconstruction


$$x_{JB} = \frac{Q_{JB}^2}{sy_{JB}}; \quad y_{JB} = \frac{(E - p_z)_h}{2E_e}; \quad Q_{JB}^2 = \frac{p_{t,h}^2}{1 - y_{JB}}$$

- Perfect detector is implemented

$h^{+/-}$ + neutron p_T vs η



$h^{+/-}$ + neutron E vs η



----- Charged Current Charged Hadron + neutron -----

eta = -3.960 theta = 0.038 pT = 0.215+/-0.000 E = 0.762+/-0.291

eta = -3.880 theta = 0.041 pT = 0.215+/-0.000 E = 0.963+/-0.348

...

eta = -2.040 theta = 0.259 pT = 0.546+/-0.017 E = 2.021+/-0.069

eta = -1.960 theta = 0.280 pT = 0.590+/-0.018 E = 2.113+/-0.065

...

eta = -1.000 theta = 0.705 pT = 1.259+/-0.015 E = 1.971+/-0.023

eta = -0.920 theta = 0.758 pT = 1.350+/-0.015 E = 1.993+/-0.022

...

eta = -0.040 theta = 1.531 pT = 2.257+/-0.016 E = 2.285+/-0.016

eta = 0.040 theta = 1.611 pT = 2.350+/-0.016 E = 2.379+/-0.016

...

eta = 1.000 theta = 2.437 pT = 2.688+/-0.018 E = 4.194+/-0.028

eta = 1.080 theta = 2.487 pT = 2.683+/-0.019 E = 4.462+/-0.031

...

eta = 2.040 theta = 2.883 pT = 1.190+/-0.013 E = 4.823+/-0.050

eta = 2.120 theta = 2.903 pT = 1.081+/-0.012 E = 4.730+/-0.051

...



eta = 3.000 theta = 3.042 pT = 0.407+/-0.004 E = 4.523+/-0.040

eta = 3.080 theta = 3.050 pT = 0.373+/-0.003 E = 4.525+/-0.034

....

eta = 3.880 theta = 3.100 pT = 0.301+/-0.002 E = 8.314+/-0.035

eta = 3.960 theta = 3.103 pT = 0.304+/-0.002 E = 9.035+/-0.038

Gamma p_T vs η Gamma E vs η 

----- Charged Current Photon -----

eta = -4.000 theta = 0.037 $p_T = 0.254 \pm 0.016$ E = 1.578 ± 0.098

eta = -3.840 theta = 0.043 $p_T = 0.263 \pm 0.016$ E = 1.598 ± 0.098

...

eta = -2.080 theta = 0.249 $p_T = 0.525 \pm 0.008$ E = 0.939 ± 0.035

eta = -1.920 theta = 0.291 $p_T = 0.551 \pm 0.007$ E = 0.864 ± 0.029

...

eta = -0.000 theta = 1.571 $p_T = 1.176 \pm 0.007$ E = 0.856 ± 0.007

eta = 0.160 theta = 1.730 $p_T = 1.247 \pm 0.007$ E = 0.948 ± 0.008

...

eta = 0.960 theta = 2.410 $p_T = 1.342 \pm 0.008$ E = 1.614 ± 0.012

eta = 1.120 theta = 2.511 $p_T = 1.367 \pm 0.008$ E = 1.879 ± 0.014

...

eta = 1.920 theta = 2.850 $p_T = 0.884 \pm 0.006$ E = 2.075 ± 0.021

eta = 2.080 theta = 2.893 $p_T = 0.797 \pm 0.005$ E = 2.045 ± 0.023

...

eta = 3.040 theta = 3.046 $p_T = 0.526 \pm 0.001$ E = 2.086 ± 0.017

eta = 3.200 theta = 3.060 $p_T = 0.516 \pm 0.001$ E = 2.280 ± 0.016

...

eta = 4.000 theta = 3.105 $p_T = 0.507 \pm 0.000$ E = 4.660 ± 0.021

eta = 4.160 theta = 3.110 $p_T = 0.508 \pm 0.000$ E = 5.462 ± 0.024

Updated Timeline for YR deliverables

- Physics Object Maps (Experimentalists)
 1. Barak - inclusive electron channels
 2. Xiaoxuan – hadronic recoil channels
 3. Not currently pursuing heavy flavor tagged channels
- Detector Constraints (Experimentalists)
 1. Electron PID as a function of η in e+p - Hanjie
 2. Determine minimum thresholds and resolutions for ECAL/HCAL – Xiaoxuan
- Cross-section/Asymmetry correction and reconstruction (Experimentalists)
 1. Hanjie - σ^{NC}
 2. Matt – g_2 and higher twist
 3. Xiaoxuan - σ^{CC} and σ^{NC}
- Impact Studies (Theorists)
 1. Statistical indicators T-test, KS test for preliminary impact (fast global fit replacement)
 2. Global Fits, including possible reweighting if necessary

Special Requests for Pavia Meeting

- Meeting between inclusive group conveners and physics+detector conveners.
- Meeting with SIDIS to discuss common detector and radiative correction tools.
- Meeting with Heavy Flavor to discuss tagged cross-sections.