## XXVIII International Workshop on Deep-Inelastic Scattering and Related Subjects



Contribution ID: 555

Type: Contributed Talk

## Light meson nuclear modification factor in p-Pb collisions over an unprecedented **XT** range with ALICE

Wednesday, 14 April 2021 08:36 (18 minutes)

Light meson measurements in high-energy proton-nucleus collisions provide a probe to study the physics of strongly interacting matter and the quark-gluon plasma. In particular, measuring the nuclear modification factor, where the particle production in p-A is compared to a baseline proton-proton reference which is scaled to account for nuclear geometry, provides important insight to the modification of nucleon structure functions in nuclei and a baseline for the observed strong suppression of hadron yields at high  $p_{\rm T}$  in heavy-ion collisions.

In this talk, neutral pion and  $\eta$  meson invariant differential yields and nuclear modification factors at midrapidity are presented with a  $p_{\rm T}$  reach beyond 100 GeV/*c* and up to 40 GeV/*c*, respectively, and compared to theoretical model calculations. Both neutral mesons are measured via their diphoton decay channel in pp and p-Pb collisions at  $\sqrt{s_{\rm NN}} = 5.02$  and 8.16 TeV with the ALICE experiment at the CERN LHC. The analysis combines results from several partially independent reconstruction techniques where the decay photons were detected with the electromagnetic calorimeter, EMCal, the photon spectrometer, PHOS, or via reconstruction of  $e^+e^-$  pairs from conversions in the ALICE detector material using the central tracking system.

Primary author: KOENIG, Joshua (IKF, Universität Frankfurt)Presenter: KOENIG, Joshua (IKF, Universität Frankfurt)Session Classification: QCD with Heavy Flavors and Hadronic Final States

Track Classification: QCD with Heavy Flavors and Hadronic Final States