

XXVIII International Workshop on Deep-Inelastic Scattering and Related Subjects



Contribution ID: 462

Type: **Contributed Talk**

Measurement of differential cross sections of weak boson production in unpolarized $p+p$ collisions at STAR

We present the status for the measurement of cross sections of weak boson production as a function of the boson's kinematics, by the STAR experiment at RHIC. Results combine data from unpolarized $p+p$ collisions at $\sqrt{s} = 500$ and 510 GeV collected during the 2011, 2012, 2013, and 2017 runs, corresponding to an integrated luminosity of 700 pb^{-1} . The differential Z^0 cross section, measured as a function of the boson's p_T , provides important constraints on the energy scale dependence of transverse momentum distributions of partons inside the proton. The W^+/W^- cross-section ratio as a function of the boson's rapidity is sensitive to the non-perturbative \bar{d}/\bar{u} distribution. The momentum fraction range ($0.1 < x < 0.3$) covered by these measurements naturally complements the phase space accessed at the LHC, providing critical inputs to global fits.

Primary author: FAZIO, Salvatore (Brookhaven National Laboratory)

Presenter: FAZIO, Salvatore (Brookhaven National Laboratory)

Session Classification: Structure function and parton densities

Track Classification: Structure Functions and Parton Densities