

Systematic Extraction of Pion Parton Distributions Using Threshold Resummation and Applications to Pion TMDs

Patrick Barry, Nobuo Sato, Wally Melnitchouk, Chueng-Ryong Ji, Leonard
Gamberg, and Astrid Hiller Blin

DIS 2021: Structure function and parton densities 4/15/2021


Motivation

What to do:

- **Define** a structure of hadrons in terms of quantum field theories
- **Identify** theoretical observables that factorize into non-perturbative objects and perturbatively calculable physics
- Perform **global QCD analysis** as structures are universal and are the same in all subprocesses


Pions

- Pion is the **Goldstone boson** associated with spontaneous symmetry breaking of chiral $SU(2)_L \times SU(2)_R$ symmetry
- **Lightest hadron** as $\frac{m_\pi}{M_N} \ll 1$ and dictates the nature of hadronic interactions at low energies
- Simultaneously a pseudoscalar meson made up of q and $\bar{q}$ constituents




Experiments to Probe Pion Structure

- Drell-Yan (DY)



- Accelerating pion allows for time dilation and longer lifetime

- Leading Neutron (LN)



- Barely striking surface of a target proton knocks out an almost on-shell pion to probe

Datasets -- Kinematics


- Large x_π -- Drell-Yan (DY)
- Small x_π -- Leading Neutron (LN)
- Not much data overlap

- In DY:

$$x_\pi = \frac{1}{2} \left(x_F + \sqrt{x_F^2 + 4\tau} \right)$$


- In LN:

$$x_\pi = x_B / \bar{x}_L$$




JAM18 Pion PDFs

- Lightly shaded bands – only Drell-Yan data
- Darkly shaded bands – fit to both Drell-Yan and LN data



PCB, N. Sato, W. Melnitchouk and Chueng-Ryong Ji,
Phys. Rev. Lett. **121**, 152001 (2018).

Soft Gluon Resummation



- Fixed-target Drell-Yan notoriously has large- x_F contamination of higher orders
- Large logarithms may spoil perturbation
- Focus on corrections to the most important $q\bar{q}$ channel
- Resum contributions to all orders of α_s

Issues with Perturbative Calculations

$$\hat{\sigma} \sim \delta(1-z) + \alpha_S (\log(1-z))_+ \longrightarrow \hat{\sigma} \sim \delta(1-z) [1 + \alpha_S \log(1-\tau)]$$

- If τ is large, can potentially spoil the perturbative calculation
- Improvements can be made by resumming $\log(1-z)_+$ terms

Next-to-Leading + Next-to-Leading Logarithm Order Calculation

An NLO calculation
gathers the $\mathcal{O}(\alpha_s)$
terms

	<u>LL</u>	<u>NLL</u>	<u>...</u>	<u>N^pLL</u>
LO	1	--	...	--
NLO	$\alpha_s \log(N)^2$	$\alpha_s \log(N)$	...	--
NNLO	$\alpha_s^2 \log(N)^4$	$\alpha_s^2 (\log(N)^2, \log(N)^3)$	...	--
...	...	...	...	...
N ^k LO	$\alpha_s^k \log(N)^{2k}$	$\alpha_s^k (\log(N)^{2k-1}, \log(N)^{2k-2})$	...	$\alpha_s^k \log(N)^{2k-2p} + \dots$

Next-to-Leading + Next-to-Leading Logarithm Order Calculation

Add the columns to
the rows

	<u>LL</u>	<u>NLL</u>	<u>...</u>	<u>N^pLL</u>
LO	1	--	...	--
NLO	$\alpha_s \log(N)^2$	$\alpha_s \log(N)$	...	--
NNLO	$\alpha_s^2 \log(N)^4$	$\alpha_s^2 (\log(N)^2, \log(N)^3)$	...	--
...	...	...	...	...
N ^k LO	$\alpha_s^k \log(N)^{2k}$	$\alpha_s^k (\log(N)^{2k-1}, \log(N)^{2k-2})$	...	$\alpha_s^k \log(N)^{2k-2p} + \dots$

Next-to-Leading + Next-to-Leading Logarithm Order Calculation

Make sure only counted once!
- Subtract the matching

	<u>LL</u>	<u>NLL</u>	<u>...</u>	<u>N^pLL</u>
LO	1	--	...	--
NLO	$\alpha_s \log(N)^2$	$\alpha_s \log(N)$	...	--
NNLO	$\alpha_s^2 \log(N)^4$	$\alpha_s^2 (\log(N)^2, \log(N)^3)$	...	--
...	...	...	...	...
N ^k LO	$\alpha_s^k \log(N)^{2k}$	$\alpha_s^k (\log(N)^{2k-1}, \log(N)^{2k-2})$	...	$\alpha_s^k \log(N)^{2k-2p} + \dots$

Origin of Landau Pole

$$\alpha_S C_{\text{soft}}^{(1)}(N) = 2 \frac{C_F}{\pi} \int_0^1 dz \frac{z^{N-1} - 1}{1-z} \int_{Q^2}^{(1-z)^2 Q^2} \frac{dk_{\perp}^2}{k_{\perp}^2} \alpha_S(k_{\perp}^2)$$


- Upper Limits imply that $k_{\perp}^2$ will go to 0
- $\alpha_S(\mu^2 = 0)$ is **NOT** well-defined
- **Ambiguities** on how to deal with this provide needs for prescriptions

Methods of Resummation

- Make use of the **Minimal Prescription** to avoid Landau Pole
- Rapidity distribution $\frac{d\sigma}{dQ^2 dY}$ adds more complications
- We can perform a **Mellin-Fourier transform** to account for the rapidity
 - A cosine appears while doing Fourier transform; options:
 - 1) Take first order **expansion**, cosine ≈ 1
 - 2) Keep **cosine** intact
- Can additionally perform a **Double Mellin transform**
- **Explore** the different methods and **analyze** effects


Data and Theory Comparison – Drell-Yan

- Cosine method tends to overpredict the data at very large x_F
- Double Mellin method is qualitatively very similar to NLO
- Resummation is largely a high- x_F effect




PDF Results

- Large x behavior in valence depends on prescription



Effective β_v parameter


- $q_v(x) \sim (1 - x)^{\beta_v}$ as $x \rightarrow 1$
- Threshold resummation does not give universal behavior of β_v
- NLO and double Mellin give $\beta_v \approx 1$
- Cosine and Expansion give $\beta_v > 2$



Transverse Momentum- dependent Drell-Yan

p_T -dependent spectrum for pion data

- Small- p_T data – TMD factorization – partonic transverse momentum
- Large- p_T data – collinear factorization – recoil transverse momentum




E615 πW Drell-Yan

Phys. Rev. D **39**, 92 (1989).

p_T -dependent spectrum for pion data


- Various factorization theorems break down in certain regions of p_T
- Errors are related with $\mathcal{O}(p_T/Q)$ (low- p_T) or $\mathcal{O}(m/p_T)$ (large- p_T)




E615 πW Drell-Yan

Phys. Rev. D **39**, 92 (1989).

JAM20 Pion PDFs



N.Cao, PCB, N. Sato, and W. Melnitchouk
arXiv:2103.02159 [hep-ph]



- For the first time, we included **large p_T** -dependent Drell-Yan data, which follows collinear factorization
- Large p_T does **not** dramatically affect the PDF
- Successfully describe data with a scale $\mu = p_T/2$

TMD factorization

- In small- p_T region, Use the CSS formalism for TMD evolution


$$\begin{aligned}
 \frac{d\sigma}{dQ^2 dy dq_T^2} &= \frac{4\pi^2 \alpha^2}{9Q^2 s} \sum_{j, j_A, j_B} H_{j\bar{j}}^{\text{DY}}(Q, \mu_Q, a_s(\mu_Q)) \int \frac{d^2 \mathbf{b}_T}{(2\pi)^2} e^{i\mathbf{q}_T \cdot \mathbf{b}_T} \\
 &\times e^{-g_{j/A}(x_A, b_T; b_{\max})} \int_{x_A}^1 \frac{d\xi_A}{\xi_A} f_{j_A/A}(\xi_A; \mu_{b_*}) \tilde{C}_{j/j_A}^{\text{PDF}}\left(\frac{x_A}{\xi_A}, b_*; \mu_{b_*}^2, \mu_{b_*}, a_s(\mu_{b_*})\right) \\
 &\times e^{-g_{\bar{j}/B}(x_B, b_T; b_{\max})} \int_{x_B}^1 \frac{d\xi_B}{\xi_B} f_{j_B/B}(\xi_B; \mu_{b_*}) \tilde{C}_{\bar{j}/j_B}^{\text{PDF}}\left(\frac{x_B}{\xi_B}, b_*; \mu_{b_*}^2, \mu_{b_*}, a_s(\mu_{b_*})\right) \\
 &\times \exp \left\{ -g_K(b_T; b_{\max}) \ln \frac{Q^2}{Q_0^2} + \tilde{K}(b_*; \mu_{b_*}) \ln \frac{Q^2}{\mu_{b_*}^2} + \int_{\mu_{b_*}}^{\mu_Q} \frac{d\mu'}{\mu'} \left[2\gamma_j(a_s(\mu')) - \ln \frac{Q^2}{(\mu')^2} \gamma_K(a_s(\mu')) \right] \right\}
 \end{aligned}$$

Non-
perturbative
TMDs to extract

- Approach with **leading order** to diagnose current situation


Results – E288 pp data

- Perform simultaneous fit of **small- p_T** TMDs to E288 (pp) and E615 (πA) data
- FO is prediction using collinear PDFs and scale $\mu = p_T/2$



Results – E615 πW data

- Proton and pion TMDs are different, but g_K are same
- Perform simultaneous fit of **small- p_T** TMDs to E288 (pp) and E615 (πA) data




Future Work

- Investigate high- x behavior of valence PDF through constraints from the **lattice data**
- Perform a **simultaneous extraction** of pion PDFs and TMDs using available low- and high- p_T data
- Explore matching procedure between W and FO terms in p_T spectrum

Backup

Previous Pion PDFs

- Fits to Drell-Yan, prompt photon, or both




GRS, GRV, and SMRS

Z. Phys. C **67**, 433 (1995).


Eur. Phys. J. C **10** 313 (1997).

Phys. Rev. D **45** 2349 (1992).



Aicher's valence PDF


Phys. Rev. Lett. **105**, 114023 (2011).



xFitter

Phys. Rev. D **102**, 014040 (2020).

Drell-Yan (DY)



$$\sigma \propto \sum_{i,j} f_i^\pi(x_\pi, \mu) \otimes f_j^A(x_A, \mu) \otimes \hat{\sigma}_{i,j}(x_\pi, x_A, Q/\mu)$$

Drell-Yan (DY) Definitions

Hadronic variable

$$\tau = \frac{Q^2}{S}$$


Partonic variable

$\hat{S}$ is the center of mass
momentum squared of
incoming partons

$$z \equiv \frac{Q^2}{\hat{S}} = \frac{\tau}{x_1 x_2}$$

LO

LO: $\mathcal{O}(1)$




$$C_{q\bar{q}} = \delta(1 - z) \frac{\delta(y) + \delta(1 - y)}{2}$$

- $z = 1$ corresponds to partonic threshold
- All $\hat{S}$ is equal to Q^2
- All energy of hard partons turns into virtuality of photon

NLO Virtual

- Virtual corrections at NLO are proportional to $\delta(1 - z)$
 - Exhibit **Born kinematics**




$$C_{q\bar{q}}^{\text{virtual}} = \delta(1 - z) \frac{\delta(y) + \delta(1 - y)}{2} \left[\frac{C_F \alpha_S}{\pi} \left(\frac{3}{2} \ln \frac{Q^2}{\mu^2} + \frac{2\pi^2}{3} - 6 \right) \right]$$


NLO Real Emission

- Next to leading order, real gluon emissions

$$C_{q\bar{q}}^{\text{real}} = \frac{C_F \alpha_S}{\pi} \left[\frac{\delta(y) + \delta(1-y)}{2} \left[(1+z^2) \left(\frac{1}{1-z} \ln \frac{Q^2(1-z)^2}{\mu^2 z} \right)_+ + 1-z \right] \right. \\ \left. + \frac{1}{2} \left[\frac{(1-z)^2}{z} y(1-y) \right] \left[\frac{1+z^2}{1-z} \left(\left[\frac{1}{y} \right]_+ + \left[\frac{1}{1-y} \right]_+ \right) - 2(1-z) \right] \right]$$




NLO Real Emission



- Real quark emissions
- $C_{qg} = C_{gq}|_{y \rightarrow 1-y}$

$$C_{qg}^{\text{real}} = \frac{T_F \alpha_S}{2\pi} \left[\delta(y) \left[(z^2 + (1-z)^2) \ln \frac{Q^2(1-z)^2}{\mu^2 z} + 2z(1-z) \right] + \left[1 + \frac{(1-z)^2}{z} y(1-y) \right] \left[(z^2 + (1-z)^2) \left(\frac{1}{y} \right)_+ + 2z(1-z) + (1-z)^2 y \right] \right]$$



Leading Neutron (LN)



$$\frac{d\sigma}{dx dQ^2 dy} \sim f_{p \rightarrow \pi^+ n}(y) \times \sum_q \int_{x/y}^1 \frac{d\xi}{\xi} C(\xi) q\left(\frac{x/y}{\xi}, \mu^2\right)$$

Large x_L

- x_L is fraction of longitudinal momentum carried by neutron relative to initial proton
- For t to be close to pion pole, has to go near 0 – happens at large x_L
- In this region, one pion exchange dominates



Splitting Function and Regulators

Amplitude for proton to dissociate into a π^+ and neutron:

$$f_{\pi N}(\bar{x}_L) = \frac{g_A^2 M^2}{(4\pi f_\pi)^2} \int dk_\perp^2 \frac{\bar{x}_L [k_\perp^2 + \bar{x}_L^2 M^2]}{x_L^2 D_{\pi N}^2} |\mathcal{F}|^2,$$

$$D_{\pi N} \equiv t - m_\pi^2 = -\frac{1}{1-y} [k_\perp^2 + y^2 M^2 + (1-y)m_\pi^2]$$

$$\mathcal{F} = \begin{cases} \text{(i)} & \exp((M^2 - s)/\Lambda^2) \\ \text{(ii)} & \exp(D_{\pi N}/\Lambda^2) \\ \text{(iii)} & (\Lambda^2 - m_\pi^2)/(\Lambda^2 - t) \\ \text{(iv)} & \bar{x}_L^{-\alpha_\pi(t)} \exp(D_{\pi N}/\Lambda^2) \\ \text{(v)} & [1 - D_{\pi N}^2/(\Lambda^2 - t)^2]^{1/2} \end{cases}$$

s -dep. exponential



Best fit

t -dep. exponential

t -dep. monopole

Regge

Pauli-Villars

- We examine five regulators, and we fit Λ
- $\mathcal{F}$ is a UV regulator, which the data chooses

Bayesian Inference

- Minimize the χ^2 for each replica

$$\chi^2(\mathbf{a}, \text{data}) = \sum_e \left(\sum_i \left[\frac{d_i^e - \sum_k r_k^e \beta_{k,i}^e - t_i^e(\mathbf{a})}{\alpha_i^e} \right]^2 + \left(\frac{1 - n_e}{\delta n_e} \right)^2 + \sum_k (r_k^e)^2 \right)$$

Normalization parameter

- Perform N total χ^2 minimizations and compute statistical quantities

Expectation value

$$E[\mathcal{O}] = \frac{1}{N} \sum_k \mathcal{O}(\mathbf{a}_k),$$

Variance

$$V[\mathcal{O}] = \frac{1}{N} \sum_k [\mathcal{O}(\mathbf{a}_k) - E[\mathcal{O}]]^2,$$

TMD Non-Perturbative Parametrization


- Parametrize g_j for each hadron and g_K as in Slide 21

$$g_{j/h}(x, b_T^2) = -b_T^2 g_1 \left(\frac{1}{2} + g_3 \log(10x) \right)$$

$$g_K(b_T) = -b_T^2 \frac{g_2}{2}$$

χ^2 as a function of $x_F^{\max}$

- Phenomenologically determine where TMD factorization breaks down in large x_F data



E288 pp

- Calculate large- p_T FO term with $\mu = Q$
- **Not** a good description

