## XXVIII International Workshop on Deep-Inelastic Scattering and Related Subjects



Contribution ID: 696

Type: Contributed Talk

## Measurements of the Antiquark Flavor Asymmetry in the Proton by the Drell-Yan Experiment SeaQuest

Thursday, 15 April 2021 09:12 (18 minutes)

Naively, the amounts of  $\bar{d}$  and  $\bar{u}$  in the proton were expected to be the same based on the flavor-independence of the strong coupling.

However, the muon deep inelastic scattering experiment NMC at CERN found  $\bar{d} > \bar{u}$  in the proton. Drell–Yan experiments also obtained the results consistent with it.

The Drell–Yan experiment E866 at Fermilab showed that  $\bar{d}(x)/\bar{u}(x) > 1.0$  for 0.015 < x < 0.20.

It also showed  $\bar{d}(x)/\bar{u}(x) < 1.0$  at large x ( $x \sim 0.3$ ), although it is consistent with 1.0 within statistical uncertainty.

SeaQuest is a Drell–Yan experiment at Fermilab, that measured the antiquark flavor asymmetry  $\bar{d}/\bar{u}$  precisely in for a wide x range (0.13 < x < 0.45) including the intriguing region from E866.

It uses a 120 GeV proton beam extracted from Fermilab Main Injector colliding with liquid hydrogen and deuterium targets.

The antiquark flavor asymmetry  $\bar{d}(x)/\bar{u}(x)$  is derived from the cross section ratio of proton-deuterium to proton-proton Drell–Yan processes.

In this talk, the SeaQuest results of the  $\bar{d}(x)/\bar{u}(x)$  analysis will be presented.

Primary author: NAGAI, Kei (Los Alamos National Laboratory)

Presenter: NAGAI, Kei (Los Alamos National Laboratory)

Session Classification: Structure function and parton densities

Track Classification: Structure Functions and Parton Densities