

XXVIII International Workshop on Deep-Inelastic Scattering and Related Subjects



Contribution ID: 381

Type: **Contributed Talk**

Quark Siverson Function at Small- x : Return of a Spin-Dependent Odderon

Tuesday, 13 April 2021 12:18 (18 minutes)

We apply the formalism developed earlier [1, 2] for studying the small- x asymptotics of transverse momentum dependent parton distribution functions (TMDs) to construct the small- x asymptotics of the quark Siverson function. We explicitly construct the transversely polarized, fundamental “Wilson line” operator to sub-sub-eikonal order. We then express the Siverson function in terms of dipole scattering amplitudes containing the transversely polarized “Wilson line”, and show that the main term which contributes to the Siverson function is the spin-dependent odderon, similar to the case of the gluon Siverson function derived by Boer, Eschevarria, Mulders and Zhou [3] (see also [4]).

[1] Y. V. Kovchegov and M. D. Sievert, Phys. Rev. D99, 054032 (2019).

[2] Y. V. Kovchegov and M. D. Sievert, Phys. Rev. D99, 054033 (2019).

[3] D. Boer, M. G. Echevarria, P. Mulders, and J. Zhou, Phys. Rev. Lett. 116, 122001 (2016).

[4] L. Szymanowski and J. Zhou, Phys. Lett. B760, 249 (2016).

Primary authors: SANTIAGO, M Gabriel (Ohio State University); KOVCHEGOV, Yuri (The Ohio State University)

Presenter: SANTIAGO, M Gabriel (Ohio State University)

Session Classification: Small- x , Diffraction and Vector Mesons

Track Classification: Small- x , Diffraction and Vector Mesons