## XXVIII International Workshop on Deep-Inelastic Scattering and Related Subjects



Contribution ID: 728

Type: Contributed Talk

## **Transverse momentum dependent splitting functions in the Parton Branching method**

*Thursday, 15 April 2021 10:00 (18 minutes)* 

The Parton Branching (PB) approach provides a way to obtain transverse momentum dependent (TMD) parton densities. Its equations are written in terms of splitting functions and Sudakov form factors and can be solved with Monte Carlo methods. Even though the transverse momentum is known in every branching, the PB method currently uses the DGLAP splitting functions, which assume that the parton has no transverse momentum. We propose to extend the PB method by including TMD splitting functions, a concept from high-energy factorization.

We present the evolution equations and their solutions obtained with a Monte Carlo Simulation and show numerically the effects that TMD splitting functions have on the TMD distribution functions.

**Primary authors:** KUSINA, Aleksander (Institute of Nuclear Physics PAN); LELEK, Aleksandra (University of Antwerp); HAUTMANN, Francesco; JUNG, Hannes (DESY); KUTAK, Krzysztof (Institute of Nuclear Physics Polish Academy of Sciences); KEERSMAEKERS, Lissa (University of Antwerp); HENTSCHINSKI, Martin (Universidad de las Americas Puebla)

Presenter: KEERSMAEKERS, Lissa (University of Antwerp)

Session Classification: Structure function and parton densities

Track Classification: Structure Functions and Parton Densities