



Breakthroughs in Detector Technology (& how we can ensure they continue)

Outline:

Why? Instrumentation the Great Enabler of Science


How? The DOE HEP Detector R&D Roadmap (2020)

For latest developments see:



Ian Shipsey co-Chair, DOE HEP Detector R&D BRN

BUILDING AN UNDERSTANDING OF THE UNIVERSE: A WORK A CENTURY IN THE MAKING



.... highly predictive and rigorously tested in some cases to 1 part in 10 billion

Instrumentation the great enabler

- Discovery of the CMB
- Weak Neutral Currents
- J/Psi discovery
- Discovery of the Tau-Neutrino
- Discovery of the top quark
- Discovery of the Higgs Boson
- Discovery of the Quark Gluon Plasma
- Discovery of Neutrino Oscillations
- Accelerated Expansion of the Universe
- Gravitational Wave Detection
- Definition of SI unit for time

Instrumentation the great enabler

| • | Discovery of the CMBRadio Telescopes |
|---|---|
| | Weak Neutral Currents Bubble Chambers |
| • | J/Psi discovery Multi-wire Proportional Chambers |
| • | Discovery of the Tau-Neutrino Emulsions |
| • | Discovery of the top quark Silicon Strip Detectors |
| • | Discovery of the Higgs Boson Silicon Strip Detectors , PWO, LHC |
| • | Discovery of the Quark Gluon Plasma Time Projection Chamber |
| • | Discovery of Neutrino Oscillations Liquid Scintillator and Photodetectors |
| • | Accelerated Expansion of the Universe Charged Coupled Devices |
| • | Gravitational Wave Detection Decoupled Interferometer + Lasers |
| • | Definition of SI unit for time Atomic Clocks |
| | |

Instrumentation the great enabler


| | Discovery of the CMB | Radio Telescopes |
|---|---------------------------------------|--|
| • | Weak Neutral Currents | Bubble Chambers |
| • | J/Psi discovery | Multi-wire Proportional Chambers |
| • | Discovery of the Tau-Neutrino | Emulsions |
| • | Discovery of the top quark | Silicon Strip Detectors |
| • | Discovery of the Higgs Boson | Silicon Strip Detectors , PWO, LHC |
| • | Discovery of the Quark Gluon Plasma | Time Projection Chamber |
| • | Discovery of Neutrino Oscillations | Liquid Scintillator and Photodetectors |
| • | Accelerated Expansion of the Universe | Charged Coupled Devices |
| • | Gravitational Wave Detection | Decoupled Interferometer + Lasers |
| • | Definition of SI unit for time | Atomic Clocks |

How do we sustain this going forward?

"Transformative discovery in science is driven by innovation in technology. Our boldest undertakings in particle physics have at their foundation precision instrumentation. To reveal the profound connections underlying everything we see from the smallest scales to the largest distances in the Universe, to understand its fundamental constituents, and to reveal what is still unknown, we must invent, develop, and deploy advanced instrumentation."

DOE Basic Research Needs Study on HEP Instrumentation, September 2020

Basic Research Needs for High Energy Physics Detector Research & Development




DOE Basic Research Needs Study on HEP Instrumentation

Bonnie Fleming Ian Shipsey (co-Chairs)

Report of the Office of Science Workshop on Basic Research
Needs for HEP Detector Research and Development
December 11-14, 2019

Basic Research Needs for High Energy Physics Detector Research & Development



Report of the Office of Science Workshop on Basic Research Needs for HEP Detector Research and Development December 11-14, 2019

DOE Basic Research Needs Study on HEP Instrumentation

Bonnie Fleming Ian Shipsey (co-Chairs)

Strong overlap in detector technologies between HEP & NP for JLab12 and EIC

DOE BRN Instrumentation Panel



Opportunities for Discovery

Many mysteries to date go unanswered including:

The mystery of the Higgs boson

The mystery of Neutrinos

The mystery of Dark Matter

They mystery of Dark Energy

The mystery of quarks and charged leptons

The mystery of Matter – anti-Matter asymmetry

The mystery of the Hierarchy Problem

The mystery of the Families of Particles

The mystery of Inflation

The mystery of Gravity

Opportunities for Discovery

Many mysteries to date go unanswered including:

The mystery of the Higgs boson

The mystery of Neutrinos

The mystery of Dark Matter

They mystery of Dark Energy

The mystery of quarks and charged leptons

The mystery of Matter – anti-Matter asymmetry


The mystery of the Hierarchy Problem

The mystery of the Families of Particles

The mystery of Inflation

The mystery of Gravity

We are very much in a data driven era!





Four Grand Challenges encompass this Instrumentation revolution

- Advancing HEP detectors to new regimes of sensitivity: To make the unmeasurable
 measurable will require the development of sensors with exquisite sensitivity with the ability to
 distinguish signal from noise.... Research will be needed to develop these sensors with maximal
 coupling to the quanta to be sensed and push their sensitivities to ultimate limits.
- Using Integration to enable scalability for HEP sensors: Future HEP detectors for certain classes of experiments will require massive increases in scalability to search for and study rare phenomena ... A key enabler of scalability is integration of many functions on, and extraction of multidimensional information from, these innovative sensors.
- Building next-generation HEP detectors with novel materials & advanced techniques: Future HEP detectors will have requirements beyond what is possible with the materials and techniques which we know. This requires identifying novel materials ... that provide new properties or capabilities and adapting them & exploiting advanced techniques for design & manufacturing.
- Mastering extreme environments and data rates in HEP experiments:

 Future HEP detectors will involve extreme environments and exponential increases in data rates to explore elusive phenomena. ... To do so requires the intimate integration of intelligent computing with sensor technology.

- Survey the present state of the HEP technology landscape.
- Identify key capabilities & performance requirements.
- Identify technologies to provide or enhance such capabilities.

10 Basic Research Needs Study Charge



Department of Energy Office of Science Washington, DC 20585

10 July 2019

MEMORANDUM FOR HELMUT MARSISKE

FROM: GLEN CRAWFORD

DIRECTOR, RESEARCH AND TECT NOLOGY DIVISION

OFFICE OF HIGH ENERGY PHY ACS (HEP)

SUBJECT: Basic Research Needs Study C. r. Detector Re rarch and Development

I request that you organize and carry out a Basic Research Needs (BR. "ady to assess the present status of the HEP technology lan." and to identify stategic. "nologo" areas, aligned with the strengths of the US comm. "" future long-term resea. "Ad development (R&D) efforts should focus on in pursuit of the HL. "" end ertivers identifie . In the P5 report. For each of these areas, the study should artic late and pu. "" est of Priority Research Directions (PRDs) to push the technology well beyond "" cut." "" "" of the art, potentially leading to transformative advances with broad-t-nein applicability. "HEP and beyond. Furthermore, the study should identify a small set. "high-impact instrumentation "Grand Challenges" where promote, breakthroughs co. Id lead to game-changing experimental capabilities in pur and the HEP. S. "ex goals."

You should ... 'oo-chairs to le al "he study and wor "with them to select the core group of working group le. "ocarny" out. 1. "dy enc. mpasses responses to the specific charge elements elucidated. "w. "ald is expecte. "No everal months to complete. A focal point of the study should includ. "vorkshop, with au-endance beyond the core group, expected to be held in " " 2019 time fi. in the Washington, DC area. The study participants are to serve by invitation o.

The HEP Detecto. 'D program ams to develop cutting-edge, novel instrumentation to enable scientific leadership areas with ever increase a project of the property of the program of the project of the p

With u. *ar.ter a technical challenges of current high-priority P5 projects subsiding, the HEP Detector . "program aims to shift more emphasis towards building a long-term, high-risk high-rewar" ("Blue Sky") R&D portfolio that holds the promise of transformative advances with broad-ranging applications across HEP as well as other fields of science, medicine, and national security. Crucially, the program must take full advantage of the major advances happening in

- Survey the present state of the HEP technology landscape.
- Identify key capabilities & performance requirements.
- Identify technologies to provide or enhance such capabilities.
- Articulate Priority Research Directions (PRDs) to push well beyond the current state of the art, potentially leading to transformative technological advances with broad-ranging applicability.
- Flesh out required R&D efforts with deliverables with notional timelines & key technical milestones. Elucidate the technical infrastructure required to support these efforts.

10 Basic Research Needs Study Charge



SUBJECT:

Department of Energy Office of Science Washington, DC 20585

10 July 2019

MEMORANDUM FOR HELMUT MARSISKE

FROM GLEN CRAWFORD

DIRECTOR, RESEARCH AND TECT NOLOGY DIVISION OFFICE OF HIGH ENERGY PHY CS (HEP)

Basic Research Needs Study C. r. Detector Regarch and Development

I request that you organize and carry out a Basic Research Needs (BR) hidy to assess the present status of the HEP technology land me, and to identify strategic and mology areas, aligned with the strengths of the US comm in future long-term resear and development (R&D) efforts should focus on in pursuit of he Hr., "ee drivers identifie in the P5 report. For each of these areas, the study should artic late and Ju. eet of Priority Research Directions (PRDs) to push the technology well beyond the current of the art, potentially leading to transformative advances with broad-r. ngir, applicability a HEP and beyond. Furthermore, the study should identify a small set a high-impact instrumentation "Grand Challenges" where to amoro, breakthroughs could lead to game-changing experimental capabilities in purant of HEP sance goals.

You should ... co-chairs to lead the study and wor with them to select the core group of working group le. to carry it out. 12 andy encompasses responses to the specific charge elements elucidated . 'w "aid is expecte. "ake several months to complete. A focal point of the study should includ. vorkshop, with amendance beyond the core group, expected to be held in 2019 time ft. in the Washington, DC area. The study participants are to serve by invitation on

The HEP Detector 'D program ams to develop cutting-edge, novel instrumentation to enable scientific leadership a worldwide experimental program that is broadening into new research areas with ever increang demands in sensitivity, scale, and cost. To meet this challenge, HEP 'ms to execute a proj am appropriately balanced between incremental, near-term, low-risk tor R&D and transformative, long-term, high-risk detector R&D.

With L. oar-ter a technical challenges of current high-priority P5 projects subsiding, the HEP Detector . Torogram aims to shift more emphasis towards building a long-term, high-risk high-rewar . ("Blue Sky") R&D portfolio that holds the promise of transformative advances with broad-ranging applications across HEP as well as other fields of science, medicine, and national security. Crucially, the program must take full advantage of the major advances happening in

- Survey the present state of the HEP technology landscape.
- Identify key capabilities & performance requirements.
- Identify technologies to provide or enhance such capabilities.
- Articulate Priority Research Directions to push well beyond the current state of the art, potentially leading to transformative technological advances with broad-ranging applicability.
- Flesh out required R&D efforts with deliverables with notional timelines & key technical milestones. Elucidate the technical infrastructure required to support these efforts.
- Formulate a small set of instrumentation Grand Challenges that could result in game-changing experimental capabilities.

10 Basic Research Needs Study Charge



Department of Energy Office of Science Washington, DC 20585

10 July 2019

MEMORANDUM FOR HELMUT MARSISKE

FROM: GLEN CRAWFORD

DIRECTOR, RESEARCH AND TECY NOLOGY DIVISION OFFICE OF HIGH ENERGY PHY LCS (HEP)

OFFICE OF HIGH ENERGY PHY ACS (HEP)

SUBJECT: Basic Research Needs Study co. n. Detector Remarch and

Development

You should ... 'co-chairs to le id "he study and wor" with them to select the core group of working group le. 'o carry i' unt. 1. 'ady enc. impasses responses to the specific charge elements elucidated. 'w 'ad is expect.' 'he several months to complete. A focal point of the study should inclus. 'workshop, with au-endance beyond the core group, expected to be held in 'a' 2019 time fi. in the Washington, DC area. The study participants are to serve by invitation o.

The HEP Detecto. *D program _ms to develop cutting-edge, novel instrumentation to enable scientific leadership a worldwide experimental program that is broadening into new research areas with ever incre: ng demands in sensitivity, scale, and cost. To meet this challenge, HEP 'ms to execute a pro' am appropriately balanced between incremental, near-term, low-risk o to recreate a pro' and transformative. lone-term, high-risk detector R&D.

With L. *ar-ter*. a technical challenges of current high-priority P5 projects subsiding, the HEP Detector. * program aims to shift more emphasis towards building a long-term, high-risk high-rewar. ("Blue Sky") R&D portfolio that holds the promise of transformative advances with broad-ranging applications across HEP as well as other fields of science, medicine, and national security. Crucially, the program must take full advantage of the major advances happening in

To meet this challenge, HEP aims to execute a program appropriately balanced between incremental, near-term, low-risk detector R&D and transformative, long-term, high-risk detector R&D

Along with the five science drivers, the 2014 P5 report identifies the importance of Instrumentation R&D in one of its highest level recommendations where it calls for a "balanced mix of short term and long-term R&D" in the current era.

The BRN does:
describe SCIENCE
OPPORTUNITIES
& TECHNOLOGIES
TO REALIZE THEM

The BRN does not: rank PRD opportunities

10 Basic Research Needs Study Charge



Department of Energy Office of Science Washington, DC 20585 10 July 2019

MEMORANDUM FOR HELMUT MARSISKE

FROM: GLEN CE

GLEN CRAWFORD

DIRECTOR, RESEARCH AND TECT NOLOGY DIVISION

OFFICE OF HIGH ENERGY PHY ACS (HEP)

SUBJECT:

Basic Research Needs Study C. r. Detector Re varch and

Development

You should ... 'oo-chairs to Ir all the study and wor' with them to select the core group of working group le. '' ocarry' with... '' wdy enc, impasses responses to the specific charge elements elucidated. 'w '' ald is expecte. ''' we several months to complete. A focal point of the study should includ. 'workshop, with auendance beyond the core group, expected to be held in '' ~ 2019 time fit. 'in the Washington, DC area. The study participants are to serve by invitation to.


The HEP Detecto. 'D program ams to develop cutting-edge, novel instrumentation to enable scientific leadership areas with ever increase with ever increase with ever increase with ever increase and genands in sensitivity, scale, and cost. To meet this challenge, HEP was to execute a pro- am appropriately balanced between incremental, near-term, low-risk of total Rep and to: sformative, long-term, high-risk detector R&D.

With L. *ar-ter-a technical challenges of current high-priority P5 projects subsiding, the HEP Detector. *D-program aims to shift more emphasis towards building a long-term, high-risk high-rewar. ("Blue Sky") R&D portfolio that holds the promise of transformative advances with broad-ranging applications across HEP as well as other fields of science, medicine, and national security. Crucially, the program must take full advantage of the major advances happening in


Report should speak to the scientific community, the public, and decision makers

Panels Science 2

Science opportunities drive the next generation of experiments.



An instrumentation revolution is critical to future discoveries



Study Process and timeline

8/2019: DOE charged co-Chairs. Conveners, panel members, additional members identified.

Co-Chairs

Bonnie Fleming, Yale Ian Shipsey, Oxford

Cross-Cut Panel

Marcel Demarteau, ORNL James Fast, JLab Sunil Golwala, CalTech Young-Kee Kim, Chicago Abraham Seiden, UCSC

Energy Frontier

James Hirschauer, Fermilab (Lead) Gabriella Sciolla, Brandeis (Lead) Michael Begel, Brookhaven Meenakshi Narain, Brown

Neutrinos

Ornella Palamara, Fermilab (Lead) Kate Scholberg, Duke (Lead) Daniel Dwyer, Berkeley Lab Amy Connolly, OSU

Physics Panels

Dark MatterJodi Cooley, S

Jodi Cooley, SMU (Lead)
Dan McKinsey, Berkeley (Lead)
Andrew Sonnenschein, Fermilab
Reyco Henning, UNC

Cosmic Acceleration

Clarence Chang, Argonne (Lead) Brenna Flaugher, Fermilab (Lead) Kyle Dawson, Utah Laura Newburgh, Yale

Explore the Unknown

Sarah Demers, Yale (Lead) Monica Pepe-Altarelli, CERN, EONR (Lead) Matthew Reece, Harvard Nicola Serra, Universität Zürich

Technology Panels

Calorimetry

Francesco Lanni, Brookhaven (Lead) Roger Rusack, Minnesota (Lead) Nural Akchurin, Texas Tech Sarah Eno, UMD Paolo Rumerio, Alabama Ren-Yuan Zhu, CalTech

Noble Liquids

Roxanne Guenette, Harvard (Lead) Jocelyn Monroe, U London (Lead) Jennifer Raaf, Fermilab Andrea Pocar, UMass Jonathan Asaadi, UT, Arlington Hugh Lippincott, UCSB

Photodetectors

Lindley Winslow, MIT (Lead)
Peter Krivzan, ULJ / JSI (Lead)
Graham Giovanetti, Williams College
Adriana Lita, NIST
Felix Sefkow, DESY

Quantum Sensors

Andrew Geraci, Northwestern (Lead) Kent Irwin, Stanford (Lead) Gretchen Campbell, JQI/UMD Alexander Sushkov, BU Ronald Walsworth, Harvard Anna Grassellino, Fermilab

Readout & ASICS

Gabriella Carini, Brookhaven (Lead) Mitch Newcomer, Penn (Lead) Angelo Dragone, SLAC Maurice Garcia-Sciveres, Berkeley Lab Terri Shaw, Fermilab

Solid State & Tracking

Julia Thom-Levy, Cornell

Marina Artuso, Syracuse (Lead) Carl Haber, Berkeley Lab (Lead) Alessandro Tricoli, Brookhaven Petra Merkel, Fermilab

TDAQ

Darin Acosta, Florida (Lead)
Tulika Bose, UW, Madison (Lead)
Wesley Ketchum, Fermilab
Jinlong Zhang, Argonne
Paul O'Connor, Brookhaven
Georgia Karagiorgi. Columbia

Balance is important

Co-Chairs

Bonnie Fleming, Yale Ian Shipsey, Oxford

Cross-Cut Panel

Marcel Demarteau, ORNL James Fast, JLab Sunil Golwala, CalTech Young-Kee Kim, Chicago Abraham Seiden, UCSC

Physics Panels

Energy Frontier

James Hirschauer, Fermilab (Lead) Gabriella Sciolla, Brandeis (Lead) Michael Begel, Brookhaven Meenakshi Narain, Brown

Neutrinos

Ornella Palamara, Fermilab (Lead) Kate Scholberg, Duke (Lead) Daniel Dwyer, Berkeley Lab Amy Connolly, OSU

Dark Matter

Jodi Cooley, SMU (Lead) Dan McKinsey, Berkeley (Lead) Andrew Sonnenschein, Fermilab Reyco Henning, UNC

Cosmic Acceleration

Clarence Chang, Argonne (Lead) Brenna Flaugher, Fermilab (Lead) Kyle Dawson, Utah Laura Newburgh, Yale

Explore the Unknown

Sarah Demers, Yale (Lead) Monica Pepe-Altarelli, CERN, EONR (Lead) Matthew Reece, Harvard Nicola Serra, Universität Zürich

Technology Panels

Calorimetry

Francesco Lanni, Brookhaven (Lead) Roger Rusack, Minnesota (Lead) Nural Akchurin, Texas Tech Sarah Eno, UMD Paolo Rumerio, Alabama Ren-Yuan Zhu, CalTech

Noble Liquids

Roxanne Guenette, Harvard (Lead) Jocelyn Monroe, U London (Lead) Jennifer Raaf, Fermilab Andrea Pocar, UMass Jonathan Asaadi, UT, Arlington Hugh Lippincott, UCSB

Photodetectors

Lindley Winslow, MIT (Lead)
Peter Krivzan, ULJ / JSI (Lead)
Graham Giovanetti, Williams College
Adriana Lita, NIST
Felix Sefkow, DESY

Quantum Sensors

Andrew Geraci, Northwestern (Lead) Kent Irwin, Stanford (Lead) Gretchen Campbell, JQI/UMD Alexander Sushkov, BU Ronald Walsworth, Harvard Anna Grassellino, Fermilab

Readout & ASICS

Gabriella Carini, Brookhaven (Lead)
Mitch Newcomer, Penn (Lead)
Angelo Dragone, SLAC
Maurice Garcia-Sciveres, Berkeley Lab
Terri Shaw, Fermilab
Julia Thom-Levy, Cornell

Solid State & Tracking

Marina Artuso, Syracuse (Lead) Carl Haber, Berkeley Lab (Lead) Alessandro Tricoli, Brookhaven Petra Merkel, Fermilab

TDAQ

Darin Acosta, Florida (Lead)
Tulika Bose, UW, Madison (Lead)
Wesley Ketchum, Fermilab
Jinlong Zhang, Argonne
Paul O'Connor, Brookhaven
Georgia Karagiorgi, Columbia

Balance is important

Co-Chairs

Bonnie Fleming, Yale lan Shipsey, Oxford

Cross-Cut Panel

Marcel Demarteau, ORNL James Fast, JLab Sunil Golwala, CalTech Young-Kee Kim, Chicago Abraham Seiden, UCSC

Energy Frontier

James Hirschauer, Fermilab (Lead) Gabriella Sciolla, Brandeis (Lead) Michael Begel, Brookhaven Meenakshi Narain, Brown

Neutrinos

Ornella Palamara, Fermilab (Lead) Kate Scholberg, Duke (Lead) Daniel Dwyer, Berkeley Lab Amy Connolly, OSU

Physics Panels

Dark Matter

Jodi Cooley, SMU (Lead) Dan McKinsey, Berkeley (Lead) Andrew Sonnenschein, Fermilab Reyco Henning, UNC

Cosmic Acceleration

Clarence Chang, Argonne (Lead) Brenna Flaugher, Fermilab (Lead) Kyle Dawson, Utah Laura Newburgh, Yale

Explore the Unknown

Sarah Demers, Yale (Lead) Monica Pepe-Altarelli, CERN, EONR (Lead) Matthew Reece, Harvard Nicola Serra, Universität Zürich

Technology Panels

Calorimetry

Francesco Lanni, Brookhaven (Lead) Roger Rusack, Minnesota (Lead) Nural Akchurin, Texas Tech Sarah Eno, UMD Paolo Rumerio, Alabama Ren-Yuan Zhu, CalTech

Noble Liquids

Roxanne Guenette, Harvard (Lead) Jocelyn Monroe, U London (Lead) Jennifer Raaf, Fermilab Andrea Pocar, UMass Jonathan Asaadi, UT, Arlington Hugh Lippincott, UCSB

Photodetectors

Lindley Winslow, MIT (Lead)
Peter Krivzan, ULJ / JSI (Lead)
Graham Giovanetti, Williams College
Adriana Lita, NIST
Felix Sefkow, DESY

Quantum Sensors

Andrew Geraci, Northwestern (Lead) Kent Irwin, Stanford (Lead) Gretchen Campbell, JQI/UMD Alexander Sushkov, BU Ronald Walsworth, Harvard Anna Grassellino, Fermilab

Readout & ASICS

Gabriella Carini, Brookhaven (Lead)
Mitch Newcomer, Penn (Lead)
Angelo Dragone, SLAC
Maurice Garcia-Sciveres, Berkeley Lab
Terri Shaw, Fermilab
Julia Thom-Levy, Cornell

Solid State & Tracking

Marina Artuso, Syracuse (Lead) Carl Haber, Berkeley Lab (Lead) Alessandro Tricoli, Brookhaven Petra Merkel, Fermilab

TDAQ

Darin Acosta, Florida (Lead)
Tulika Bose, UW, Madison (Lead)
Wesley Ketchum, Fermilab
Jinlong Zhang, Argonne
Paul O'Connor, Brookhaven
Georgia Karagiorgi, Columbia

Study Process and timeline

- Fall 2019: Regular telecons began to conduct the ground work leading up to the December BRN workshop
- Important input:

arXiv.org > physics > arXiv:1908.00194

Physics > Instrumentation and Detectors

New Technologies for Discovery

A report of the 2018 DPF Coordinating Panel for Advanced Detectors (CPAD) Community Workshop

```
Z. Ahmed,<sup>a</sup> A. Apresyan,<sup>b</sup> M. Artuso,<sup>a</sup> P. Barry,<sup>d</sup> E. Bielejec,<sup>e</sup> F. Blaszczyk,<sup>ah</sup> T. Bose,<sup>f</sup> D. Braga,<sup>b</sup> S.A. Charlebois,<sup>ai</sup> A. Chatterjee,<sup>j</sup> A. Chavarria,<sup>g</sup> H.-M. Cho,<sup>a</sup> S. Dalla Torre,<sup>h</sup> M. Demarteau*,<sup>ar</sup> D. Denisov,<sup>v</sup> M. Diefenthaler,<sup>ag</sup> A. Dragone,<sup>a</sup> F. Fahim,<sup>b</sup> C. Gee,<sup>af</sup> S. Habib,<sup>d</sup> G. Haller,<sup>a</sup> J. Hogan,<sup>f</sup> B.J.P. Jones,<sup>j</sup> M. Garcia-Sciveres,<sup>k</sup> G. Giacomini,<sup>v</sup> K. Gilmore,<sup>an,p</sup> G.K. Giovanetti,<sup>al</sup> D. Glenzinski,<sup>b</sup> S. Gleyzer,<sup>l</sup> A.H. Goldan,<sup>ad</sup> S. Gollapinni,<sup>m</sup> C. Grace,<sup>k</sup> R. Guenette,<sup>n</sup> O. Gutsche,<sup>b</sup> U. Heintz,<sup>o</sup> S.A. Hertel,<sup>aj</sup> N. R. Hutzler,<sup>ac</sup> S. Kolkowitz,<sup>f</sup> T. Kovachy,<sup>ak</sup> F. Leonard,<sup>am</sup> R. Lipton,<sup>b</sup> M. Liu,<sup>b</sup> J.F. Low,<sup>l</sup> P. Madigan,<sup>s,k</sup> S. Malik,<sup>aq</sup> J. Mates,<sup>p</sup> Y. Mei,<sup>k</sup> P. Merkel,<sup>b</sup> T. Mohayai,<sup>b</sup> A. Nomerotski,<sup>v</sup> E. Oliveri,<sup>q</sup> K. Palladino,<sup>f</sup> E. Pantic,<sup>r</sup> A. Para,<sup>b</sup> K. Perez,<sup>aa</sup> M. Pyle,<sup>s</sup> P. Riedler,<sup>q</sup> L. Ropelewski,<sup>q</sup> R. Rusack,<sup>t</sup> M. Schleier-Smith,<sup>f</sup> I. Shipsey*,<sup>as</sup> K. Scholberg,<sup>u</sup> B. A. Schumm,<sup>af</sup> A. Slosar,<sup>v</sup> W. Smith,<sup>f</sup> B. Surrow,<sup>w</sup> A. O. Sushkov,<sup>ah</sup> A. Suzuki,<sup>k</sup> M. Szydagis,<sup>ae</sup> D. Temples,<sup>ak</sup> J. Thom,<sup>z</sup> M. Titov,<sup>y</sup> L. Tvrznikova,<sup>aa</sup> E. Usai,<sup>a</sup> R. Van Berg,<sup>z</sup> V. Velan,<sup>s</sup> D.W. Whittington,<sup>c</sup> L. Winslow,<sup>aa</sup> T. Wongjirad,<sup>ab</sup> Q. Xia<sup>ap</sup> J. Xie,<sup>d</sup> Z.F. You,<sup>f</sup> A. Zani,<sup>q</sup> J. Zhang,<sup>d</sup> R.Y. Zhu.<sup>ac</sup>
```

 BRN Interim report laid the foundations of the panel's work and informed interactions at CPAD2019 arXiv.org > physics > arXiv:1908.00194

Physics > Instrumentation and Detectors

New Technologies for Discovery

A report of the 2018 DPF Coordinating Panel for Advanced Detectors (CPAD) Community Workshop



Goal: provide a community forum to communicate with the BRN panel, timed to be just before the BRN workshop.

12 plenary speakers Day 1 were mostly BRN panel members. Townhalls provided further opportunities for dialog

Report of the 2018 CPAD workshop was a primary input to the 2019 DOE BRN study on HEP Detector R&D

| | Monona Terrace Convention Center | 08:00 - 09:00 |
|----|---|-----------------------|
| 00 | Welcome | Kimberly PALLADINO 🗎 |
| | Monona Terrace Convention Center | 09:00 - 09:15 |
| | The Higgs as a tool for discovery | Jim HIRSCHAUER 📋 |
| | Monona Terrace Convention Center | 09:15 - 09:30 |
| | Dark Matter | Jodi COOLEY 🛅 |
| | Monona Terrace Convention Center | 09:30 - 09:45 |
| | DE and Inflation Instrumentation BRN working group | Dr. BRENNA FLAUGHER 📋 |
| | Monona Terrace Convention Center | 09:45 - 10:00 |
| 00 | Exploring the Unknown | Sarah DEMERS |
| | Monona Terrace Convention Center | 10:00 - 10:15 |
| | Neutrinos and Neutrino Mass | Amy CONOLLY |
| | Monona Terrace Convention Center | 10:15 - 10:30 |
| | Photodetectors | Jungi XIE |
| | Monona Terrace Convention Center | 11:40 - 11:52 |
| | Quantum Sensors | Tim KOVACHY |
| 00 | Monona Terrace Convention Center | 11:52 - 12:04 |
| | Noble Liquid detectors | Dr. Hugh LIPPINCOTT |
| | Monona Terrace Convention Center | 12:04 - 12:16 |
| | Trigger and DAQ | Prof. Tulika BOSE 📋 |
| | Solid State and tracking | Carl HABER 📋 |
| | Meeting Rooms K-R, Monona Terrace Convention Center | 13:30 - 13:45 |
| | Calorimetry | Roger RUSACK |
| | Meeting Rooms K-R, Monona Terrace Convention Center | 13:45 - 14:00 |
| 00 | Readout and ASICs | Mitch NEWCOMER 📄 |
| | Meeting Rooms K-R, Monona Terrace Convention Center | 14:00 - 14:15 |
| | Plenary Townhall: BRN process | |
| | Monona Terrace Convention Center | 14:15 - 15:00 |

HEP Basic Research Needs Workshop on Detector Research and Development


December 11 - 14, 2019

Hilton Washington DC/Rockville Hotel & Executive Meeting Center

1750 Rockville Pike


Rockville, MD 20852

- Attendees: Panel conveners and members, Agencies (DOE, NSF)
- Plenaries on the first day live streamed to community
- Parallel sessions and working groups to work through the substance and first draft of the report
- Post workshop: spring 2020 streamlining and refining
- July 2020 Presentation to HEPAP
- September 2020 Report published




Physics to Technologies

PRDs (Priority Research Directions)




DIS2021-Shipsey 34

Physics to



DIS2021-Shipsey 35



Four Grand Challenges encompass this Instrumentation revolution

- Advancing HEP detectors to new regimes of sensitivity: To make the unmeasurable measurable will require the development of sensors with exquisite sensitivity with the ability to distinguish signal from noise.... Research will be needed to develop these sensors with maximal coupling to the quanta to be sensed and push their sensitivities to ultimate limits.
- Using Integration to enable scalability for HEP sensors: Future HEP detectors for certain classes of experiments will require massive increases in scalability to search for and study rare phenomena ... A key enabler of scalability is integration of many functions on, and extraction of multidimensional information from, these innovative sensors.
- Building next-generation HEP detectors with novel materials & advanced techniques: Future HEP detectors will have requirements beyond what is possible with the materials and techniques which we know. This requires identifying novel materials ... that provide new properties or capabilities and adapting them & exploiting advanced techniques for design & manufacturing.
- Mastering extreme environments and data rates in HEP experiments:

 Future HEP detectors will involve extreme environments and exponential increases in data rates to explore elusive phenomena. ... To do so requires the intimate integration of intelligent computing with sensor technology.

Instrumentation Development Ecosystem

Key to the success of this tool driven revolution are **people**, **facilities and resources**, and **connections and collaborations**

- 1. Advanced workforce
- 2. Unique capabilities and facilities
- 3. Connections to other programs, other offices, other agencies, private foundations, commercial partners, global collaborations

A Commitment to Equality, Diversity and Inclusion in HEP Instrumentation R&D

Excellence and innovation come most effectively from diverse teams of people.



A Commitment to Equality, Diversity and Inclusion in HEP Instrumentation R&D

Excellence and innovation come most effectively from diverse teams of people.



To accomplish the best science, we must commit, as a community, to action, to overcome the social injustices in our own backyard, and realize the impact of a diverse workforce. We must find, develop, and invent new ways to attract, encourage, recruit, and support a diverse community. We must enact an inclusive environment within instrumentation and within particle physics at all levels and across the areas in which we work and that we touch including academia, in universities and national laboratories, and in industry.

Some small steps are being taken to draw young people from diverse backgrounds to instrumentation.

- Outreach programs at universities and national labs and to the public
- Undergraduate research opportunities (NSF Research Experience for Undergraduates)
- Graduate Instrumentation Research Awards (award winners and honorable mentions, 2018 GIRA)



Recipients of the DPF
Instrumentation Awards
2019. Left to Right
Hanguo Wang (UCLA),
Ettore Segreto, and Anna
Amelia Machado (both of
the University of
Campinas in Brazil)



Some small steps are being taken to draw young people from diverse backgrounds to instrumentation.

- Outreach programs at universities and national labs and to the public
- Undergraduate research opportunities (NSF Research Experience for Undergraduates)
- Graduate Instrumentation Research Awards (award winners and honorable mentions, 2018 GIRA)



The scale of these initiatives must be dramatically increased and new initiatives developed with urgency.

Recipients of Instrumentation

2019. Left to Right
Hanguo Wang (UCLA),
Ettore Segreto, and Anna
Amelia Machado (both of
the University of
Campinas in Brazil)



Workforce requirements

Many areas require expertise and cross-disciplinary work (electronics, CS, DAQ, Mechanical engineering, cryogenic systems, composites design and fabrication, microfab and assembly, analytic chemistry, materials science, ...

To succeed in creating tools and new technologies, we need to succeed in excellence in the current and next generation of people

- diverse pipeline (in US, international)
- University/lab partnerships
- connections to other disciplines
- appropriate recognition

These experts, in turn, educate the next generation in advanced HEP instrumentation techniques and development transforming not only HEP but other fields too.

Instrumentation Development Ecosystem

Key to the success in this tool revolution are **people**, **facilities and resources**, and **connections and collaborations**

- 1. Advanced workforce
- 2. Unique capabilities and facilities
- 3. Connections to other programs, other offices, other agencies, private foundations, commercial partners, global collaborations

Maintain core facilities

- SiDET
- Noble Liquid Test Facility
- Micro Systems Lab
- FTBTF
- Test beams at SLAC



| | Higgs and Energy Frontier | Neutrinos | Dark Matter | Cosmic Acceleration | Unknown |
|--|---------------------------------|-----------|----------------|---------------------|----------|
| Irradiation, ionizing and non-ionizing | ✓ | ✓ | | | ✓ |
| Test Beams | ✓ | ✓ | | | ✓ |
| Test Stands at Ultra- low Temperature | | | ✓ | ✓ | ✓ |
| Calibration Facilities | ✓ | ✓ | ✓ | ✓ | ✓ |
| Low Background Materials and Assay | | ✓ | ✓ | | ✓ |
| Ultra-light Composites | ✓ | | | | ✓ |
| Novel CCD Development | | | ✓ | ✓ | |
| Superconducting Detector and Device Foundry | | | ✓ | ✓ | |
| Microelectronics Engineering and Foundry Access | √ | ✓ | ✓ | √ | ✓ |
| Simulation Framework | √ | √ | √ | √ | √ |

Table 23: Capability needs for the five science drivers.

Develop new capabilities, collaborate where possible

45



Instrumentation Development Ecosystem

Key to the success in this tool revolution are **people**, **facilities and resources**, and **connections and collaborations**

- 1. Advanced workforce
- 2. Unique capabilities and facilities
- 3. Connections to other programs, other offices, other agencies, private foundations, commercial partners, global collaborations

R&D connections and collaborations

- Connections within OHEP, with the DOE Office of Science, between federal agencies, Universities & National Labs, with industry, & philanthropic foundations
- Need organizational structures to bring together technical areas
- Rotating leadership, National labs provide homes








- Models: CERN R&D collaborations,
- DOE NNSA R&D consortia


NNSA/DNN University Consortia



Connections to other disciplines: Benefits to Society

The development of the manufacturing process of BGO crystals for the calorimeter of the L3 experiment at the LEP collider at CERN (left) has contributed significantly to the advancement of Positron Emission Tomography (PET) scanners





(photo credit: CERN and S.R. Cherry/U.C. Davis)





The development of large-area hybrid pixel detectors for high energy physics experiments led to the realization of the potential of this new technology to provide noise-hit-free single-photon counting impactful for development of sophisticated integrated circuits with timing. The circuit is being used in medical imaging, X-ray science, materials analysis, space dosimetry and climate studies among others

The instrumentation plan described in this report will lead to the development of new technologies that hold the promise to be as broadly applicable and equally transformative.

Higgs and the Energy Frontier:

Next generation energy frontier colliders & detectors are precision measurement machines & discovery machines. The transformative physics goals include 4 inspiring & distinct directions:


Higgs properties with sub-percent precision

Higgs self-coupling with 5% precision

Higgs connection to dark matter

New particles and phenomena at multi-TeV scale

Technical Requirements to enable the physics program for Higgs and the Energy Frontier and map to Priority Research Directions.




| Science | Measurement | Technical Requirement | PRD |
|---|---|--|--|
| Higgs properties with sub-percent precision Higgs self-coupling with 5% precision | TR 1.1: Tracking for e^+e^- | TR 1.1.1: $p_{\rm T}$ resolution: $\sigma_{p_{\rm T}}/p_{\rm T}=0.2\%$ for tracks with $p_{\rm T}<100$ GeV, $\sigma_{p_{\rm T}}/p_{\rm T}^2=2\times10^{-5}/{\rm GeV}$ for tracks with $p_{\rm T}<100$ GeV TR 1.1.2: Impact parameter resolution: $\sigma_{r\phi}=5\bigoplus 15\ (p\ [{\rm GeV}]\ \sin^{\frac{3}{2}}\theta)^{-1}\ \mu{\rm m}$ TR 1.1.3: Granularity: $25\times50\ \mu{\rm m}^2$ pixels TR 1.1.4: $5\ \mu{\rm m}$ single hit resolution TR 1.1.5: Per track timing resolution of 10 ps | 18, 19 20, 23 |
| Higgs connection to dark matter | TR 1.2: Tracking for 100 TeV pp | Generally same as e^+e^- (TR 1.1) except TR 1.2.1: Radiation tolerant to 300 MGy and $8\times 10^{17}~\rm n_{eq}/cm^2$ TR 1.2.2: $\sigma_{pr}/p_{\rm T}=0.5\%$ for tracks with $p_{\rm T}<100$ GeV TR 1.2.3: Per track timing resolution of 5 ps rejection and particle identification | 16, 17 18, 19 20, 23 26 |
| New particles and phenomena at multi-TeV scale | TR 1.3: Calorimetry for e^+e^- | TR 1.3.1: Jet resolution: 4% particle flow jet energy resolution TR 1.3.2: High granularity: EM cells of 0.5×0.5 cm ² , hadronic cells of 1×1 cm ² TR 1.3.3: EM resolution: $\sigma_E/E = 10\%/\sqrt{E} \bigoplus 1\%$ TR 1.3.4: Per shower timing resolution of 10 ps | 1, 3, 7, 10, 11, 23 |
| | TR 1.4: Calorimetry for 100 TeV pp | Generally same as e^+e^- (TR 1.3) except TR 1.4.1: Radiation tolerant to 4 (5000) MGy and $3 \times 10^{16} (5 \times 10^{18}) \text{ n}_{eq}/\text{cm}^2$ in endcap (forward) electromagnetic calorimeter TR 1.4.2: Per shower timing resolution of 5 ps | 1, 2, 3 7, 9, 1 11, 16 17, 23 26 |
| | TR 1.5: Trigger and readout | TR 1.5.1: Logic and transmitters with radiation tolerance to 300 MGy and $8 \times 10^{17} \text{ n}_{\rm eq}/\text{cm}^2$ | 16, 17 21, 26 |

Jim Hirschauer Gabriella Sciolla (leads)


Michael Begel Meenakshi Narain

Timeline: Higgs → Technologies to Discovery



Neutrinos

- Push the three-flavor paradigm into the regime of high-precision measurements of all parameters.
- Explore unknown territory in neutrino energy range, types of neutrino sources, and faint source intensities.
- Hunt for evidence of new particles and phenomena in the neutrino sector, and in other sectors using neutrino detectors.



| Science | Measurement | Technical Requirement | PRD |
|---|---|---|---|
| Neutrino mixing matrix unitarity | Measure tau neutrino appearance with high efficiency/purity | TR 2.1: Resolve short tracks (0.1 mm at 10 GeV) in 10 kton detectors | 4,6,10, 11,16,21,22, 23,25,26 |
| Measure neutrinos at macroscopic energies from cosmic distances for BSM searches | Sensitivity to neutrino fluxes $1/\mathrm{km}^2/\mathrm{decade}$ at low energy threshold | TR 2.2: Low power ($\ll 1$ W) digitizers sampling at >3 GHz, triggering at $\mathcal{O}(1)$ S/N | 16,17 |
| Resolve solar/ reactor Δm_{12}^2 tension | Measure solar ⁸ B, hep and neutrino regeneration in the Earth with S/B>1 above a few MeV | TR 2.3: Radiogenic background reduction by a factor of 100-1000 in argon TR 2.4: <1 cm spatial, TR 2.5: <10% energy resolution at kton scale | 4,6,21,22, 23,24,25,26 |
| Measure all flavor components of a supernova burst in real time | Flavor tagging with >90% efficiency, 5-50 MeV; measure CEvNS glow/buzz in large LAr or scintillator | TR 2.3, 2.4, 2.5 TR 2.6: Photodetector efficiency improvement by factor of 10 TR 2.7: Photosensor dark noise reduction by factor of 100 | 4,6,7,9,10, 11,16,21,22 23,24,25,26 |
| BSM physics with sub-MeV (or sub-keV) neutrinos (geoneutrinos, pp neutrinos, solar thermal neutrinos, artificial radioactive sources) | Sensitivity to very low energy nuclear or electronic recoils in real time | TR 2.8: 10 eV nuclear recoil threshold at multi-ton to kton scale TR 2.9 Few degree recoil directionality | 5,6,7,9,11, 12,14,24,25,26 |
| Cosmic relic neutrino background, test of cosmological models | Measure cosmic relic neutrino capture on nuclei | TR 2.10: 10 meV energy resolution at beta endpoint, $\mathcal{O}(1 \text{ kg})$ source with TR 2.11: <10 meV energy loss distortion at endpoint | 12,14,26 |

Table 3: Technical Requirements to enable example neutrino physics topics and map to Priority Research Directions.

Ornella Palamara Kate Scholberg (leads) Amy Connolly Dan Dwyer





Figure II: Timeline of neutrino experiments.

Dark Matter

- Search for WIMP dark matter towards the neutrino floor
- Searching for particle dark matter with low masses
- Searching for wave-like dark matter
- Searching for the annihilation or decay products of dark matter interactions




Jodi Cooley Dan McKinsey (leads)
Reyco Henning Andrew Sonnenshein

| Science | Measurement | Technical Requirement | PRDs |
|---|---|--|---------------------------|
| Test for dark matter particles with mass >1 GeV | Search for nuclear recoils arising from scattering of >1 GeV dark matter with normal matter via spin-independent and spin-dependent couplings to nucleons | Mass 1 - 10 GeV TR 3.1(SI),TR 3.7(SD): Background rate < coherent scattering rate of solar neutrinos TR 3.2(SI), TR 3.8(SD): Target mass ~100 kg TR 3.3(SI), TR 3.9(SD): Energy Threshold: ~100 eV | 5, 6, 24, 25 |
| | | Mass > 10 GeV TR 3.4(SI), TR 3.10(SD): Background rate < coherent scattering rate of atmospheric neutrinos TR 3.5(SI), TR 3.11(SD): Target mass ~100 tonnes TR 3.6(SI), TR 3.12(SD): Energy Threshold: ~10 keV | 6, 7, 9, 11, 25, 26 |


| Science | Measurement | Technical Requirement | PRDs |
|--|---|---|------------|
| Test for peV-neV QCD axion dark matter | Search for peV-neV QCD axion dark matter via axion-nucleon coupling with nuclear magnetic | Near Term: TR 3.21 $P \ge 0.05$ TR 3.23 $N\tau = 10^{24}\mathrm{sec.}$ | 12, 13, 15 |
| | resonance | Long Term: TR 3.22 $P \ge 0.3$ TR 3.24 $N\tau = 10^{25} {\rm sec.}$ | 12, 13, 15 |
| Test for neV- μ eV QCD axion dark matter | Search for neV- μ eV QCD axion dark matter using axion-photon conversion in lumped-element electromagnetic resonators | Near Term: TR 3.25 $Q_L \ge 10^6 {\rm GeV}$ TR 3.27 $\eta \le 20$ TR 3.29 $BV \ge 4 {\rm T \cdot m^3}$ | 12, 15 |
| | | Long Term: TR 3.26 $Q_L \ge 10^8$ TR 3.28 $\eta \le 0.1$ TR 3.30 $BV \ge 10\mathrm{T\cdot m^3}$ | 12, 15 |
| Test for μeV -meV QCD axion dark matter | Search for µeV-meV QCD axion dark matter using axion-photon conversion in cavity electromagnetic resonators | Near Term TR 3.31 $Q_C \ge 10^5$ TR 3.33 $\eta \le 1$ TR 3.35 $B \ge 10$ T, $V \ge 100l$ | 12, 15 |
| | | $\begin{array}{l} \textbf{Long Term:} \\ \text{TR } 3.32 \ Q_C \geq 10^6 \\ \text{TR } 3.34 \ \eta \leq 10^{-6} \\ \text{TR } 3.36 \ B \geq 30 \ \text{T}, \ V \geq 1l \end{array}$ | 12, 15 |


Cosmic Acceleration: Dark Energy & Inflation

- Drive cosmological measurements to new spatial and temporal scales
- Explore the properties of inflation, dark energy, and dark matter
- Study neutrino physics in a context complementing terrestrial techniques
- Test our concordance cosmological model in new regimes



| Science Goal | Measurement | Technical Requirement | PRD |
|--|--|---|-------------------|
| Fully sample the epoch of late-time cosmic acceleration | 500M Galaxy spectra (R~3000) to z<4 | For Optical/IR spectroscopy TR 4.1: Sensitivity at wavelengths beyond the 1eV Silicon cutoff. TR 4.2: Ten-fold increase in multi- plexing relative to current experiments | 7, 11, 26 |
| Distinguish between single vs. multi-field inflation by measuring f_{NL} down to 1 | Multiple Intensity mapping surveys to measure flux from 2.9B galaxies to z<6 | For 21-cm Intensity Mapping: TR 4.3: Pico-second timing synchronization across ~km TR 4.4: Direct digitization and real-time calibration | 21, 22, 23, 26 |
| | | For mm-wave Intensity Mapping: TR 4.5: On-chip mm spectrometers with R>200 TR 4.6: Fabrication and readout of 1M detectors | 7, 8, 26 |






Clarence Chang Brenna Flaugher (leads)

Kyle Dawson Laura Newburgh


Explore the Unknown

- Precision measurements in heavy flavor decays
- Searching for charged lepton flavor violation in rare decays of muons and kaons
- Tests of CP violation through electric dipole moment searches
- Probes of the dark sector and hunts for new fundamental forces



| Science | Timescale | Technical Requirement | PRD |
|--|----------------|--|-------------------------|
| | medium term | TR 5.1: Timing resolution at the level of $10-30$ ps per hit in the silicon-pixel vertex detectors and $10-30$ ps per track for both PID detectors (RICH, TORCH) and electromagnetic calorimeters | 2, 10, 18 |
| Search for new physics though rare flavor interactions | medium term | TR 5.2: Development of radiation-hard, fast and cost-effective photosensors for TORCH and RICH detectors and tracking systems with optical readout | 9, 11 |
| | medium term | TR 5.3: Development of the next generation ASICS to extract the large data rate (and possibly pre-process it) out of inner pixel layer detectors in a very challenging radiation environment | 16, 17 |
| | medium term | TR 5.4: Radiation-hard silicon pixel detectors (fluences of $5 \times 10^{16} n_{\rm eq}/{\rm cm}^2)$ | 18, 20 |
| Tests of the CKM quark mixing matrix description | medium term | TR 5.5: Cost-effective electromagnetic calorimeter with granularity of typically 2×2 cm ² , resolution of $\frac{\sigma(E)}{E} \sim \frac{10\%}{\sqrt{E}} \oplus 1\%$ and timing resolution of a few tens of ps; total radiation dose of ~ 200 Mrad | 1 |
| | medium term | TR 5.6: Real-time processing of large amount of data (400-500 Tb/sec) and development of radiation-hard, high-rate optical links, with tight constraints of low-power consumption and low mass | 16, 17, 21, 22 |
| | long term | TR 5.7: Fast-timing resolution at the level of 1 ps per track for $\pi/K/p$ separation up to 50 GeV | 3, 10 |
| | long term | TR 5.8: Further ASICS development to extract and pre-process on detector the large data rate of inner layers detectors in an extreme radiation environment | 16, 17 |
| Studies of Lepton Flavor Universality | long term | TR 5.9: Radiation-hard, ultra-fast silicon pixel detectors (fluences of $10^{18}~\rm n_{eq}/cm^2)$ | 18, 19, 20 |
| | long term | TR 5.10: Very high granularity calorimeters preserving an energy resolution of $\frac{\sigma(E)}{E}\sim \frac{10\%}{\sqrt{E}}$ | 1, 2, 7, 9 |
| | | TR 5.11: Real-time processing of large amount of data | 16, 17, |
| Sarah Demers Monica Pepe-Altarelli (leads) Matthew Reece Nicola Serra | | | |

Technology Panels Priority Research Directions Thrusts delineated Actionable Research plans



DIS2021-Shipsey 61


| Priority Research Direction | Technical Requirements |
|--|---------------------------------|
| PRD 1: Enhance calorimetry energy resolution for precision elec- | TR 1.3, TR 1.4, TR 5.1, TR 5.5, |
| troweak mass and missing-energy measurements | TR 5.10 |
| PRD 2: Advance calorimetry with spatial and timing resolution | TR 1.4, TR 5.7 |
| and radiation hardness to master high-rate environments | |
| PRD 3: Develop ultrafast media to improve background rejection | TR 1.3, TR 1.4, TR 5.7 |
| in calorimeters and improve particle identification | |

Connections outside of HEP:

- The detection of photons, electrons, and hadrons beyond HEP. Eg: experiments at EIC
- Development of organic scintillators for medicine and national security

Facilities and Capabilities (existing and needed)

- Detailed, reliable simulation studies (GEANT4)
- Irradiation facilities to qualify materials, test beams
- Characterizing precision timing systems.
- Studies of data rate, rad tolerance, improved or alternate power delivery systems.
- Expertise: Research scientists at universities




| Priority Research Direction | Technical Requirements |
|--|--|
| PRD 4: Enhance and combine existing modalities to in- | TR 1.3.3, 2.1, 2.4, 2.5, 2.7, 2.9, 3.3, 3.6, |
| crease signal-to-noise and reconstruction fidelity | 3.9, 3.12, 3.13, 3.15, 3.17, 3.19 |
| PRD 5: Develop new modalities for signal detection | |
| PRD 6: Improve the understanding of detector micro- | TR 2.8, 2.9, 3.3, 3.6, 3.9, 3.12, 3.13, |
| physics and characterization to increase signal-to-noise and | 3.15, 3.17, 3.19 |
| reconstruction fidelity | |
| PRD 25: Advance material purification and assay methods | TR 2.3, 3.1, 3.4, 3.7, 3.10 |
| to increase sensitivity | |
| PRD 26: Addressing challenges in scaling technologies | TR 2.1, 2.3, 2.4, 2.7, 2.9, 3.2, 3.5, 3.8, |
| | 3.11, 3.14, 3.16, 3.18, 3.20, 3.45a, 3.45b |


Connections outside of HEP:

- Double beta decay experiments (NP)
- Impact on Astrophysics (eg: SN and solar nus)
- Dedicated R&D for medical imaging

Facilities and Capabilities (existing and needed)

- Low background screening
- Cryogenic platforms (materials, optical properti HV....)
- Test beams
- Engineering expertise





Roxanne Guenette Jocelyn Monroe (Leads)


Jennifer Raaf Andrea Pocar Jonathan Asaadi Hugh Lippincott

| Priority Research Direction | Technical Requiremen | ts | |
|--|---|-------|--|
| PRD 7: Extend wavelength range and develop new single-photon counters to enhance photodetector sensitivity | TR 3.6, TR 4.1, | a) b) | |
| PRD 8: Advance high-density spectroscopy and polarimetry to extract all photon properties | TR 4.1, TR 4.2 | | |
| PRD 9: Adapt photodetectors for extreme environments | TR 1.4, TR 2.3, TR 2.9, TR 2.1 | d) | |
| PRD 10: Design new devices and architectures to enable picosecond timing and event separation | TR 5.10, TR 5.1 TR 1.3, TR 1.4, TR 2.7, TR 4.3, | | |
| PRD 11: Develop new optical coupling paradigms for enhanced or dynamic light collection | TR 1.3, TR 1.4, TR 2.7, TR 2.8, ' TR 3.5, TR 3.6, | f) | |
| | | | |

Connections outside of HEP

- physics experiments and detectors at the light sources and in Astronomy
- Time-Of-Flight (TOF) PET medical imaging, biology, quantum computers, national security Facilities and Capabilities
 - close connections to industry for fabrication of devices and the procurement of materials.
 - new infrastructure through upgrades at existing DOE facilities or partnerships with other federal facilities and industry. (eg: Ge CCD R&D, development of readout and ASICs

Priority Research Direction Technical Requirements $\overline{\mathrm{TR}}$ PRD 12: Advance quantum devices to meet and surpass the Standard Quan-TR TRtum Limit PRD 13: Enable the use of quantum TR TR ensembles and sensor networks for fun-TRdamental physics $\overline{\mathrm{TR}}$ PRD 14: Advance the state of the art in low-threshold quantum calorimeters TRTR PRD 15: Advance enabling technolo-TRgies for quantum sensing TR TRTR




Connections outside of HEP:


quantum information science, quantum computing, materials science, and biology

Facilities and Capabilities (existing and needed

- Large volume high field magnets in solenoidal and toroidal geometries
- Faster turnaround, cheaper, larger mK dilution refrigerators



Andrew Geraci Kent Irwin (Leads) Gretchen Campbell
Alexander Sushkov Ronald Walsworth Anna Grassellino



Connections outside of HEP:

Instrumentation for **Basic Energy** Sciences, NASA, stockpile stewardship program


Facilities and Capabilities (existing and needed)


- Foundry access, including design tools and third party intellectual property
- Long-term, diverse, **HEP** workforce
- Collaboration with other sponsors


PRD PRD 18: Develop high spatial resolution pixel detectors with high per-pixel time resolution to resolve individual interactions in high-collision-density environments

Technical Requirements Thrust

Thrust 1:Lepton colliders, re-







PRD 19: Adapt new materials and fabrication/integration techniques for particle tracking

dustrial partnerships

Thrust 2: Development of rea out electronics matched to r sensor characteristics, includ new processing such as integration

PRD 20: Realize scalable. irreducible-mass trackers

Highly integrat Thrust 1: monolithic, active sensors

Thrust 2: Scaling of low-m detector system

Thrust 3: Systems for spec applications: space-based tra detectors and dedica

Marina Artuso Carl Haber (Leads) Alessandro Tricoli Petra Merkel




Connections outside of HEP:

Nuclear (eg: EIC), astroparticle, medical, materials, homeland security science and engineering.

Facilities and Capabilities (existing and needed)

- specialized infrastructure: test beam and irradiation facilities, silicon processing labs, electronic packaging and assembly, metrology, and composites fabrication facilities
- engineering expertise in ASIC design and test, simulation, verification, and low power systems, and mechanical design and composite fabrication.



Connections outside of HEP:

- DOE Nuclear Physics and DOE Basic Energy Sciences.
- Machine-learning and implementation overlap with technology industry: Aeronautics, smart power grids, autonomous vehicles...

Facilities and Capabilities (existing and needed)

- partnerships between U.S. national laboratories and universities for tool, ASIC, and TDAQ development
- irradiation facilities, integration test facilities

Darin Acosta Tulika Bose (Leads)
Wesley Ketchum Jinlong Zhang Paul O'Connor Georgia Karagiorgi



MPGD Technologies @ RD51 Collaboration

MPGDs in the LHC Experiments:

| Experiment / Timescale | Application Domain | MPGD Technology | Total detector size / Single module size | Operation Characteristics / Performance | Special Requirements / Remarks |
|-----------------------------------|---|-----------------------------------|---|--|---|
| COMPASS TRACKING > 2002 | Fixed Target Experiment (Tracking) | 3-GEM Micromegas w/ GEM preampl. | Total area: 2.6 m ² Single unit detect: 0.31x0.31 m ² Total area: ~ 2 m ² Single unit detect: 0.4x0.4 m ² | Max.rate: ~100kHz/mm² Spatial res.: ~70-100μm (strip), ~120μm (pixel) Time res.: ~ 8 ns Rad. Hard.: 2500 mC/cm² | Required beam tracking (pixelized central / beam area) |
| TOTEM TRACKING: > 2009 | Hadron Collider / Forward Physics (5.3≤ η ≤ 6.5) | 3-GEM (semicircular shape) | Total area: ~ 4 m² Single unit detect: up to 0.03m² | Max.rate:20 kHz/cm ² Spatial res.: ~120μm Time res.: ~ 12 ns Rad. Hard.: ~ mC/cm ² | Operation in pp, pA and AA collisions. |
| LHCb MUON DETECTOR > 2010 | Hadron Collider / B-physics (triggering) | 3-GEM | Total area: ~ 0.6 m ² Single unit detect: 20-24 cm ² | Max.rate:500 kHz/cm² Spatial res.: ~ cm Time res.: ~ 3 ns Rad. Hard.: ~ C/cm² | Redundant triggering |
| COMPASS RICH UPGRADE > 2016 | Fixed Target Experiment (RICH - detection of single VUV photons) | Hybrid (THGEM + Csl and MM) | Total area: ~ 1.4 m² Single unit detect: ~ 0.6 x 0.6 m² | Max.rate:100 Hz/cm ² Spatial res.: <~ 2.5 mm Time res.: ~ 10 ns | Production of large area THGEM of sufficient quality |
| ATLAS MUON UPGRADE CERN LS2 | Hadron Collider (Tracking/Triggering) | Resistive Micromegas | Total area: 1200 m^2 Single unit detect: $(2.2x1.4\text{m}^2) \sim 2-3 \text{ m}^2$ | Max. rate:15 kHz/cm² Spatial res.: <100µm Time res.: ~ 10 ns Rad. Hard.: ~ 0.5C/cm² | Redundant tracking and triggering; Challenging constr. in mechanical precision |
| CMS MUON UPGRADE CERN LS2 | Hadron Collider (Tracking/Triggering) | 3-GEM | Total area: ~ 143 m² Single unit detect: 0.3-0.4m² | Max. rate:10 kHz/cm ² Spatial res.: ~100μm Time res.: ~ 5-7 ns Rad. Hard.: ~ 0.5 C/cm ² | Redundant tracking and triggering |
| ALICE TPC UPGRADE CERN LS2 | Heavy-Ion Physics (Tracking + dE/dx) | 4-GEM / TPC | Total area: ~ 32 m² Single unit detect: up to 0.3m² | Max.rate:100 kHz/cm² Spatial res.: ~300µm Time res.: ~100 ns dE/dx: 11 % | - 50 kHz Pb-Pb rate; - Continues TPC readout - Low IBF and good energy resolution |
| | JINST15 C | 10023 (202 | 20) | Rad. Hard.: 50 mC/cm ² | chargy recording |

CERN – RD51 collaboration): https://rd51-public.web.cern.ch/welc

- ~ 90 RD51 institutes in 25 countries 9 Institutes from the USA
- ✓ Scaling up MPGD detectors, while preserving the typical properties of small prototypes, allowed their use in the LHC upgrades → Many emerged from the R&D studies within the


CERN-RD51 Collaboration

MPGDs not in BRN scope but an important technology

For current status see talk by Maxim Titov @CPAD2021 including application to EIC where all proposed detector concepts include large area **MPGDs**

- Future RD51 activities for advanced **MPGD** concepts includes Generic « Blue-Sky » R&Ds:
 - Resitive materials & architectures:
 - Fast and precise timing;
 - Hybrid detectors (MPGDs + CMOS. optical readout of MPGDs, ...);
 - Novel materials & fabrication techniques (MEMS, nanotechnology, sputtering, novel PCs, 3D printing)
 - arXiv: 1806.09955





CPAD Instrumentation Frontier Workshop 2021

Virtual Event @ Stony Brook University, March 18-22, 2021



What to do with this report and why now #1

During the course of this BRN study the Division of Particles and Fields of the American Physical Society announced the U.S. Particle Physics Community Planning Exercise Snowmass 2021. This will be followed by a new meeting of the Particle Physics Project Prioritization Panel (P5).

We encourage the particle physics community to build on the research plans presented in this BRN study by developing and refining them further and introduce and develop new instrumentation ideas during Snowmass 2021.



Welcome to Snowmass 2021

What to do with this report and why now #2

The ESU states: The community should define a global detector R&D roadmap that should be used to support proposals at the European and national levels."

We support the stance the ESU articulates towards instrumentation.

We encourage the U.S. particle physics community through the Snowmass process to play a role in the ECFA Detector R&D Roadmap which is set in a global context.

After Snowmass CPAD should continue to play a role in developing this international roadmap and can be the vehicle for the realization of the program outlined in this report.



Impact of BRNs & You Part 1

Following the recent microelectronics BRN multiple SC program offices issued a joint DOE National Laboratory Program Announcement focused on multi-disciplinary co-design approaches to basic research that could enable transformative innovation in microelectronic technologies for sensing, communication, and computing.

MICROELECTRONICS CO-DESIGN RESEARCH

DOE NATIONAL LABORATORY PROGRAM ANNOUNCEMENT NUMBER: LAB 21-2491

24 March, 2021



https://science.osti.gov/-/media/grants/pdf/lab-announcements/2021/LAB_21-2491.pdf

Impact of BRNs & You Part 2

Following the recent HEP Instrumentation BRN, HEP issued an Instrumentation Traineeship FOA to support graduate student training in the areas of sensors, front-end electronics and data acquisition, and systems design and engineering; modeled closely after the Accelerator Traineeship FOA.



Department of Energy to Provide \$5 Million to Advance Workforce Development for High Energy Physics Instrumentation

Efforts Will Support Graduate-level Traineeships in Particle Detector Technology

1 April, 2021

https://science.osti.gov/hep/Funding-Opportunities?utm_medium=email&utm_source=govdelivery

The report has already been influential inside the Office of Science. To further increase the impact of the report we are preparing a BRN brochure, will organize a community letter writing campaign (with APS), and work with the FNAL-UEC to carry the report message to Washington.

Acknowledgments

Many contributed at various stages of the Basics Research Needs study that led to this report. We are grateful to those who played roles beyond the report authors. The 66 panel members acknowledge with gratitude:

The 142 other members of the particle physics community who contributed their time and ideas to the BRN study in the months leading up to the workshop. (See back-up for individual names.)

The Report's "readers" gave us critical feedback and provided fact checking during the final stages of preparation. Dan Akerib (SLAC National Laboratory), Myron Campbell (University of Michigan), Andy Lankford (University of California Irvine), Ritchie Patterson (Cornell University), Steve Ritz (University of California Santa Cruz) and Heidi Schellman (University of Oregon).

Our report benefited enormously from professional editing assistance by Tiffani Conner, (Oak Ridge Associated Universities). DOE staff and contractors were always responsive to logistical requests. We especially thank Donna Nevels and Christie Ashton who provided outstandingly professional support at the workshop and contributed importantly to the immensely positive and constructive atmosphere that was highly conducive to productivity.



142 Additional Contributors

Additional Contributors

In addition to the members of the BRN panel many other members of the particle physics community contributed their time and ideas to the BRN study in the months leading up to the workshop. We acknowledge with gratitude:

Energy Frontier: Artur Apresyan (Fermi National Accelerator Laboratory), James Brau (University of Oregon), Martin Breidenbach (SLAC National Accelerator Laboratory), Sarah Eno (University of Maryland), Zhen Liu (University of Maryland), Alessandro Tricoli (Brookhaven National Laboratory);

Neutrinos: Phil Barbeau (Duke University), Flavio Cavanna (Fermilab), André de Gouvêa (Northwestern University), Roni Harnik (Fermilab), Bryce Littlejohn (Illinois Institute of Technology), Georgia Karagiorgi (Columbia University), Josh Klein (University of Pennsylvania), Pedro Machado (Fermilab), Stephen Magill (Argonne National Laboratory), Pedro Ochoa-Ricoux (University of California, Irvine). Gabriel Orebi Gann (University of California, Berkeley, Lawrence Berkeley National Laboratory), Roberto Petti (University of South Carolina), Grayson Rich (University of Chicago), Andres Romero-Wolf (Jet Propulsion Laboratory,

Caltech), Federico Sanchez (University of Geneva), Peter Shanahan (Fermilab), Nick Solomey (Wichita State University), Gensheng Wang (Argonne National Laboratory), Elizabeth Worcester (Brookhaven National Laboratory), Katsuva Yonehara (Fermilab)

Dark Matter: A. Albert (Los Alamos National Laboratory), J. Buckley (University of Washington), G.P. Carosi (Lawrence Livermore National Laboratory), A. Chou (Fermi National Accelerator Laboratory), P. Chu (Los Alamos National Laboratory), E. Dahl (Northwestern University), B. Dingus (Los Alamos National Laboratory), J. Estrada (Fermi National Accelerator Laboratory), C. Galbiati (Princeton University), D. Grant (Michigan State University), C. Hall (University of Maryland), N. Kurinsky (Fermi National Accelerator Laboratory), K. Lesko (Lawrence Berkeley National Laboratory), J. Orrell(Pacific Northwest National Laboratory), M. Pyle (University of California, Berkeley), J. Tieffenberg (Fermi National Accelerator Laboratory), C. Tunnell (Rice University), T.T. Yu (University of Oregon);

Cosmic Acceleration: Elisabeth Krause (University of Arizona), Erik Shirokoff (University of Chicago), Anze Slosar (Brookhaven National Lab)

Explore the Unknown: John M. Doyle (Harvard University), Nicholas R. Hutzler (California Institute of Technology and Harvard University), Takeyasu M. Ito (Los Alamos National Laboratory), Andrew Jayich (University of California Santa Barbara), Edward J. Stephenson (Indiana University), Paula Collins (CERN, European Organization for Nuclear Research), Francesco Forti (Istituto Nazionale di Fisica Nucleare Sezione di Pisa and Universita' di Pisa), Andrey Golutvin (Imperial College London and National University of Science and Technology "MISIS"), Mike Williams (Massachusetts Institute of Technology), Augusto Ceccucci (European Organization for Nuclear Research), Pavel A. Murat (Fermi National Accelerator Laboratory), Gianantonio Pezzullo (Yale University)

142 Additional Contributors

Additional Contributors

In addition to the members of the BRN panel many other members of the particle physics community contributed their time and ideas to the BRN study in the months leading up to the workshop. We acknowledge with gratitude:

Calorimetry: M. Aleska (CERN, The European Organization for Nuclear Research), A. Apresyan (Fermi National Accelerator Laboratory), F. Bedeschi (Istituto Nazionale di Fisica Nucleare, Sezione di Pisa), J. Brau (The University of Oregon), M. Breidenbach (SLAC National Accelerator Laboratory), S. Derenzo (University of California, Berkeley), V. Fadeyev (University of California, Santa Cruz), R. Ferari (Istituto Nazionale di Fisica Nucleare, Sezione di Pavia and Università di Pavia), G. Gaudio (Istituto Nazionale di Fisica Nucleare, Sezione di Pavia and Università di Pavia), P. Giacomelli (Istituto Nazionale di Fisica Nucleare, Sezione di Bologna), D. Hitlin (Caltech), P. Lecoq (CERN, The European Organization for Nuclear Research), H. Ma (Brookhaven National Laboratory), M. Mannelli (CERN, The European Organization for Nuclear Research), C. Melcher (The University of Tennessee, Knoxville), A. Para (Fermi National Accelerator Laboratory), L. Pezotti (Istituto Nazionale di Fisica Nucleare, Sezione di Pavia and Università di Pavia), A. Tricoli (Brookhaven National Laboratory), C. Woody (Brookhaven National Laboratory).

Noble Liquids: R. Acciarri (Fermi National Accelerator Laboratory), N. Bostan (University of Iowa), Flavio Cavanna (Fermi National Accelerator Laboratory), Serge Charlebois (University of Sherbrooke), Eric Dahl (Northwestern University), C. O. Escobar (Fermi National Accelerator Laboratory), A. Fava (Fermi National Accelerator Laboratory), C. Galbiati (Princeton University), C. Hall (University of Maryland), B.J.P. Jones (University of Texas at Arlington), S. Kravitz (Lawrence Berkeley National Laboratory), R. Lang (Purdue University), K. Lesko (Lawrence Berkeley National Lab), B. Littlejohn (Illinois Institute of Technology), D. McKinsey (University of California, Berkeley / Lawrence Berkeley National Lab), C. S. Montanari (Fermi National Accelerator Laboratory), H. Nelson (University California, Santa Barbara), K. Ni (University of California, San Diego), X. Qian (Brookhaven National Laboratory), P. C. Rowson (SLAC National Accelerator Laboratory), M. Szydagis (University at Albany – State University of New York), C. Tunnell (Rice University), G. Visser (Indiana University), C. Zhang (Brookhaven National Laboratory)

Photodetectors: S. Cho (Stanford University), A. Elagin (University of Chicago), R. Hirosky (University of Virginia), D. Hitlin (Caltech), S. Korpar (University of Maribor and JSI, Ljubljana), Sae Woo Nam (NIST), C. Rockosi (University of California, Santa Cruz), J. Xie (Argonne National Lab)

Quantum Sensors: Derek Jackson-Kimball (California State University - East Bay), Cindy Regal (JILA, University of Colorado - Boulder), Kater Murch (Washington University - Saint Louis), Swati Singh (University of Delaware), David DeMille (Yale University), Yoni Kahn (University of Illinois at Urbana-Champaign), Tim Kovachy (Northwestern University), Nick Hutlzer (Caltech), Hartmut Haffner (University of California-Berkeley).

142 Additional Contributors

Additional Contributors

In addition to the members of the BRN panel many other members of the particle physics community contributed their time and ideas to the BRN study in the months leading up to the workshop. We acknowledge with gratitude:

Readout & ASICs: T. Affolder (University of California, Santa Cruz), G. Deptuch (Brookhaven National Lab), G. Drake (Fermi National Accelerator Laboratory), C. Grace (Lawrence Berkeley National Laboratory), W. Hansford (IMEC USA), J. Cressler (Georgia Institute of Technology), P. Merkel (Fermi

National Accelerator Laboratory), P. O'connor (Brookhaven National Laboratory), A. Suzuki (Lawrence Berkeley National Laboratory)

Solid State: R. Brenner (University of Uppsala), V.Fadeyev (University of California, Santa Cruz), T.Heim (Lawrence Berkeley National Lab), S.C. Hsu (University of Washington), K. Krizka (Lawrence Berkeley National Laboratory), J. Metcalfe (Argonne National Laboratory), S. Seidel (University of New Mexico), D.Stuart (University of California, Santa Barbara), C. da Via (University of Mancester, SBU)

TDAQ: K. Chen (Brookhaven National Laboratory), K. Ecklund (Rice University), J. Eisch (Iowa State University), P. Harris (Massachusetts Institute of Technology), M. Liu (Fermi National Accelerator Laboratory), I. Ojalvo (Princeton University), A. Slosar (Brookhaven National Laboratory), N. Tran (Fermi National Accelerator Laboratory), M. Wetstein (Iowa State University), M. Williams (Massachusetts Institute of Technology), P. Wittich (Cornell University).