




LHC⁻¹ = ILC?

Part I

Carola F. Berger

Stanford Linear Accelerator Center

BNL 2007: New Horizons at Colliders – May 30th, 2007




[The LHC Inverse
Problem –
Summary](#)

[SUSY at the ILC](#)

[Sample Results –
Slepton Analyses](#)

[Summary and
Outlook](#)

$\text{LHC}^{-1} = \text{ILC} ?$



The LHC Inverse
Problem –
Summary


SUSY at the ILC

Sample Results –
Slepton Analyses

Summary and
Outlook

$\text{LHC}^{-1} = \text{ILC} ?$

Yes, of course!



The LHC Inverse
Problem –
Summary

SUSY at the ILC


Sample Results –
Slepton Analyses

Summary and
Outlook

$\text{LHC}^{-1} = \text{ILC} ?$

~~Yes, of course!~~
but


To appear, arXiv/0706.xxxx [hep-ph], featuring CFB, James Gainer,
JoAnne L. Hewett, Ben Lillie, Thomas G. Rizzo



Outline

- **Summary of results of “The LHC Inverse Problem”, how to map from observables to a Lagrangian**
- **SUSY at the ILC**
Standard Model background and analyses for sleptons
- **Results for sleptons (preliminary!)**
- **Summary and outlook**

Part II – charginos \Rightarrow Tom Rizzo’s talk



Introduction

The LHC Inverse
Problem –
Summary

- Introduction
- A Typical Degenerate Pair – Example 1
- A Typical Degenerate Pair – Example 2
- A Typical Degenerate Pair – Example 3

SUSY at the ILC

Sample Results –
Slepton Analyses

Summary and
Outlook


What's in a name?


Theorists: parameters \Rightarrow observables

Experimentalists: observables \Rightarrow parameters

N. Arkani-Hamed, G. L. Kane, J. Thaler, L.-T. Wang, *Supersymmetry and the LHC inverse problem*, JHEP 0608, 070 (2006) [hep-ph/0512190]

Question: parameters \Leftrightarrow observables – a one-to-one map?





Introduction

The LHC Inverse
Problem –
Summary

- Introduction
- A Typical Degenerate Pair – Example 1
- A Typical Degenerate Pair – Example 2
- A Typical Degenerate Pair – Example 3

SUSY at the ILC

Sample Results –
Slepton Analyses

Summary and
Outlook


What's in a name?


Theorists: parameters \Rightarrow observables

Experimentalists: observables \Rightarrow parameters

N. Arkani-Hamed, G. L. Kane, J. Thaler, L.-T. Wang, *Supersymmetry and the LHC inverse problem*, JHEP 0608, 070 (2006) [hep-ph/0512190]

Question: parameters \Leftrightarrow observables – a one-to-one map?





Introduction

The LHC Inverse
Problem –
Summary

- Introduction
- A Typical Degenerate Pair – Example 1
- A Typical Degenerate Pair – Example 2
- A Typical Degenerate Pair – Example 3

SUSY at the ILC

Sample Results –
Slepton Analyses

Summary and
Outlook


What's in a name?


Theorists: parameters \Rightarrow observables

Experimentalists: observables \Rightarrow parameters

N. Arkani-Hamed, G. L. Kane, J. Thaler, L.-T. Wang, *Supersymmetry and the LHC inverse problem*, JHEP 0608, 070 (2006) [hep-ph/0512190]

Question: parameters \Leftrightarrow observables – a one-to-one map?





Introduction contd.

hep-ph/0512190: scan over (restricted) MSSM parameter space, simulation of different “models”.

Existence proof: there are distinct MSSM models that are indistinguishable at LHC with standard set of observables even without SM background.

The LHC can only measure mass differences accurately, initial state (= c.m. energy per collision) not well defined.

In addition, multiparton final states dominate. In general, “messy” final states \Rightarrow soft particles not visible.

The LHC Inverse Problem – Summary

● Introduction

● A Typical Degenerate Pair – Example 1


● A Typical Degenerate Pair – Example 2

● A Typical Degenerate Pair – Example 3

SUSY at the ILC

Sample Results – Slepton Analyses

Summary and Outlook



Introduction contd.

hep-ph/0512190: scan over (restricted) MSSM parameter space, simulation of different “models”.

Existence proof: there are distinct MSSM models that are indistinguishable at LHC with standard set of observables even without SM background.

The LHC can only measure mass differences accurately, initial state (= c.m. energy per collision) not well defined.

In addition, multiparton final states dominate. In general, “messy” final states \Rightarrow soft particles not visible.

Question: given the list of MSSM “models” that turned out to be indistinguishable at the LHC in hep-ph/0512190, can the ILC distinguish them, using which observables?

The LHC Inverse Problem – Summary

● Introduction

● A Typical Degenerate Pair – Example 1


● A Typical Degenerate Pair – Example 2

● A Typical Degenerate Pair – Example 3

SUSY at the ILC

Sample Results – Slepton Analyses

Summary and Outlook



A Typical Degenerate Pair – Example 1



The LHC Inverse Problem – Summary


- Introduction
- A Typical Degenerate Pair – Example 1
- A Typical Degenerate Pair – Example 2
- A Typical Degenerate Pair – Example 3

SUSY at the ILC


Sample Results – Slepton Analyses

Summary and Outlook





A Typical Degenerate Pair – Example 1



“Flipper”: mass of LSP fixed, but identity changes between mostly bino/wino and mostly Higgsino


The LHC Inverse Problem – Summary

- Introduction
- A Typical Degenerate Pair – Example 1
- A Typical Degenerate Pair – Example 2
- A Typical Degenerate Pair – Example 3

SUSY at the ILC

Sample Results – Slepton Analyses

Summary and Outlook



A Typical Degenerate Pair – Example 2



The LHC Inverse Problem – Summary


- Introduction
- A Typical Degenerate Pair – Example 1
- A Typical Degenerate Pair – Example 2
- A Typical Degenerate Pair – Example 3

SUSY at the ILC


Sample Results – Slepton Analyses

Summary and Outlook





A Typical Degenerate Pair – Example 2



“Slider”: mass differences roughly the same, but shifted by about 100 GeV.


The LHC Inverse Problem – Summary

- Introduction
- A Typical Degenerate Pair – Example 1
- A Typical Degenerate Pair – Example 2
- A Typical Degenerate Pair – Example 3

SUSY at the ILC

Sample Results – Slepton Analyses

Summary and Outlook



A Typical Degenerate Pair – Example 3



The LHC Inverse Problem – Summary


- Introduction
- A Typical Degenerate Pair – Example 1
- A Typical Degenerate Pair – Example 2
- A Typical Degenerate Pair – Example 3

SUSY at the ILC

Sample Results – Slepton Analyses

Summary and Outlook





A Typical Degenerate Pair – Example 3



“Squeezer”: mass differences between electroweakinos too similar – decay products too soft


The LHC Inverse Problem – Summary

- Introduction
- A Typical Degenerate Pair – Example 1
- A Typical Degenerate Pair – Example 2
- A Typical Degenerate Pair – Example 3

SUSY at the ILC

Sample Results – Slepton Analyses

Summary and Outlook



Our Project – Overview

- Simulate signal events with Pythia, and feed in appropriate beamspectrum generated via Whizard/GuineaPig, including ISR, beamstrahlung, beamspread
- Add SM background (1016 different processes), produced by Tim Barklow, stored on SLAC tape
- Pipe through detector simulation
Java-based detector simulation, code developed by SLAC ILC group: `org.lcsim`, SiD detector concept
- Analyze 500 fb^{-1} of “data” with appropriate cuts
Several iterations necessary to find best cuts


The LHC Inverse Problem – Summary

SUSY at the ILC

- Our Project – Overview
- Characteristics of Signal Models
- Analyses
- Cuts

Sample Results – Slepton Analyses

Summary and Outlook



Characteristics of Signal Models

The LHC Inverse
Problem –
Summary


SUSY at the ILC

- Our Project – Overview
- Characteristics of Signal Models
- Analyses
- Cuts


Sample Results –
Slepton Analyses

Summary and
Outlook

Sparticle counts



Accessible at 500 GeV, 1 TeV c.m. energy



Analyses

The LHC Inverse
Problem –
Summary


SUSY at the ILC

- Our Project – Overview
- Characteristics of Signal Models
- Analyses
- Cuts

Sample Results –
Slepton Analyses

Summary and
Outlook

- **Selectrons – electron-positron pair plus LSPs (missing E_T)**
- **Smuons – muon-pair plus LSPs (missing E_T)**
- **Staus – tau-pair plus LSPs (missing E_T) - tau ID!**
- **Charginos – several sub-scenarios depending on $\Delta m_{\tilde{\chi}} = m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0}$**
 - ◆ $\Delta m_{\tilde{\chi}} > 1 \text{ GeV}$ – jets and/or μ s plus LSPs
 - ◆ $m_\pi \leq \Delta m_{\tilde{\chi}} < 1 \text{ GeV}$ – radiative, tag on high-E photon
 - ◆ $\Delta m_{\tilde{\chi}} < m_\pi$ – massive, stable, charged track
- **Radiative Neutralinos**



Cuts

$\gamma\gamma$ and $e\gamma$ SM background nontrivial!
Need good particle ID and tracking down to low angles.

Cuts adapted and expanded from

- U. Nauenberg, *et al.*, Colorado SUSY group,
<http://hep-www.colorado.edu/SUSY/susynlc.html>
- H. U. Martyn, arXiv:hep-ph/0408226
- P. Bambade, M. Berggren, F. Richard and Z. Zhang, arXiv:hep-ph/0406010
- G. Abbiendi *et al.* [OPAL Collaboration], Eur. Phys. J. C 35, 1 (2004)
[arXiv:hep-ex/0401026]
- C. H. Chen, M. Drees and J. F. Gunion, Phys. Rev. Lett. 76, 2002 (1996)
[arXiv:hep-ph/9512230]; J. F. Gunion and S. Mrenna, Phys. Rev. D 64, 075002 (2001)
[arXiv:hep-ph/0103167]
- H. K. Dreiner, O. Kittel and U. Langenfeld, arXiv:hep-ph/0703009


The LHC Inverse
Problem –
Summary

SUSY at the ILC

- Our Project – Overview
- Characteristics of Signal Models
- Analyses
- Cuts

Sample Results –
Slepton Analyses

Summary and
Outlook



SM Background – Selectrons


The LHC Inverse
Problem –
Summary

SUSY at the ILC


Sample Results –
Slepton Analyses

- SM Background – Selectrons
- SM Background contd. – p_T – Selectrons
- Importance of Forward Coverage
- Signal – Selectrons
- Signal + Background – Selectrons
- Sample Model Comparison

Summary and
Outlook



Bigger than “standard” background



SM Background contd. – p_T – Selectrons

Comparison Whizard \Leftrightarrow PYTHIA:

The LHC Inverse
Problem –
Summary


SUSY at the ILC

Sample Results –
Slepton Analyses


- SM Background – Selectrons
- SM Background contd. – p_T – Selectrons
- Importance of Forward Coverage
- Signal – Selectrons
- Signal + Background – Selectrons
- Sample Model Comparison

Summary and
Outlook

$$e_{pol}^- = 0 \quad \sqrt{s} = 500 \text{ GeV} \quad 250 \text{ fb}^{-1}$$




new Whizard EPA



9

Thanks to Tim Barklow



SM Background contd. – p_T – Selectrons

The LHC Inverse
Problem –
Summary

SUSY at the ILC


Sample Results –
Slepton Analyses

- SM Background – Selectrons
- SM Background contd. – p_T – Selectrons
- Importance of Forward Coverage
- Signal – Selectrons
- Signal + Background – Selectrons
- Sample Model Comparison

Summary and
Outlook


Comparison Whizard \Leftrightarrow PYTHIA:

$$e_{pol}^- = 0 \quad \sqrt{s} = 500 \text{ GeV} \quad 250 \text{ fb}^{-1}$$



10

Thanks to Tim Barklow



Importance of Forward Coverage


The LHC Inverse Problem – Summary

SUSY at the ILC


Sample Results – Slepton Analyses

- SM Background – Selectrons
 - SM Background contd. – p_T – Selectrons
 - Importance of Forward Coverage
 - Signal – Selectrons
 - Signal + Background – Selectrons
 - Sample Model Comparison

Summary and Outlook



$$20 \text{ mrad} \xrightarrow{-9\%} 5 \text{ mrad} \xrightarrow{-3\%} 3.5 \text{ mrad}$$



Signal – Selectrons


The LHC Inverse
Problem –
Summary


SUSY at the ILC

Sample Results –
Slepton Analyses

- SM Background – Selectrons
- SM Background contd. – p_T – Selectrons
- Importance of Forward Coverage
- Signal – Selectrons
- Signal + Background – Selectrons
- Sample Model Comparison

Summary and
Outlook





Signal – Selectrons


The LHC Inverse Problem – Summary


SUSY at the ILC

Sample Results – Slepton Analyses

- SM Background – Selectrons
 - SM Background contd. – p_T – Selectrons
 - Importance of Forward Coverage
 - Signal – Selectrons
 - Signal + Background – Selectrons
 - Sample Model Comparison

Summary and Outlook





Signal – Selectrons


The LHC Inverse
Problem –
Summary


SUSY at the ILC

Sample Results –
Slepton Analyses

- SM Background – Selectrons
- SM Background contd. – p_T – Selectrons
- Importance of Forward Coverage
- Signal – Selectrons
- Signal + Background – Selectrons
- Sample Model Comparison

Summary and
Outlook





Signal – Selectrons


The LHC Inverse Problem – Summary


SUSY at the ILC

Sample Results – Slepton Analyses

- SM Background – Selectrons
- SM Background contd. – p_T – Selectrons
- Importance of Forward Coverage
- Signal – Selectrons
- Signal + Background – Selectrons
- Sample Model Comparison

Summary and Outlook





Signal + Background – Selectrons


The LHC Inverse
Problem –
Summary


SUSY at the ILC

Sample Results –
Slepton Analyses

- SM Background – Selectrons
- SM Background contd. – p_T – Selectrons
- Importance of Forward Coverage
- Signal – Selectrons
- Signal + Background – Selectrons
- Sample Model Comparison

Summary and
Outlook





Signal + Background – Selectrons


The LHC Inverse
Problem –
Summary


SUSY at the ILC

Sample Results –
Slepton Analyses

- SM Background – Selectrons
- SM Background contd. – p_T – Selectrons
- Importance of Forward Coverage
- Signal – Selectrons
- Signal + Background – Selectrons
- Sample Model Comparison

Summary and
Outlook





Signal + Background – Selectrons


The LHC Inverse
Problem –
Summary


SUSY at the ILC

Sample Results –
Slepton Analyses

- SM Background – Selectrons
- SM Background contd. – p_T – Selectrons
- Importance of Forward Coverage
- Signal – Selectrons
- Signal + Background – Selectrons
- Sample Model Comparison

Summary and
Outlook





Sample Model Comparison

Particle masses:

particle	Model 2631	Model 16195
$\tilde{e}_L, \tilde{\mu}_L$	190.2	459.1
$\tilde{e}_R, \tilde{\mu}_R$	759.8	708.7
$\tilde{\tau}_1$	177.4	528.9
$\tilde{\tau}_2$	842.2	552.5
$\tilde{\chi}_1^0$	102.7	151.3
$\tilde{\chi}_1^\pm$	759.9	767.6
\tilde{q}_L	750	655
\tilde{d}_R, \tilde{s}_R	630.5	984.3
\tilde{u}_R, \tilde{c}_R	700.4	787.1
\tilde{g}	1003.9	1014.8


The LHC Inverse Problem – Summary

SUSY at the ILC

Sample Results – Slepton Analyses

- SM Background – Selectrons
- SM Background contd. – p_T – Selectrons
- Importance of Forward Coverage
- Signal – Selectrons
- Signal + Background – Selectrons
- Sample Model Comparison

Summary and Outlook



Sample Model Comparison contd.


The LHC Inverse Problem – Summary


SUSY at the ILC

Sample Results – Slepton Analyses

- SM Background – Selectrons
- SM Background contd. – p_T – Selectrons
- Importance of Forward Coverage
- Signal – Selectrons
- Signal + Background – Selectrons
- Sample Model Comparison

Summary and Outlook





Summary


- First analysis of “random” points in SUSY parameter space looking at a variety of observables with full SM background.
- First “user” analysis with `org.lcsim` and SiD detector concept.
- **Sleptons:** ILC can in principle distinguish between models that are degenerate at the LHC, by using polarized beams and due to accurate mass measurements.
- In practise cuts need to be optimized, and detector coverage as close to the beam direction as possible is very desirable.
Low-angle tracking possibly very important
- There are regions in SUSY parameter space that are challenging due to low number of signal events and SM background.

The LHC Inverse Problem – Summary

SUSY at the ILC

Sample Results – Slepton Analyses

Summary and Outlook
● Summary
● Outlook



Outlook

- Tom's talk: **special feature on charginos**
- Study 1 TeV case
- Merge with information from cosmology and precision flavor studies
- **Work in progress!**
Stay tuned for the final results

The LHC Inverse
Problem –
Summary

SUSY at the ILC

Sample Results –
Slepton Analyses

Summary and
Outlook

- Summary
- Outlook