Charged particles scattering in a finite volume

Silas Beane

BNL 2/5/2015

Outline

- Motivation
- Effective range theory / with Coulomb
- Review of A_1^+ QC in FV
- Coulomb in a FV w/ M.J.Savage, arXiv:1407.4846
- Conclusions

✦ Motivation

✓ Lattice QCD calculates scattering parameters



✓ $\alpha = 0$ phase shift prediction



✓ Can one get proton-proton scattering *ab initio* ?



Di-neutron binding energy: world data



Di-neutron binding energy: world data



✦ Effective Range Theory (s-wave)

(Bethe, 1949)

✓ R is the range of the interaction

✓ $p \ll R^{-1}$ gives expansion of interaction in local operators:

$C(E^*) =$  $= C_0 + C_2 M T^* + \ldots$

center-of-mass energy

$$E^* = 2M + \frac{p^2}{M} = 2M + T^*$$

✦ Effective Range Theory (s-wave)

(Bethe, 1949)

✓ R is the range of the interaction

✓ $p \ll R^{-1}$ gives expansion of interaction in local operators:

$C(E^*) =$  $= C_0 + C_2 M T^* + \ldots$

center-of-mass energy

$$E^* = 2M + \frac{p^2}{M} = 2M + T^*$$

✓ In practice, remove momentum dependence in EFT:

$$-\hat{\theta}\, \psi^T (\overleftarrow{\nabla} - \overrightarrow{\nabla})^2 \psi = 4M\, \hat{\theta} \left[i\partial_0 + \frac{\nabla^2}{4M} \right] \psi^T \psi \equiv 4M\, \hat{\theta}\, \mathcal{O}_{E^*} \psi^T \psi$$

✓ Scattering amplitude: interaction summed to all orders:

$T_S =$ 

$$T_S = \frac{C(E^*)}{1 - C(E^*) J_0^\infty(E^*)} = -\frac{4\pi}{M} \frac{1}{p \cot\delta - ip}$$

✓ Matching achieved through renormalization ($\overline{MS}$):

$J_0^\infty(E^*) =$  $= M \int \frac{d^3q}{(2\pi)^3} \frac{1}{p^2 - q^2 + i\epsilon}$ Green function

$$p \cot\delta = -\frac{1}{a} + \frac{1}{2} r_0 p^2 + \ldots$$

✓ Analytic structure of the scattering amplitude



✓ Analytic structure of the scattering amplitude



✓ Analytic structure of the scattering amplitude



✦ Effective Range Theory with Coulomb

✓ Coulomb is enhanced in the infrared: (Bethe, 1949)

$$p \sim \alpha M \sim a_{\text{bohr}}^{-1}$$



$$\frac{\alpha}{(\mathbf{p}-\mathbf{p}')^2} \sim \mathcal{O}(\alpha^{-1})$$



$$\int d^3\mathbf{k}dk_0 \frac{1}{\left(\frac{E}{2}+k_0-\frac{\mathbf{k}^2}{2M}+i\epsilon\right)} \frac{1}{\left(\frac{E}{2}-k_0-\frac{\mathbf{k}^2}{2M}+i\epsilon\right)} \frac{\alpha}{(\mathbf{k}-\mathbf{p})^2} \frac{\alpha}{(\mathbf{k}-\mathbf{p}')^2} \sim \mathcal{O}(\alpha^{-1})$$

Coulomb is non-perturbative

✦ Effective Range Theory with Coulomb

✓ Coulomb is enhanced in the infrared: (Bethe, 1949)

$$p \sim \alpha M \sim a_{\rm bohr}^{-1}$$



$$\frac{\alpha}{(\mathbf{p}-\mathbf{p}')^2} \sim \mathcal{O}(\alpha^{-1})$$



$$\int d^3\mathbf{k}dk_0 \frac{1}{\left(\frac{E}{2}+k_0-\frac{\mathbf{k}^2}{2M}+i\epsilon\right)} \frac{1}{\left(\frac{E}{2}-k_0-\frac{\mathbf{k}^2}{2M}+i\epsilon\right)} \frac{\alpha}{(\mathbf{k}-\mathbf{p})^2} \frac{\alpha}{(\mathbf{k}-\mathbf{p}')^2} \sim \mathcal{O}(\alpha^{-1})$$

Coulomb is non-perturbative

✓ Radiation is suppressed



$$\sim \mathcal{O}(1)$$

✦ Effective Range Theory with Coulomb

✓ Coulomb ladders treated to all orders:



$$\sigma_0 = \arg\Gamma(1+i\eta)$$ is Coulomb phase shift

$$\boxed{T_{SC} = C^2_{\eta(p)} \frac{C(E^*)e^{i2\sigma_0}}{1 - C(E^*)J_0^\infty(E^*)} = -\frac{4\pi}{M}\frac{e^{2i\sigma_0}}{p\cot\delta - ip}}$$

$$C^2_{\eta(p)} = \frac{2\pi\eta(p)}{e^{2\pi\eta(p)} - 1} \qquad \eta \equiv \frac{\alpha M}{2p} \sim \frac{\alpha}{v} \sim 1$$

$$J_0^\infty(E^*) \;=\; M\int \frac{d^3q}{(2\pi)^3}\frac{C^2_{\eta(q)}}{p^2-q^2+i\epsilon}$$

QED "dressed" Green function

✓ Match to effective range parameters with $\overline{MS}$:

$$C^2_{\eta(p)}\, p\cot\delta + \alpha M h(\eta) \;=\; -\frac{1}{a_C} \;+\; \frac{1}{2}r_0 p^2 \;+\; \ldots$$

$$\downarrow$$

$$\ln\left(\frac{M\alpha}{p}\right) \;+\; \mathcal{O}(\alpha)$$

Removes non-analytic piece!

$$(pp: \quad a_c \;=\; -7.82 \text{ fm})$$

✓ Analytic structure of the scattering amplitude



✓ Analytic structure of the scattering amplitude



✓ Analytic structure of the scattering amplitude



✦ Review of A_1^+ QC in FV

$b = 0,\ T = \infty,\ s - wave$

(Lüscher, 1990)



$$R < L/2$$

✓ Two-particle correlation function in FV

Source

Sink

$$S^\dagger \left[J_0^L(E^*) \; + \; C^L(E^*) \left(J_0^L(E^*)\right)^2 \; + \; C^L(E^*)^2 \left(J_0^L(E^*)\right)^3 \; + \; \ldots \right] S$$

$$= \; S^\dagger \frac{1}{1/J_0^L(E^*) - C^L(E^*)} S$$

Singularities of the correlation function

$$\boxed{\frac{1}{C^L(E^*)} \; = \; J_0^L(E^*)}$$



$$J_0^L(E^*) \; = \; -\frac{M}{4\pi^2 L} \sum_{\mathbf{n}}^{\Lambda_n} \frac{1}{|\mathbf{n}|^2 - \tilde{p}^2}$$

UV same as the continuum

✓ Quantization condition:

$$\boxed{p \cot \delta \;=\; \frac{1}{\pi L} \mathcal{S}(\tilde{p})} \;+\; \mathcal{O}\left(e^{-m_\pi L}\right)$$

Valid in QFT up to inelastic threshold



$$\mathcal{S}(x) \;\equiv\; \sum_{\mathbf{n}}^{\Lambda_n} \frac{1}{|\mathbf{n}|^2 - x^2} \;-\; 4\pi\Lambda_n$$

✓ Quantization condition in practice:



✦ A_1^+ QC in FV with Coulomb



A prior there are many issues at FV…

- QED ill defined with PBCs — omit zero mode (ZD Talk)
- Increasingly complex integer sums .. many infrared problems
- Inelastic thresholds suppressed relative to Coulomb?
- Kinematics altered by power law effects
- Lack of universality

✓ Absence of zero mode: gap in momentum operator

$$p_{min} = \frac{2\pi}{L}$$

(ZD Talk)

$$L \to \infty$$



✓ Absence of zero mode: gap in momentum operator

$$p_{min} = \frac{2\pi}{L}$$

(ZD Talk)

$$L \to \infty$$



Gap is a good thing: $\eta \equiv \frac{\alpha M}{2p} \sim \alpha M L$

$\eta \ll 1$ if $1 \ll ML \ll 1/\alpha$

Coulomb is perturbative!

✓ Zero mode removed



✓ Kinematics altered by power-law corrections:

$$E^* = 2M^L + T^{*L}$$

NRQED

$$p \cot \delta \quad \to -\frac{1}{a'_C} + \frac{1}{2} r'_0 M T^{*L} + r'_1 M^2 (T^{*L})^2 \; + \; \ldots$$

$$\frac{1}{a'_C} = \frac{1}{a_C} - \frac{\alpha \, r_0 M \, \mathcal{I}}{2\pi L} \; + \; \mathcal{O}(\alpha^2; \alpha/L^2)$$

$$r'_0 = r_0 + \frac{4 \, \alpha \, r_1 M \, \mathcal{I}}{\pi L} \; + \; \mathcal{O}(\alpha^2; \alpha/L^2)$$

Kinematically shifted effective range parameters

$$(\mathcal{I} = \pi c_1)$$

Source

Sink

$$= S^\dagger \frac{1}{1/J_0^L(E^*) - C^L(E^*)} S \quad \longrightarrow \quad \boxed{\frac{1}{C^L(E^*)} = J_0^L(E^*)}$$

$$J_0^\infty(E^*) =$$



UV same as the continuum

$$- M \int \frac{d^3q}{(2\pi)^3} \frac{1}{q^2 - p^2} + 4\pi\alpha M^2 \int \frac{d^3q}{(2\pi)^3} \int \frac{d^3k}{(2\pi)^3} \frac{1}{q^2 - p^2} \frac{1}{k^2 - p^2} \frac{1}{|\mathbf{q} - \mathbf{k}|^2}$$

FV with PBCs

$$- \frac{M}{4\pi^2 L} \sum_{\mathbf{n}}^{\Lambda_n} \frac{1}{|\mathbf{n}|^2 - \tilde{p}^2} + \frac{\alpha M^2}{16\pi^5} \sum_{\mathbf{n}}^{\Lambda_n} \sum_{\mathbf{m} \neq \mathbf{n}}^{\infty} \frac{1}{|\mathbf{n}|^2 - \tilde{p}^2} \frac{1}{|\mathbf{m}|^2 - \tilde{p}^2} \frac{1}{|\mathbf{n} - \mathbf{m}|^2} \qquad \tilde{p} = Lp/2\pi$$

✓ Radiation *not* suppressed by α



$$\sim \left(\sqrt{M/L}\right)^n$$

Infrared suppressed!



$$\delta C^{(FV)}(E^*) = -\alpha \left(\frac{4a_C^2 r_0}{L} \mathcal{I} + \ldots \right)$$

$$J_0^L \to J_0^L + (J_0^L)^2 \, \delta C^{(FV)}$$

$$(\mathcal{I} = \pi c_1)$$

Infrared suppressed but let's keep it..

✓ Quantization condition:

$$\boxed{-\frac{1}{a'_C} + \frac{1}{2} r'_0 p^2 + \ldots = \frac{1}{\pi L} \mathcal{S}^C(\tilde{p}) + \alpha M \left[\ln\left(\frac{4\pi}{\alpha M \mathrm{L}} \right) - \gamma_E \right]} + \ldots$$

Valid up to inelastic threshold!

$$\mathcal{S}^C(x) \equiv \mathcal{S}(x) - \frac{\alpha M L}{4\pi^3} \mathcal{S}_2(x) + \frac{\alpha\, M a_C^2 r_0}{\pi^2 L^2} \mathcal{I}\left[\mathcal{S}(x)\right]^2$$

$$\mathcal{S}_2(x) \equiv \sum_{\mathbf{n}}^{\Lambda_n} \sum_{\mathbf{m} \neq \mathbf{n}}^{\infty} \frac{1}{|\mathbf{n}|^2 - x^2} \frac{1}{|\mathbf{m}|^2 - x^2} \frac{1}{|\mathbf{n} - \mathbf{m}|^2} - 4\pi^4 \ln \Lambda_n$$

✓ Note the breakdown of universality

+ ... contains infrared suppressed, subleading QED, and 1/M effects

✓ Define a scale-dependent scattering length:

$$\frac{1}{a(\mu)} \equiv \frac{4\pi}{MC(0;\mu)} = \frac{1}{a_C} + \alpha M \left[\ln\left(\frac{2\mu}{\alpha M}\right) - \gamma_E \right]$$

$\overline{MS}_{FV}$

exhibits the impossibility of separating QCD and QED contributions

$$p \cot \bar{\delta}(p,\mu) \equiv -\frac{1}{a(\mu)} + \frac{1}{2} r_0 p^2 + \ldots$$

✓ Alternate form of QC:

$$p \cot \bar{\delta}'(p, 2\pi/L) \equiv -\frac{1}{a'(2\pi/L)} + \frac{1}{2} r_0' p^2 + \ldots = \frac{1}{\pi L} \mathcal{S}^C(\tilde{p})$$

leading QED effects from distance scales $> L$ are removed

Approximate formulas

✓ Ground state energy level:

$$\Delta E_0^C = \frac{4\pi\, a_C}{M\, L^3}\left\{1 - \left(\frac{a_C}{\pi\, L}\right)\mathcal{I} + \left(\frac{a_C}{\pi\, L}\right)^2 \left[\mathcal{I}^2 - \mathcal{J}\right] + \ldots\right\}$$

$$- \frac{2\alpha\, a_C}{L^2\pi^2}\left\{\mathcal{J} + \left(\frac{a_C}{\pi\, L}\right)\left[\mathcal{K} - \mathcal{I}\mathcal{J} - \tilde{\mathcal{R}}/2\right]\right.$$

$$+ \left(\frac{a_C}{\pi\, L}\right)^2 \left[\tilde{\mathcal{R}}\mathcal{I} + \mathcal{I}^2\mathcal{J} - 2\mathcal{J}^2 - 2\mathcal{I}\mathcal{K} + \mathcal{L} - \mathcal{R}_{24}\right]$$

$$\left. + \frac{a_C r_0 \pi^2}{L^2}\mathcal{I} + \ldots\right\}$$

geometric constants: $\mathcal{I}, \mathcal{J}, \mathcal{K}, \mathcal{L}, \mathcal{R}, \mathcal{R}_{24}$

$$\tilde{\mathcal{R}} \equiv \mathcal{R} - 4\pi^4\left[\ln\left(\frac{4\pi}{\alpha M L}\right) - \gamma_E\right]$$

✓ First excited energy level:

$$\Delta E_1^C = \frac{4\pi^2}{ML^2} - \frac{12\tan\overline{\delta}'}{ML^2}\left(1 + c_1'\tan\overline{\delta}' + c_2'\tan^2\overline{\delta}' + \ldots\right)$$
$$+ \frac{9\alpha}{4\pi \mathrm{L}}\left(1 + c_{1\alpha}'\tan\overline{\delta}' + \left(c_{2\alpha}' + \frac{8}{3}\log(\alpha M L)\right)\tan^2\overline{\delta}' + \ldots\right)$$

geometric constants: $c_1', c_2', c_{1\alpha}', c_{2\alpha}'$

$$\frac{9\alpha}{4\pi \mathrm{L}} \xrightarrow[\mathrm{L} = 10\ \mathrm{fm}]{} \sim 100\ \mathrm{keV}$$

Coulomb effects are very small!

Bound state

✓ Bound state QC with binding momentum κ :

$$-\frac{1}{a_C} - \frac{1}{2} r_0 \kappa^2 = -\kappa - \alpha M \left(\gamma_E + \log\left(\frac{\alpha M}{4\kappa}\right) \right) - \frac{\alpha M}{2\pi\kappa L}(1-\kappa r_0)\mathcal{I}$$

$$\boxed{B^C = \frac{\kappa_0^2}{M} - \frac{2\alpha\kappa_0}{1-\kappa_0 r_0}\left[\gamma_E + \log\left(\frac{\alpha M}{4\kappa_0}\right)\right] - \frac{\alpha}{\pi L}\mathcal{I} + \ldots}$$

Bound state

✓ Bound state QC with binding momentum κ :

$$-\frac{1}{a_C} - \frac{1}{2} r_0 \kappa^2 \;=\; -\kappa - \alpha M \left(\gamma_E + \log\left(\frac{\alpha M}{4\kappa}\right) \right) \;-\; \frac{\alpha M}{2\pi\kappa L}(1-\kappa r_0)\mathcal{I}$$

$$\boxed{B^C \;=\; \frac{\kappa_0^2}{M} \;-\; \frac{2\alpha\kappa_0}{1-\kappa_0 r_0}\left[\gamma_E \;+\; \log\left(\frac{\alpha M}{4\kappa_0}\right)\right] \;-\; \frac{\alpha}{\pi L}\mathcal{I} \;+\; \ldots}$$

✓ Two contributions in the *compact* limit:

$$\delta M_{BS}^{(FV)} \;=\; 2\delta M^{(FV)} - \delta B^C \;=\; 2\left(\frac{\alpha}{2\pi L}\mathcal{I}\right) + \frac{\alpha}{\pi L}\mathcal{I} \;+\; \ldots \;=\; \boxed{\frac{2\alpha}{\pi L}\mathcal{I}} + \;\ldots$$

2() + 

✓

Summary

- Electromagnetism essential for *ab initio* nuclear physics
- QED in a FV with PBCs is consistent without zero mode
- Two particle QC is tricky: violates Lüscher condition
- There is hierarchy of scales that allows perturbative treatment of Coulomb
- It would be very interesting to look at a simple system in lattice QCD.. $\pi^+\pi^+$?